arXiv:0912.2762v4 [math.AG] 29 Jul 2011

**Claude Sabbah**

---

# INTRODUCTION TO STOKES STRUCTURES

## LECTURE NOTES (LISBOA, JANUARY 2009)

---

*Claude Sabbah*
UMR 7640 du CNRS, Centre de Mathématiques Laurent Schwartz,
École polytechnique, F–91128 Palaiseau cedex, France.
*E-mail :* `sabbah@math.polytechnique.fr`
*Url :* `http://www.math.polytechnique.fr/~sabbah`

**2000 *Mathematics Subject Classification.*** — 34M40, 32C38, 35A27.

***Key words and phrases.*** — Meromorphic connection, real blowing-up, Riemann-Hilbert correspondence, Stokes filtration, Stokes-perverse sheaf.

This research was supported by the grant ANR-08-BLAN-0317-01 of the Agence nationale de la recherche.
*August 1, 2011*

# INTRODUCTION TO STOKES STRUCTURES

## LECTURE NOTES (LISBOA, JANUARY 2009)

**Claude Sabbah**

***Abstract.*** — The purpose of these lectures is to introduce the notion of a Stokes-perverse sheaf as a receptacle for the Riemann-Hilbert correspondence for holonomic $\mathscr{D}$-modules. They develop the original idea of P. Deligne in dimension one, and make it enter the frame of perverse sheaves. They also give a first step for a general definition in higher dimension, and make explicit particular cases of the Riemann-Hilbert correspondence, relying on recent results of T. Mochizuki.

# CONTENTS

**Introduction** 1

**1. $\mathcal{I}$-filtrations** 7
  1.a. Étale spaces of sheaves 7
  1.b. Étale spaces of sheaves of ordered abelian groups 8
  1.c. The category of pre-$\mathcal{I}$-filtrations 10
  1.d. Traces for étale maps 15
  1.e. Pre-$\mathcal{I}$-filtrations of sheaves 16
  1.f. $\mathcal{I}$-filtrations of sheaves (the Hausdorff case) 18
  1.g. $\mathcal{I}$-filtered local systems (the Hausdorff case) 20
  1.h. $\mathcal{I}$-filtered local systems (the stratified Hausdorff case) 20
  1.i. Comments 22

**Part I. Dimension one** 25

**2. Stokes-filtered local systems in dimension one** 27
  2.a. Non-ramified Stokes-filtered local systems 27
  2.b. Pull-back and push-forward 32
  2.c. Stokes filtrations on local systems 34
  2.d. Extension of scalars 37
  2.e. Stokes-filtered local systems and Stokes data 38

**3. Abelianity and strictness** 41
  3.a. Strictness and abelianity 41
  3.b. Level structure of a Stokes-filtered local system 42
  3.c. Proof of Theorem 3.5 44
  3.d. Proof of the stability by extension 49
  3.e. More on the level structure 49
  3.f. Comments 50

**4. Stokes-perverse sheaves on curves** 51
4.a. The setting 51
4.b. The category of Stokes-$\mathbb{C}$-constructible sheaves on $\widetilde{X}$ 52
4.c. Derived categories and duality 54
4.d. The category of Stokes-perverse sheaves on $\widetilde{X}$ 57
4.e. Direct image to $X$ 58
4.f. Stokes-perverse sheaves on $\underline{\widetilde{X}}$ 59
4.g. Associated perverse sheaf on $X$ 60

**5. The Riemann-Hilbert correspondence for holonomic $\mathscr{D}$-modules on curves** 63
5.a. Some basic sheaves 63
5.b. The Riemann-Hilbert correspondence for germs 65
5.c. The Riemann-Hilbert correspondence in the global case 71
5.d. Compatibility with duality for meromorphic connections 73

**6. Applications of the Riemann-Hilbert correspondence to holonomic distributions** 75
6.a. The Riemann-Hilbert correspondence for meromorphic connections of Hukuhara-Turrittin type 75
6.b. The Hermitian dual of a holonomic $\mathscr{D}_X$-module 76
6.c. Asymptotic expansions of holonomic distributions 77
6.d. Comments 83

**7. Riemann-Hilbert and Laplace on the affine line (the regular case)** 85
7.a. Introduction 85
7.b. Direct image of the moderate de Rham complex 87
7.c. Topological spaces 89
7.d. Topological Laplace transform 91
7.e. Proof of Theorem 7.6 and compatibility with Riemann-Hilbert 95
7.f. Compatibility of Laplace transformation with duality 97
7.g. Compatibility of topological Laplace transformation with Poincaré-Verdier duality 99
7.h. Comparison of both duality isomorphisms 102

**Part II. Dimension two and more** 105

**8. Real blow-up spaces and moderate de Rham complexes** 107
8.a. Real blow-up 107
8.b. The sheaf of functions with moderate growth on the real blow-up space 109
8.c. The moderate de Rham complex 112
8.d. Examples of moderate de Rham complexes 117

**9. Stokes-filtered local systems along a divisor with normal crossings** 121
9.a. The sheaf $\mathcal{I}$ on the real blow-up (smooth divisor case) 121
9.b. The sheaf $\mathcal{I}$ on the real blow-up (normal crossing case) 122

9.c. Goodness ........ 123
9.d. Stokes filtrations on local systems ........ 127
9.e. Behaviour by pull-back ........ 131
9.f. Partially regular Stokes-filtered local systems ........ 134

**10. The Riemann-Hilbert correspondence for good meromorphic connections (case of a smooth divisor)** ........ 137
10.a. Good formal structure of a meromorphic connection ........ 137
10.b. The Riemann-Hilbert functor ........ 138
10.c. Proof of the full faithfulness in Theorem 10.7 ........ 140
10.d. Elementary and graded equivalences ........ 141
10.e. Proof of the essential surjectivity in Theorem 10.7 ........ 143

**11. Good meromorphic connections** ........ 147
11.a. Preliminary notation ........ 147
11.b. Good formal decomposition ........ 148
11.c. Good lattices ........ 152
11.d. Proof of Theorem 11.16 ........ 154
11.e. The Hukuhara-Turrittin-Majima theorem ........ 156

**12. Leaky pipes** ........ 161
12.a. The setting ........ 161
12.b. Direct images and Riemann-Hilbert ........ 163

**13. The Riemann-Hilbert correspondence for good meromorphic connections** ........ 169
13.a. The Riemann-Hilbert functor ........ 169
13.b. Local theory ........ 170
13.c. Proof of the Malgrange-Sibuya theorem 13.4 ........ 173
13.d. Application to Hermitian duality of holonomic $\mathscr{D}$-modules ........ 175

**14. Nearby cycles of Stokes-filtered local systems** ........ 177
14.a. Preliminaries ........ 177
14.b. Adjunction ........ 178
14.c. Nearby cycles along a function (the good case) ........ 179
14.d. Nearby cycles along a function (dimension two) ........ 186
14.e. Moderate nearby cycles of holonomic $\mathscr{D}$-modules ........ 187
14.f. Comparison ........ 189

**Bibliography** ........ 193

**Index of notation** ........ 199

**Index** ........ 201

# INTRODUCTION

On a Riemann surface $X$, the Riemann-Hilbert correspondence for meromorphic connections with regular singularities on a discrete set $D$, or more generally for regular holonomic $\mathscr{D}$-modules with singularities at $D$, induces an equivalence of the corresponding category with the category of "monodromy data", which can be presented

- sheaf-theoretically as the category of locally constant sheaves of finite dimensional $\mathbb{C}$-vector spaces on $X^* = X \smallsetminus D$ (first case) or perverse sheaves with singularities at $D$ (second case),
- quiver-theoretically as the data of local monodromies and connection matrices (first case), together with canonical and variation morphisms (second case).

While the second presentation is suited to describing moduli spaces, the first one is suited to sheaf theoretic operations on such objects. Each of these objects can be defined over subfields $\boldsymbol{k}$ of $\mathbb{C}$, giving rise to a $\boldsymbol{k}$-structure on the meromorphic connection with regular singularities, or regular holonomic $\mathscr{D}$-module.

When the singularities are not regular any more, such a correspondence with both aspects also exists. The second one is the most popular, with Stokes data consisting of Stokes matrices instead of local monodromy data. The first one, initiated by P. Deligne [**15**] (case of meromorphic connections) and [**16**] (holonomic $\mathscr{D}$-modules), has also been developed by B. Malgrange [**46, 49**] and D. Babbitt & V.S. Varadarajan [**2**]. Moreover, the Poincaré duality has been expressed by integrals on "rapid decay cycles" by various authors [**30, 9**].

The purpose of these lectures is to develop the original idea of P. Deligne and B. Malgrange, and make it enter the frame of perverse sheaves, so that it can be extended to arbitrary dimensions. This has been motivated by recent beautiful results of T. Mochizuki [**65, 60**], who has rediscovered it and shown the powerfulness of this point of view in higher dimension.

One of the sought applications of this sheaf-theoretic approach, named Stokes-perverse sheaf, is to answer a question that S. Bloch asked me some years ago: to define a sheaf-theoretical Fourier transform over $\mathbb{Q}$ (say) taking into account the Stokes data. Note that the unpublished manuscript [**7**] gave an answer to this question (see also the recent work [**62**] of T. Mochizuki). The need of such an extension to dimension bigger than one also shows up in [**18**, p. 116](1).

---

(1) Deligne writes: "On aimerait dire (mais ceci nous obligerait à quitter la dimension 1)...".

One of the main problems in the "perverse" approach is to understand on which spaces the sheaves are to be defined. In dimension one, Deligne replaces first a Riemann surface by its real oriented blowing up at the singularities of the meromorphic connection, getting a surface with boundary, and endows the extended local system of horizontal sections with a "Stokes filtration" on the boundary. This is a filtration indexed by an ordered local system. We propose to regard such objects as sheaves on the étale space of the ordered local system (using the notion of étale space as in [**23**]). In order to obtain a perfect correspondence with holonomic $\mathscr{D}$-modules, Deligne fills the boundary with discs together with perverse sheaves on them, corresponding to the formal part of the meromorphic connection. The gluing at the boundary between the Stokes-filtered local system and the perverse sheaf is defined through grading the Stokes filtration.

The road is therefore a priori well-paved and the program can be clearly drafted:

(1) To define the notion of Stokes-constructible sheaf on a manifold, and a $t$-structure in its derived category, in order to recover the category of Stokes-perverse sheaves on a complex manifold as the heart of this $t$-structure.

(2) To exhibit a Riemann-Hilbert correspondence RH between holonomic $\mathscr{D}$-modules and Stokes-perverse sheaves, and to prove that it is an equivalence of categories.

(3) To define the direct image functor in the derived category of Stokes-constructible sheaves and prove the compatibility of RH when taking direct images of holonomic $\mathscr{D}$-modules.

An answer to the latter question would give a way to compute Stokes data of the asymptotic behaviour of integrals of multivalued functions which satisfy themselves a holonomic system of differential equations.

While we realize the first two points of the program in dimension one, by making a little more explicit the contents of [**15, 16**], we do not go to the end in dimension bigger than one, as we only treat the Stokes-perverse counterpart of meromorphic connections, not holonomic $\mathscr{D}$-modules. The reason is that some new phenomena appear, which were invisible in dimension one.

In order to make them visible, let us consider a complex manifold $X$ endowed with a divisor $D$. In dimension one, the topological space to be considered is the oriented real blow-up $\widetilde{X}$ of $X$ along $D$, and meromorphic connections on $X$ with poles on $D$ are in one-to-one correspondence with local systems on $X \smallsetminus D$ whose extension to $\widetilde{X}$ is equipped with a Stokes filtration along $\partial\widetilde{X}$. If $\dim X \geqslant 2$, in order to remain in the realm of local systems, a simplification of the underlying geometric situation seems unavoidable in general, so that we treat the case of a divisor with normal crossings (and all components smooth), and a generic assumption has also to be made on the connection, called *goodness*. Variants of this genericity condition have occurred in asymptotic analysis (e.g. in [**42**]) or when considering the extension of the Levelt-Turrittin theorem to many variables (e.g. in [**51**]). We define the notion of good stratified $\mathcal{I}$-covering of $\partial\widetilde{D}$. To any good meromorphic connection and to any good Stokes-filtered local system are associated in a natural way such a good stratified $\mathcal{I}$-covering, and the categories to be put into Riemann-Hilbert correspondence are those subcategories of objects having an associated stratified $\mathcal{I}$-covering contained in a fixed good one.

This approach remains non intrinsic, that is, while the category of meromorphic connections with poles along an arbitrary divisor is well defined, we are able to define a Stokes-topological counterpart only with the goodness property. This is an obstacle to define intrinsically a category of Stokes-perverse sheaves. This should be overcome together with the fact that such a category should be stable by direct images, as defined in Lecture 1 for pre-$\mathcal{I}$-filtrations.

Therefore, our approach still remains non-complete with respect to the program above, but already gives strong evidence of its feasibility.

Compared to the approach of T. Mochizuki in [**60, 65**], which is nicely surveyed in [**61**], we regard a Stokes-filtered object as an abstract “topological” object, while Mochizuki introduces the Stokes filtration as a filtration of a flat vector bundle. In the recent preprint [**62**], T. Mochizuki has developed the notion of a Betti structure on a holonomic $\mathscr{D}$-module and proved many functorial properties. Viewing the Betti structure as living inside a pre-existing object (a holonomic $\mathscr{D}$-module) makes it a little easier to analyze its functorial properties, since the functorial properties of holonomic $\mathscr{D}$-modules are already understood. On the other hand, this gives a strong evidence of the existence of a category of Stokes-perverse sheaves with good functorial properties.

### *Contents lecture by lecture*

Part one is mainly concerned with dimension one, although Lecture 7 anticipates some results in dimension two, according to the footnote on page 1.

In Lecture 1, we develop the notion of Stokes filtration in a general framework under the names of pre-$\mathcal{I}$-filtration and $\mathcal{I}$-filtration, with respect to an ordered sheaf of abelian group $\mathcal{I}$. The sheaf $\mathcal{I}$ for the Stokes filtration in dimension one consists of polar parts of multivalued meromorphic functions of one variable, as originally introduced by Deligne. Its étale space is Hausdorff, which makes the understanding of a filtration simpler with respect to taking the associated graded sheaf. This lecture may be skipped in a first reading.

In Lecture 2, we essentially redo more concretely the same work as in Lecture 1, in the context of Stokes-filtered local systems on a circle. We prove abelianity of the category in Lecture 3, a fact which follows from the Riemann-Hilbert correspondence, but is proved here directly over the base field $\boldsymbol{k}$. In doing so, we introduce the level structure, which was a basic tool in the higher dimensional analogue developed by

T. Mochizuki [**65**], and which was previously considered together with the notion of multisummability [**3, 55, 39**].

In Lectures 4 and 5 we develop the notion of a Stokes-perverse sheaf, mainly by following P. Deligne [**15, 16**] and B. Malgrange [**49**, Chap. IV.3], and prove the Riemann-Hilbert correspondence. We made explicit the behaviour with respect to duality, at least at the level of Stokes-filtered local systems, and the main tools are explained in Lecture 4.

Lecture 6 gives two analytic applications of the Riemann-Hilbert correspondence in dimension one. Firstly, the Hermitian dual of a holonomic $\mathscr{D}$-module (i.e., the conjugate module of the module of distribution solutions of the original one) on a Riemann surface is shown to be holonomic. Secondly, the local structure of distributions solutions of a holonomic system is analyzed.

Lecture 7 presents another application, with a hint of the theory in dimension two, by computing the Stokes filtration of the Laplace transform of a regular holonomic $\mathscr{D}$-module on the affine line. We introduce the topological Laplace transformation, and we make precise the relation with duality, of $\mathscr{D}$-modules on the one hand, Poincaré-Verdier on the other hand, and their relations. For this lecture, we use tools in dimension 2 which are fully developed in the next lecture.

Part two considers dimension $\geqslant 2$. Lecture 8 defines the real blow-up space along a family of divisors and the relation between various real blow-up spaces. We pay attention to the global existence of these spaces. The basic sheaf on such real blow-ups spaces is the sheaf of holomorphic functions with moderate growth along the divisor. It leads to the moderate de Rham complex of a meromorphic connection. We give some examples of such de Rham complexes, showing how non-goodness can produce higher dimensional cohomology sheaves.

Lecture 9 takes up Lectures 2 and 3 and introduces the goodness assumption. The construction of the sheaf $\mathcal{I}$ is given with some care, to make it global along the divisor.

The first approach to the Riemann-Hilbert correspondence in dimension $\geqslant 2$ is given in Lecture 10, along a smooth divisor. It can be regarded as obtained by putting a (good) parameter in Lecture 5. The main new argument is the local constancy of the *Stokes sheaf* (Stokes matrices can be chosen locally constant with respect to the parameter).

Lecture 11 analyzes the properties of good meromorphic connections, following T. Mochizuki [**65**], and gives a proof of the Hukuhara-Turrittin-Majima theorem in this context, which asymptotically lifts a formal decomposition of the connection. A short account of Mochizuki's proof has already been given by M. Hien in [**26**, Appendix].

Before considering the general case of the Riemann-Hilbert correspondence for meromorphic connections in Lecture 13, and answering a question of Kashiwara by proving that the Hermitian dual of a holonomic $\mathscr{D}$-module is holonomic (see Lecture 6 in dimension one), we make explicit in Lecture 12 a calculation of the Stokes filtration of an exponentially twisted Gauss-Manin system (such a system has already been analyzed by C. Roucairol [**69, 70, 71**]). However, the method is dependent on the simple geometric situation, so can hardly be extended directly to the general case.

Lastly, we give in Lecture 14 a definition of the nearby cycle functor relative to a holomorphic function, and compare it with the notion of moderate nearby cycles of a holonomic $\mathscr{D}$-module (through the theory of the $V$-filtration or some variants) through the Riemann-Hilbert correspondence. We restrict our study to the case of a meromorphic connection with poles along a divisor with normal crossings and a holomorphic function whose zero set is equal to this divisor.

*Acknowledgements.* — I thank A. Beilinson and the university of Chicago where part of this work was achieved, and T. Monteiro Fernandes and O. Neto in Lisbon (University of Lisbon, CAUL and CMAF) for giving me the opportunity to lecture on it in January 2009, as well as the audience of these lectures (whose content corresponds approximately to the present first six lectures) for many interesting questions and remarks. Many discussions with S. Bloch, H. Esnault, C. Hertling, M. Hien, T. Mochizuki and G. Morando have been very stimulating and helpful. I thank especially T. Mochizuki for letting me know his ongoing work on the subject, which strengthens the approach given here. Discussions with him have always been very enlightening. Needless to say, this work owes much to P. Deligne and B. Malgrange, and to D. Bertrand who asked me to help him when editing the volume [**19**], giving me the opportunity of being more familiar with this correspondence.

# LECTURE 1

# $\mathcal{I}$-FILTRATIONS

**Summary.** This lecture introduces the general framework for the study of the Stokes phenomenon in a sheaf-theoretic way. The underlying topological spaces are étale spaces of sheaves of ordered abelian groups $\mathcal{I}$. The general notion of pre-$\mathcal{I}$-filtration is introduced as a convenient abelian category to work in. The notion of $\mathcal{I}$-filtration is first considered when the étale space of $\mathcal{I}$ is Hausdorff. We will soon restrict to $\mathcal{I}$-filtrations of locally constant sheaves of $\boldsymbol{k}$-vector spaces and we will extend the definition to the case where $\mathcal{I}$ satisfies the stratified Hausdorff property. Most of the notions introduced in this lecture will be taken up as a more concrete approach to Stokes filtrations in Lecture 2, and this lecture may be skipped in a first reading. It contains nevertheless many guiding principles for Lectures 2, 3 and 9.

**1.a. Étale spaces of sheaves.** — In the following, $Y$ will be a locally compact and locally connected topological space (e.g. $Y = S^1$) and $\mathcal{I}$ a sheaf of abelian groups on $Y$. Let $\mu : \mathcal{I}^{\text{ét}} \to Y$ be the étale space associated to $\mathcal{I}$ (see [**23**, §II.1.2, p. 110]). Recall that, for any $y \in Y$, $\mu^{-1}(y) = \mathcal{I}_y$ with its discrete topology and any germ $\varphi_y \in \mathcal{I}_y$ has a fundamental system of open neighbourhoods consisting of the sets $\{\varphi_z \mid z \in U\}$, where $U$ is an open neighbourhood of $y$ in $Y$ on which the germ $\varphi_y$ extends as a section $\varphi \in \mathcal{I}(U)$. Then $\mu$ is a local homeomorphism, in particular it is an open map and $\mathcal{I}^{\text{ét}}$ is locally connected (because $Y$ is so).

At some places in this lecture, we will assume that $\mathcal{I}^{\text{ét}}$ is Hausdorff. This means that, for any open set $U \subset Y$ and any $\varphi, \psi \in \Gamma(U, \mathcal{I})$, if $\varphi_y \neq \psi_y$ for some $y \in U$, then $\varphi_z \neq \psi_z$ for any $z$ in some neighbourhood of $y$, that is, the maximal open subset $U(\varphi, \psi)$ on which $\varphi \equiv \psi$ satisfies $\overline{U(\varphi, \psi)} \not\ni y$. This is equivalent to saying that $U(\varphi, \psi)$ is also closed, that is, is empty or a connected component of $U$. In such a case, $\mathcal{I}^{\text{ét}}$ is then locally compact.

***Examples 1.1***

(1) If $Y$ is locally connected and $\mathcal{I}$ is a locally constant sheaf, then $\mathcal{I}^{\text{ét}}$ is Hausdorff.

(2) If $\mathcal{J}$ is a subsheaf of $\mathcal{I}$ and if $\mathcal{I}^{\text{ét}}$ is Hausdorff, then so is $\mathcal{J}^{\text{ét}}$ (as $\mathcal{J}^{\text{ét}}$ is open in $\mathcal{I}^{\text{ét}}$).

(3) Let $X$ be a complex manifold and let $D$ be a reduced divisor in $X$. Let $\mathscr{O}_X(*D)$ be the sheaf of meromorphic functions on $X$ with poles along $D$ at most. Then if $D$ is locally irreducible, the sheaf $\mathcal{I}=\mathscr{O}_X(*D)/\mathscr{O}_X$, regarded as a sheaf on $D$, has a Hausdorff étale space: indeed, if a germ $\varphi_x\in\mathscr{O}_{X,x}(*D)$ $(x\in D)$, defined on some open set $U\subset X$ such that $(D\smallsetminus\mathrm{Sing}(D))\cap U$ is connected, is holomorphic on some nonempty open set of $D\cap U$, it is holomorphic on $(D\smallsetminus\mathrm{Sing}(D))\cap U$, and thus all over $D\cap U$ by Hartogs.

(4) If $f:Y'\to Y$ is continuous and $Y'$ is Hausdorff, then $f^{-1}\mathcal{I}$ satisfies the Hausdorff property when $\mathcal{I}$ does so. If moreover $f$ is proper, then the converse holds, that is, $\mathcal{I}$ satisfies the Hausdorff property when $f^{-1}\mathcal{I}$ does so.

(5) Let $i:Y\hookrightarrow Y'$ be a closed immersion (with $Y'$ Hausdorff). If the sheaf $\mathcal{I}$ on $Y$ satisfies the Hausdorff property, then in general $i_*\mathcal{I}$ on $Y'$ does not.

***Example 1.2*** **(Geometry in étale spaces).** — Let $Y$ be a complex analytic manifold and let $\mathcal{F}$ be a locally free $\mathscr{O}_Y$-module of finite rank. Let $p:F\to Y$ be the associated holomorphic bundle. If $\mu:\mathcal{F}^{\text{ét}}\to Y$ denotes the étale space of $\mathcal{F}$, then there is a natural commutative diagram

$$\begin{tikzcd}
\mathcal{F}^{\text{ét}} \arrow[rr, "\mathrm{ev}"] \arrow[dr, "\mu"'] & & F \arrow[dl, "p"] \\
& Y &
\end{tikzcd}$$

where the evaluation map ev associates to any germ $s_y$ of holomorphic section of $p:F\to Y$ at $y$ its value $s_y(y)$. The map ev is continuous.

We will also deal with the following situation. Assume that we are given a closed subset $\Sigma$ of $\mathcal{F}^{\text{ét}}$ such that $\mu:\Sigma\to Y$ is a finite covering. Then $\Sigma$ is naturally equipped with the structure of a complex manifold. The image $\Sigma'=\mathrm{ev}(\Sigma)$ is an analytic subset of $F$, locally equal to the finite union of local holomorphic sections of $p:F\to Y$, the map $p:\Sigma'\to Y$ is finite and the map $\mathrm{ev}:\Sigma\to\Sigma'$ is the normalization.

### 1.b. Étale spaces of sheaves of ordered abelian groups.

— Assume now that $\mathcal{I}$ is a sheaf of ordered abelian groups, i.e., a sheaf with values in the category of ordered abelian groups. Hence, for every open set $U$ of $Y$, $\mathcal{I}(U)$ is an ordered abelian group, with order denoted by $\leqslant_U$, and the restriction maps $(\mathcal{I}(U),\leqslant_U)\to(\mathcal{I}(V),\leqslant_V)$, for $V\subset U$, are morphisms of ordered sets. For every $y\in Y$, the germ $\mathcal{I}_y$ is then ordered: for $\varphi,\psi\in\mathcal{I}_y$, we have $\varphi\leqslant_y\psi$ iff there exists an open neighbourhood $U\ni y$ such that $\varphi,\psi\in\mathcal{I}(U)$ and $\varphi\leqslant_U\psi$. For every open set $V\subset U$ we then have $\varphi_{|V}\leqslant_V\psi_{|V}$.

For $\varphi,\psi\in\Gamma(U,\mathcal{I})$, we then have $\varphi\leqslant_U\psi$ iff $\varphi-\psi\leqslant_U 0$. Giving an order compatible with addition on a sheaf of abelian groups is equivalent to giving a subsheaf $\mathcal{I}_{\leqslant 0}\subset\mathcal{I}$ of abelian groups such that $-\mathcal{I}_{\leqslant 0}\cap\mathcal{I}_{\leqslant 0}=0$. Then, for $\varphi,\psi\in\Gamma(U,\mathcal{I})$, we have $\varphi\leqslant_U\psi$ iff $\varphi-\psi\in\Gamma(U,\mathcal{I}_{\leqslant 0})$.

We also set $<_{U} = (\leqslant_{U}$ and $\neq)$. For every $y \in Y$ and for $\varphi, \psi \in \mathcal{I}_y$, we have $\varphi <_{y} \psi$ iff for any sufficiently small open neighbourhood $V \ni y$ such that $\varphi, \psi \in \mathcal{I}(V)$, $\varphi_{|V} <_{V} \psi_{|V}$, that is, $\varphi_{|V} \leqslant_{V} \psi_{|V}$ and $\varphi_{|V} \neq \psi_{|V}$.

***Definition 1.3* (Exhaustivity).** — We say that a sheaf $\mathcal{I}$ satisfies the *exhaustivity property* if, for any $y \in Y$ and any finite set $\Phi_y \subset \mathcal{I}_y$, there exists $\varphi_y, \psi_y \in \mathcal{I}_y$ such that $\psi_y <_{y} \Phi_y$ and $\Phi_y \leqslant \varphi_y$.

***Example 1.4*.** — Let us already remark that the constant sheaf $\mathcal{I}$ of Example 1.6 below satisfies the exhaustivity property: this is seen by choosing $\psi_\theta$ and $\varphi_\theta$ with a pole of order bigger than the orders of the pole of the elements of $\Phi_\theta$, and suitable dominant coefficients.

***Notation 1.5*.** — Let $U$ be an open subset of $Y$ and let $\varphi, \psi \in \Gamma(U, \mathcal{I})$. We denote by $U_{\psi \leqslant \varphi}$ the subset of $U$ defined by $y \in U_{\psi \leqslant \varphi}$ iff $\psi_y \leqslant_{y} \varphi_y$. Then $U_{\psi \leqslant \varphi}$ is open in $U$ (hence in $Y$). The boundary of $U_{\psi \leqslant \varphi}$ is denoted by $\mathrm{St}(\psi, \varphi)$, with the convention that $\mathrm{St}(\varphi, \varphi) = \varnothing$.

Let $(\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}})_{\leqslant} \subset \mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}}$ (resp. $(\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}})_{<}$) be the subset consisting of those $(\varphi, \psi) \in \mathcal{I}_y \times \mathcal{I}_y$ such that $\varphi \leqslant_{y} \psi$ (resp. $\varphi <_{y} \psi$) $(y \in Y)$. These are open subsets (with the Hausdorff assumption in the $<$ case). We denote by $j_{\leqslant}, j_{<}$ the corresponding inclusions. Let us note that the diagonal inclusion $\delta : \mathcal{I}^{\text{ét}} \hookrightarrow \mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}}$ is open, and also closed if we assume that $\mathcal{I}^{\text{ét}}$ is Hausdorff, and $(\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}})_{\leqslant} = (\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}})_{<} \cup \delta(\mathcal{I}^{\text{ét}})$. The two projections $p_1, p_2 : \mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}} \to \mathcal{I}^{\text{ét}}$ also are local homeomorphisms (both are the étale space corresponding to the sheaf $\mu^{-1}\mathcal{I}$).

We regard $U_{\psi \leqslant \varphi}$ as the pull-back of $(\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}})_{\leqslant}$ by the section $(\varphi, \psi) : U \to \mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}}$. Similarly, $U_{\psi < \varphi}$ denotes the subset defined by $y \in U_{\psi < \varphi}$ iff $\psi_y <_{y} \varphi_y$. If we assume that $\mathcal{I}^{\text{ét}}$ is Hausdorff, $U_{\psi < \varphi}$ is open in $U$ (because its complementary set in $U_{\psi \leqslant \varphi}$, which is the pull-back by $(\varphi, \psi)$ of the diagonal, is then closed in $U_{\psi \leqslant \varphi}$). If $U$ is connected, then either $\varphi \equiv \psi$ on $U$ and $U_{\psi < \varphi} = \varnothing$, or $\psi < \varphi$ everywhere on $U$ and $U_{\psi < \varphi} = U_{\psi \leqslant \varphi}$.

***Example 1.6* (Main example in dimension one).** — Let $X$ be the open disc centered at 0 and radius 1 in $\mathbb{C}$, with coordinate $x$. Let $X^*$ be the punctured disc $X \smallsetminus \{0\}$ and let us denote by $\varpi : \widetilde{X} \to X$ the real oriented blowing-up of $X$ at the origin, so that $\widetilde{X} = S^1 \times [0, 1)$. In the following, we denote by $S^1$ the boundary $S^1 \times \{0\}$ of $X$ and we forget about $X$. We still denote by $\varpi$ the constant map $S^1 \to \{0\}$.

Set $\mathscr{O} = \mathbb{C}\{x\}$ and let $\mathcal{I}$ be the constant sheaf $\varpi^{-1}\mathcal{P}$ on $S^1$, with $\mathcal{P} = \mathscr{O}(*0)/\mathscr{O}$ (polar parts of Laurent expansions). For any connected open set $U$ of $S^1$ and any $\varphi, \psi \in \mathcal{P} = \Gamma(U, \mathcal{I})$, we define $\psi \leqslant_{U} \varphi$ if $e^{\psi - \varphi}$ has *moderate growth* on a neighbourhood of $U$ in $S^1 \times [0, 1)$, that is, for any compact set $K \in U$, there exists $C_K > 0$ and $N_K \in \mathbb{N}$ such that, for some open neighbourhood $\mathrm{nb}(K)$ of $K$ in $S^1 \times [0, 1)$ and for some representative of $\varphi, \psi$ in $\mathscr{O}(*0)$, the inequality $|e^{\psi - \varphi}| \leqslant C_K |x|^{N_K}$ holds on $\mathrm{nb}(K) \cap S^1 \times (0, 1)$ (this can also be expressed in polar coordinates $x = re^{i\theta}$).

Let us check that this is indeed an order on $\Gamma(U,\mathcal{I})$. The only point to check, due to the additivity property, is that, for $\eta\in\mathcal{P}$, $\eta\leqslant_U 0$ and $0\leqslant_U\eta$ imply $\eta=0$. If $\eta\neq 0$, let us write $\eta=u_n(x)x^{-n}$ with $n\geqslant 1$ and $u_n(0)\neq 0$. For a given $\theta\in U$, we have

$$\eta\leqslant_\theta 0\Longleftrightarrow \eta=0 \text{ or } \arg u_n(0)-n\theta\in(\pi/2,3\pi/2)\mod 2\pi. \tag{1.6 $*$}$$

It clearly follows that the relation $\leqslant_\theta$ is an order relation on $\mathcal{I}_\theta$, and since for $\eta\in\Gamma(U,\mathcal{I})$ we have $\eta\leqslant_U 0$ if and only if $\eta\leqslant_\theta 0$ for any $\theta\in U$, the relation $\leqslant_U$ is an order relation on $\Gamma(U,\mathcal{I})$.

It clearly follows from (1.6 $*$) that $\psi\leqslant_\theta\varphi$ iff $\mathrm{Re}(\psi-\varphi)\leqslant 0$ in some neighbourhood of $(\theta,0)\in S^1\times[0,1)$. We also have

$$\eta<_\theta 0\Longleftrightarrow \eta\neq 0 \text{ and } \arg u_n(0)-n\theta\in(\pi/2,3\pi/2)\mod 2\pi. \tag{1.6 $**$}$$

Given $\varphi\neq\psi\in\mathcal{P}$, there exists at most a finite subset of $S^1$ where neither $\psi<_\theta\varphi$ nor $\varphi<_\theta\psi$: if $\varphi-\psi=u_n(x)x^{-n}$ as above, it consists of the $\theta$ such that $\arg u_n(0)-n\theta=\pm\pi/2\mod 2\pi$. It is called the set of *Stokes directions* of $(\varphi,\psi)$. It is the boundary $\mathrm{St}(\varphi,\psi)$ of $S^1_{\psi\leqslant\varphi}$

**1.c. The category of pre-$\mathcal{I}$-filtrations.** — This category will be the ambient category where objects are defined. Its main advantage is to be abelian. We assume that $\mathcal{I}$ is as in §1.b. Recall that we have a commutative diagram of étale maps

$$\begin{tikzcd} \mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}} \arrow[r, "p_1"] \arrow[d, "p_2"'] & \mathcal{I}^{\text{ét}} \arrow[d, "\mu"] \\ \mathcal{I}^{\text{ét}} \arrow[r, "\mu"] & Y \end{tikzcd} \tag{1.7}$$

and an open subset $(\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}})_\leqslant\subset\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}}$. We denote by $\beta_\leqslant$ the functor on the category of sheaves on $\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}}$ which restricts to $(\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}})_\leqslant$ and extends by zero. In [**35**, Prop. 2.3.6], it is denoted by putting $(\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}})_\leqslant$ as an index. This is an exact functor, and there is a natural morphism of functors $\beta_\leqslant\to\mathrm{Id}$.

Let $\mathscr{F}$ be a sheaf of $\boldsymbol{k}$-vector spaces on $Y$. We then have $p_1^{-1}\mu^{-1}\mathscr{F}=p_2^{-1}\mu^{-1}\mathscr{F}$ and therefore a functorial morphism $\beta_\leqslant p_1^{-1}\mu^{-1}\mathscr{F}\to p_2^{-1}\mu^{-1}\mathscr{F}$. We call $\mu^{-1}\mathscr{F}$ (together with this morphism) the *constant pre-$\mathcal{I}$-filtration on $\mathscr{F}$*.

***Definition 1.8* (Pre-$\mathcal{I}$-filtration)**

(1) By a *pre-$\mathcal{I}$-filtration* we will mean a sheaf $\mathscr{F}_\leqslant$ of $\boldsymbol{k}$-vector spaces on $\mathcal{I}^{\text{ét}}$ equipped with a morphism $\beta_\leqslant p_1^{-1}\mathscr{F}_\leqslant\to p_2^{-1}\mathscr{F}_\leqslant$ such that, when restricted to the diagonal, $\beta_\leqslant p_1^{-1}\mathscr{F}_\leqslant\to p_2^{-1}\mathscr{F}_\leqslant$ is the identity.

(2) A morphism $\lambda:\mathscr{F}_\leqslant\to\mathscr{F}'_\leqslant$ of pre-$\mathcal{I}$-filtrations is a morphism of the corresponding sheaves, with the obvious compatibility relation: the following diagram

should commute:

$$\begin{tikzcd}
\beta_{\leqslant} p_1^{-1}\mathscr{F}_{\leqslant} \arrow[r] \arrow[d, "\beta_{\leqslant} p_1^{-1}\lambda"'] & p_2^{-1}\mathscr{F}_{\leqslant} \arrow[d, "p_2^{-1}\lambda"] \\
\beta_{\leqslant} p_1^{-1}\mathscr{F}'_{\leqslant} \arrow[r] & p_2^{-1}\mathscr{F}'_{\leqslant}
\end{tikzcd}$$

We therefore define in this way a category, denoted by $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. It comes with a "forgetting" functor to the category $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$ of sheaves of $\boldsymbol{k}$-vector spaces on $\mathcal{I}^{\text{ét}}$. We will usually denote an object of both categories by $\mathscr{F}_{\leqslant}$, hoping that this does not lead to any confusion.

Given a sheaf $\mathscr{F}_{\leqslant}$ on $\mathcal{I}^{\text{ét}}$, we denote its germ at $\varphi_y \in \mathcal{I}_y$ by $\mathscr{F}_{\leqslant\varphi_y}$. The supplementary data of $\beta_{\leqslant} p_1^{-1}\mathscr{F}_{\leqslant} \to p_2^{-1}\mathscr{F}_{\leqslant}$ induces, for each pair of germs $\varphi_y, \psi_y \in \mathcal{I}_y$ such that $\varphi_y \leqslant_y \psi_y$ a morphism $\mathscr{F}_{\leqslant\varphi_y} \to \mathscr{F}_{\leqslant\psi_y}$, justifying the name of "filtration", although we do not impose injectivity in order to get an abelian category. The condition on morphisms is that they are compatible with the "filtration". Given a section $\varphi \in \mathcal{I}(U)$ on some open set $U \subset Y$, we also denote by $\mathscr{F}_{\leqslant\varphi}$ the pull-back $\varphi^{-1}\mathscr{F}_{\leqslant}$ on $U$. This is a sheaf on $U$, whose germ at $y \in U$ is the germ $\mathscr{F}_{\leqslant\varphi_y}$ of $\mathscr{F}_{\leqslant}$ at $\varphi_y \in \mathcal{I}^{\text{ét}}$.

***Remark 1.9.*** — The space $\mathcal{I}^{\text{ét}}$ has an open covering formed by the $\varphi(U)$, where $U$ runs among open subsets of $Y$ and, for each $U$, $\varphi$ belongs to $\Gamma(U, \mathcal{I})$. Giving a sheaf on $\mathcal{I}^{\text{ét}}$ is then equivalent to giving a sheaf $\mathscr{F}_{\leqslant\varphi}$ on $U$ for each pair $(U, \varphi)$, together with compatible isomorphisms $(\mathscr{F}_{\leqslant\varphi})_{|V} \xrightarrow{\sim} \mathscr{F}_{\leqslant(\varphi_{|V})}$ for $V \subset U$. The supplementary data of a morphism $\beta_{\leqslant} p_1^{-1}\mathscr{F}_{\leqslant} \to p_2^{-1}\mathscr{F}_{\leqslant}$ is equivalent to giving, for each pair $\varphi, \psi \in \Gamma(U, \mathcal{I})$, of a morphism $\beta_{\varphi\leqslant\psi}\mathscr{F}_{\leqslant\varphi} \to \mathscr{F}_{\leqslant\psi}$, and such morphisms should commute with the isomorphisms above for each $V \subset U$.

***Definition 1.10* (Twist).** — Let $\mathscr{F}_{\leqslant}$ be a pre-$\mathcal{I}$-filtration and let $\eta \in \Gamma(Y, \mathcal{I})$ be a global section of $\mathcal{I}$. The *twisted pre-$\mathcal{I}$-filtration* $\mathscr{F}[\eta]_{\leqslant}$ is defined by $\mathscr{F}[\eta]_{\leqslant\varphi} = \mathscr{F}_{\leqslant\varphi-\eta}$ for any local section $\varphi$ of $\mathcal{I}$. It is a pre-$\mathcal{I}$-filtration. A morphism $\lambda : \mathscr{F}_{\leqslant} \to \mathscr{G}_{\leqslant}$ induces a morphism $\mathscr{F}[\eta]_{\leqslant} \to \mathscr{G}[\eta]_{\leqslant}$, so that $[\eta]$ is a functor from $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ to itself, with inverse functor $[-\eta]$.

***Lemma 1.11.*** — *The category* $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *of pre-$\mathcal{I}$-filtrations is abelian and a sequence is exact if and only if the sequence of the underlying sheaves is exact.*

*Proof.* — This follows from the fact that $\beta_{\leqslant} p_1^{-1}$ and $p_2^{-1}$ are exact functors. ☐

***Remark 1.12* (Tensor product).** — We will not try to define at this level of generality an internal tensor product of the category $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. However, we can easily define a tensor product operation $\mathrm{Mod}(\boldsymbol{k}_Y) \times \mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \to \mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. To $\mathscr{F}'$ in $\mathrm{Mod}(\boldsymbol{k}_Y)$ and $\mathscr{F}_{\leqslant}$ in $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, we associate $\mu^{-1}\mathscr{F}' \otimes_{\boldsymbol{k}} \mathscr{F}_{\leqslant}$, where $\mu^{-1}\mathscr{F}'$ is the constant pre-$\mathcal{I}$-filtration on $\mathscr{F}'$. The associated morphism is obtained as

$$\begin{aligned}\beta_{\leqslant}p_1^{-1}(\mu^{-1}\mathscr{F}'\otimes_{\boldsymbol{k}}\mathscr{F}_{\leqslant}) &= p_1^{-1}\mu^{-1}\mathscr{F}'\otimes_{\boldsymbol{k}}\beta_{\leqslant}p_1^{-1}\mathscr{F}_{\leqslant}\\ &\longrightarrow p_1^{-1}\mu^{-1}\mathscr{F}'\otimes_{\boldsymbol{k}}p_2^{-1}\mathscr{F}_{\leqslant} = p_2^{-1}\mu^{-1}\mathscr{F}'\otimes_{\boldsymbol{k}}p_2^{-1}\mathscr{F}_{\leqslant} = p_2^{-1}(\mu^{-1}\mathscr{F}'\otimes_{\boldsymbol{k}}\mathscr{F}_{\leqslant}).\end{aligned}$$

*Push-forward.* — Our purpose is now to define the *push-forward of a pre-$\mathcal{I}$-filtration* by a continuous map $f : Y'\to Y$. We first consider the case of a cartesian diagram

$$\begin{tikzcd} \widetilde{\mathcal{I}^{\text{ét}}} \arrow[r, "\widetilde{f}"] \arrow[d, "\widetilde{\mu}"'] & \mathcal{I}^{\text{ét}} \arrow[d, "\mu"] \\ Y' \arrow[r, "f"] & Y \end{tikzcd} \tag{1.13}$$

with $\widetilde{\mathcal{I}} := f^{-1}\mathcal{I}$ equipped with the pull-back order, so that $(\widetilde{\mathcal{I}^{\text{ét}}}\times_{Y'}\widetilde{\mathcal{I}^{\text{ét}}})_{\leqslant}$ is the pull-back of the open set $(\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}})_{\leqslant}\subset\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}}$ by $\widetilde{f}\times\widetilde{f}$. As a consequence,

$$\begin{aligned}(\widetilde{f}\times\widetilde{f})_*\circ\widetilde{\beta}_{\leqslant} &= \beta_{\leqslant}\circ(\widetilde{f}\times\widetilde{f})_*\\ (\widetilde{f}\times\widetilde{f})^{-1}\circ\beta_{\leqslant} &= \widetilde{\beta}_{\leqslant}\circ(\widetilde{f}\times\widetilde{f})^{-1}.\end{aligned}\tag{1.14}$$

***Lemma 1.15.*** — *There are natural functors*

$$\begin{aligned}\widetilde{f}_* &: \operatorname{Mod}(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant})\longrightarrow\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})\\ \widetilde{f}^{-1} &: \operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})\longrightarrow\operatorname{Mod}(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant})\end{aligned}$$

*and natural morphisms of functors*

$$\begin{aligned}\widetilde{f}^{-1}\widetilde{f}_*\longrightarrow\operatorname{Id}\quad &\textit{in}\ \operatorname{Mod}(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant})\\ \operatorname{Id}\longrightarrow\widetilde{f}_*\widetilde{f}^{-1}\quad &\textit{in}\ \operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}).\end{aligned}$$

*Moreover,*

$$\operatorname{Hom}_{\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}}(\mathscr{F}_{\leqslant},\widetilde{f}_*\widetilde{\mathscr{G}}_{\leqslant})\simeq\operatorname{Hom}_{\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant}}(\widetilde{f}^{-1}\mathscr{F}_{\leqslant},\widetilde{\mathscr{G}}_{\leqslant}).$$

*Proof.* — At the level of sheaves the result is standard. One has only to check the compatibility with the morphisms $\beta_{\leqslant}p_1^{-1}\to p_2^{-1}$ and $\widetilde{\beta}_{\leqslant}\widetilde{p}_1^{-1}\to\widetilde{p}_2^{-1}$. It follows from (1.14). For instance, the adjunction morphisms are obtained from the similar ones for $\widetilde{f}\times\widetilde{f}$. □

***Corollary 1.16.*** — *The category* $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *has enough injectives.*

*Proof.* — Let $Y_{\text{disc}}$ be $Y$ equipped with its discrete topology and $\delta : Y_{\text{disc}}\to Y$ be the canonical morphism. Let us consider the diagram (1.13) for $\delta$. Since $\widetilde{\mu}$ is a local homeomorphism, $\widetilde{\mathcal{I}^{\text{ét}}}$ is nothing but $\mathcal{I}^{\text{ét}}_{\text{disc}}$ and $\widetilde{\delta}$ is the corresponding canonical morphism. Using the adjunction for $\widetilde{\delta}$, one shows as in [**35**, Prop. 2.4.3] that it is enough to check the corollary for $Y_{\text{disc}}$, that is, stalkwise for $\mathcal{I}^{\text{ét}}$. Assume then that we are given vector spaces $\mathscr{F}_{\leqslant\varphi_y}$ for any $\varphi_y\in\mathcal{I}^{\text{ét}}$, together with morphisms $\mathscr{F}_{\leqslant\varphi_y}\to\mathscr{F}_{\leqslant\psi_y}$ whenever $\varphi_y\leqslant_y\psi_y$. Choose a monomorphism $\mathscr{F}_{\leqslant\varphi_y}\hookrightarrow\mathscr{I}_{\leqslant\varphi_y}$

into an injective object in the category of $\boldsymbol{k}$-vector spaces, for any $\varphi_y \in \mathcal{I}^{\text{ét}}$. By injectivity of $\mathscr{I}_{\leqslant \psi_y}$, the composed morphism $\mathscr{F}_{\leqslant \varphi_y} \to \mathscr{I}_{\leqslant \psi_y}$ extends to a morphism $\mathscr{I}_{\leqslant \varphi_y} \to \mathscr{I}_{\leqslant \psi_y}$, providing a monomorphism $\mathscr{F}_{\leqslant} \hookrightarrow \mathscr{I}_{\leqslant}$ in $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}_{\text{disc}},\leqslant})$. ☐

Arguing as in [**35**, Prop. 2.4.6(vii)], one checks that injectives are flabby (i.e., the underlying sheaf is flabby). Going back to the setting of Lemma 1.15, we deduce:

***Corollary 1.17.*** — *The subcategory of* $\operatorname{Mod}(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant})$ *whose objects have a flabby underlying sheaf is injective with respect to* $\widetilde{f}_*$ *(see* [**35**, Def. 1.8.2]*).* ☐

Notice also that the construction of Godement ([**23**, §II.4.3]) can be extended to the present setting: using the notation of the proof of Corollary 1.16, the adjunction $\mathrm{Id} \to \widetilde{\delta}_*\widetilde{\delta}^{-1}$ gives an injective morphism $\widetilde{\mathscr{F}_{\leqslant}} \to \widetilde{\delta}_*\widetilde{\delta}^{-1}\widetilde{\mathscr{F}_{\leqslant}}$ and the latter has a flabby underlying sheaf.

We will denote by $D^+(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ (resp. $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$) the left-bounded (resp. bounded) derived category of $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. We have "forgetting" functors to $D^+(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$, resp. $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$.

***Remark 1.18* (Tensor product).** — According to Remark 1.12, we have a tensor product bifuntor $D^b(\boldsymbol{k}_Y) \times D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \to D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$.

***Corollary 1.19.*** — *The derived functor* $\boldsymbol{R}\widetilde{f}_* : D^+(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant}) \to D^+(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *is well defined and compatible with the similar functor* $D^+(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}}}) \to D^+(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$*, and if $f$ has finite cohomological dimension, so does* $\boldsymbol{R}\widetilde{f}_* : D^b(\boldsymbol{k}_{\widetilde{\mathcal{I}^{\text{ét}}},\leqslant}) \to D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. ☐

In general, given $\mathcal{I}'$ on $Y'$, $\mathcal{I}$ on $Y$ and $f : Y' \to Y$, we wish to define the pushforward of an object of $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}'^{\text{ét}},\leqslant})$ as an object of $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, and similarly at the level of derived categories. In order to do so, we need to assume the existence of a morphism $q_f : \widetilde{\mathcal{I}} := f^{-1}\mathcal{I} \to \mathcal{I}'$. We now consider a diagram

$$\begin{tikzcd}
\mathcal{I}'^{\text{ét}} \arrow[d, "\mu'"'] & \widetilde{\mathcal{I}^{\text{ét}}} \arrow[l, "q_f"'] \arrow[r, "\widetilde{f}"] \arrow[dl, "\widetilde{\mu}"] & \mathcal{I}^{\text{ét}} \arrow[d, "\mu"] \\
Y' \arrow[rr, "f"] & & Y
\end{tikzcd} \tag{1.20}$$

and we will also make the natural assumption that $q_f$ is compatible with the order, that is, given any two germs $\varphi_y, \psi_y \in \mathcal{I}_y$ ($y \in Y$), defining germs $\varphi_y, \psi_y \in (\widetilde{f}^{-1}\mathcal{I})_{y'}$ at $y' \in f^{-1}(y)$, we have $\varphi_y \leqslant_y \psi_y \Rightarrow q_f(\varphi_y) \leqslant_{y'} q_f(\psi_y)$ for any such $y'$. This is equivalent to asking that $(\widetilde{\mathcal{I}^{\text{ét}}} \times_{Y'} \widetilde{\mathcal{I}^{\text{ét}}})_{\leqslant}$ is contained in the pull-back of $(\mathcal{I}'^{\text{ét}} \times_{Y'} \mathcal{I}'^{\text{ét}})_{\leqslant}$ by $q_f \times q_f$. Under this assumption, we get a morphism of functors

$$\widetilde{\beta}_{\leqslant}(q_f \times q_f)^{-1} \longrightarrow (q_f \times q_f)^{-1}\beta'_{\leqslant} \tag{1.21}$$

Therefore, for any object $\mathscr{F}'_{\leqslant}$ of $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}'^{\text{ét}},\leqslant})$ with corresponding morphism $\beta'_{\leqslant}p_1'^{-1}\mathscr{F}'_{\leqslant} \to p_2'^{-1}\mathscr{F}'_{\leqslant}$, $q_f^{-1}\mathscr{F}'_{\leqslant}$ comes equipped with a morphism $\widetilde{\beta}_{\leqslant}\widetilde{p}_1^{-1}q_f^{-1}\mathscr{F}'_{\leqslant} \to$

$\widetilde{p}_2^{-1}q_f^{-1}\mathscr{F}'_{\leqslant}$, according to (1.21), and $\widetilde{f}_*q_f^{-1}\mathscr{F}'_{\leqslant}$ with a morphism $\beta_{\leqslant}p_1^{-1}\widetilde{f}_*q_f^{-1}\mathscr{F}'_{\leqslant} \to p_2^{-1}\widetilde{f}_*q_f^{-1}\mathscr{F}'_{\leqslant}$, according to (1.14).

***Definition 1.22.*** — Under these assumptions, the push-forward by $f$ is defined by $\widetilde{f}_*q_f^{-1}$.

The functor $q_f^{-1}$ is exact, while $\widetilde{f}_*$ is a priori only left exact. We will now extend the push-forward functor to the derived categories. For the sake of simplicity, we assume that $f$ has finite cohomological dimension.

***Definition 1.23* (Push-forward).** — Under these assumptions, the push-forward functor $f_+ : D^b(\boldsymbol{k}_{\mathcal{I}'^{\text{ét}},\leqslant}) \to D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ is defined as the composed functor $\boldsymbol{R}f_*q_f^{-1}$.

*The functor $\mathscr{H}om(\iota^{-1}\bullet,\mu^{-1}\mathscr{G})$.* — We denote by $\iota$ the involution $\varphi \mapsto -\varphi$ on $\mathcal{I}^{\text{ét}}$ and by $\tau$ the permutation $(\varphi,\psi) \mapsto (\psi,\varphi)$ on $\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}}$. We have $p_2 = p_1 \circ \tau$. Let $\mathscr{G}$ be a sheaf on $Y$. Recall that $\mu^{-1}\mathscr{G}$ is the constant pre-$\mathcal{I}$-filtration on $\mathscr{G}$. We will define a contravariant functor $\mathscr{H}om(\iota^{-1}\bullet,\mu^{-1}\mathscr{G})$ from $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ to itself, over the corresponding functor on $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$.

Given an object $\mathscr{F}_{\leqslant}$ of $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, we apply $\sigma^{-1}$ to the corresponding morphism $\beta_{\leqslant}p_1^{-1}\mathscr{F}_{\leqslant} \to p_2^{-1}\mathscr{F}_{\leqslant}$ to get a morphism $\beta_{\geqslant}p_2^{-1}\mathscr{F}_{\leqslant} \to p_1^{-1}\mathscr{F}_{\leqslant}$ (with obvious notation $\beta_{\geqslant}$). Applying then $(\iota \times_Y \iota)^{-1}$, we get a morphism $\beta_{\leqslant}p_2^{-1}\iota^{-1}\mathscr{F}_{\leqslant} \to p_1^{-1}\iota^{-1}\mathscr{F}_{\leqslant}$. Since $p_2^{-1}\mu^{-1}\mathscr{G} = p_1^{-1}\mu^{-1}\mathscr{G}$, we obtain, since $p_1$ and $p_2$ are local homeomorphisms, a morphism

$$\begin{aligned}p_1^{-1}\,\mathscr{H}om_{\boldsymbol{k}}(\iota^{-1}\mathscr{F}_{\leqslant},\mu^{-1}\mathscr{G}) &= \mathscr{H}om_{\boldsymbol{k}}(p_1^{-1}\iota^{-1}\mathscr{F}_{\leqslant},p_1^{-1}\mu^{-1}\mathscr{G})\\ &\longrightarrow \mathscr{H}om_{\boldsymbol{k}}(\beta_{\leqslant}p_2^{-1}\iota^{-1}\mathscr{F}_{\leqslant},p_2^{-1}\mu^{-1}\mathscr{G}) = j_{\leqslant,*}j_{\leqslant}^{-1}p_2^{-1}\,\mathscr{H}om_{\boldsymbol{k}}(\iota^{-1}\mathscr{F}_{\leqslant},\mu^{-1}\mathscr{G}),\end{aligned}$$

where $j_{\leqslant}$ is the open inclusion $(\mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}})_{\leqslant} \hookrightarrow \mathcal{I}^{\text{ét}} \times_Y \mathcal{I}^{\text{ét}}$ (see [**35**, (2.3.18)] for the second equality). Since for any two sheaves $\mathscr{A}$ and $\mathscr{B}$ and an open set $Z$, the natural morphism $\mathrm{Hom}(\mathscr{A}_Z,\mathscr{B}) \to \mathrm{Hom}(j_Z^{-1}\mathscr{A},j_Z^{-1}\mathscr{B}) = \mathrm{Hom}(\mathscr{A},j_{Z,*}j_Z^{-1}\mathscr{B})$ is an isomorphism (see loc. cit.), we deduce a morphism

$$\beta_{\leqslant}p_1^{-1}\,\mathscr{H}om_{\boldsymbol{k}}(\iota^{-1}\mathscr{F}_{\leqslant},\mu^{-1}\mathscr{G}) \longrightarrow p_2^{-1}\,\mathscr{H}om_{\boldsymbol{k}}(\iota^{-1}\mathscr{F}_{\leqslant},\mu^{-1}\mathscr{G}),$$

as wanted. That it is the identity on the diagonal is clear. The construction is also clearly functorial (in a contravariant way). By Lemma 1.11, this functor is exact if $\mathscr{G}$ is injective in $\mathrm{Mod}(\boldsymbol{k}_Y)$.

***Corollary 1.24.*** — *For any object $\mathscr{G}$ of $D^+(\boldsymbol{k}_Y)$ there is a well-defined functor*

$$\boldsymbol{R}\mathscr{H}om(\iota^{-1}\bullet,\mu^{-1}\mathscr{G}) : D^{b,\mathrm{op}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \longrightarrow D^+(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}),$$

*and for any $\varphi \in \Gamma(U,\mathcal{I})$, we have $\varphi^{-1}\boldsymbol{R}\mathscr{H}om(\iota^{-1}\mathscr{F}_{\leqslant},\mu^{-1}\mathscr{G}) = \boldsymbol{R}\mathscr{H}om(\mathscr{F}_{\leqslant-\varphi},\mathscr{G})$.* □

**1.d. Traces for étale maps.** — Let us recall the definition of the *trace map* on sheaves of $\boldsymbol{k}$-vector spaces, where $\boldsymbol{k}$ is a field (more generally, a ring). Let $q : Z \to Z'$ be an étale map between topological spaces. For any $z' \in Z'$, the fibre $q^{-1}(z')$ is discrete with the induced topology. Let $\mathscr{F}'$ be a sheaf on $Z'$ and let $\widetilde{\mathscr{F}}$ be a subsheaf of $q^{-1}\mathscr{F}'$. We denote by $\mathrm{Tr}_q(\widetilde{\mathscr{F}}, \mathscr{F}')$ the subsheaf of $\mathscr{F}'$ defined by the following condition: for any open subset $U'$ of $Z'$, $\Gamma(U', \mathrm{Tr}_q(\widetilde{\mathscr{F}}, \mathscr{F}')) \subset \Gamma(U', \mathscr{F}')$ consists of those sections $f \in \Gamma(U', \mathscr{F}')$ such that, for any $z' \in U'$, the germ $f_{z'}$ belongs to $\sum_{z\in q^{-1}(z')} \widetilde{\mathscr{F}}_z$ (where we canonically identify $(q^{-1}\mathscr{F}')_z$ with $\mathscr{F}'_{q(z)}$, so that $\widetilde{\mathscr{F}}_z \subset \mathscr{F}'_{q(z)}$ and the sum is taken in $\mathscr{F}'_{q(z)}$). This clearly defines a subsheaf of $\mathscr{F}'$.

***Lemma 1.25.*** — *The germ* $\mathrm{Tr}_q(\widetilde{\mathscr{F}}, \mathscr{F}')_{z'}$ *is equal to* $\sum_{z\in q^{-1}(z')} \widetilde{\mathscr{F}}_z$ *(the sum is taken in* $\mathscr{F}'_{z'}$*).*

*Proof.* — Indeed, the inclusion $\subset$ is clear. For the converse inclusion, let $f'_{z'} \in \sum_{z\in q^{-1}(z')} \widetilde{\mathscr{F}}_z$. Then there exists $n \in \mathbb{N}$, $z_1, \dots, z_n \in q^{-1}(z)$ and $f_{z_i} \in \widetilde{\mathscr{F}}_{z_i}$ $(i = 1, \dots, n)$ such that $f'_{z'} = \sum_i f_{z_i}$. As $q$ is étale, there exist open neighbourhoods $V'_{z'}$ and $V_i$ of $z_i$ such that $q_i : V_i \to V'_{z'}$ is an homeomorphism for each $i$ and there exist representatives $f' \in \Gamma(V'_{z'}, \mathscr{F}')$ and $f_i \in \Gamma(V_i, \widetilde{\mathscr{F}})$ such that $f' = \sum_i f_i$ (by identifying $(q^{-1}\mathscr{F}')_{|V_i}$ to $\mathscr{F}'_{|V'_{z'}}$ via $q_i$). This shows that $f' \in \Gamma(V'_{z'}, \mathrm{Tr}_q(\widetilde{\mathscr{F}}, \mathscr{F}'))$ and therefore $f'_{z'} \in \mathrm{Tr}_q(\widetilde{\mathscr{F}}, \mathscr{F}')_{z'}$. ☐

If $Z$ is locally compact, then the functor $q_!$ is defined (see [**35**, §2.5]), and $\mathrm{Tr}_q$ is a functorial morphism $q_!q^{-1} \to \mathrm{Id}$.

There is a related notion of *saturation with respect to an order*, an operation that we denote by $\mathrm{Tr}_\leqslant$. Let $Y, \mathcal{I}, \mu$ be as in §1.b and let $\mathscr{F}$ be a sheaf of $\boldsymbol{k}$-vector spaces on $Y$ (where $\boldsymbol{k}$ is a field). If $\mathscr{F}'$ is a subsheaf of $\mu^{-1}\mathscr{F}$, the subsheaf $\mathrm{Tr}_\leqslant(\mathscr{F}', \mu^{-1}\mathscr{F})$ of $\mu^{-1}\mathscr{F}$ is defined by the following condition: for any open subset $V$ of $\mathcal{I}^{\text{ét}}$, $\Gamma(V, \mathrm{Tr}_\leqslant(\mathscr{F}', \mu^{-1}\mathscr{F}))$ consists of those sections $f \in \Gamma(V, \mu^{-1}\mathscr{F})$ such that, for any $\varphi_y \in V$, the germ of $f$ at $\varphi_y$ belongs to $\sum_{\psi_y \leqslant_y \varphi_y} \mathscr{F}'_{\psi_y}$.

***Lemma 1.26.*** — *The germ* $\mathrm{Tr}_\leqslant(\mathscr{F}', \mu^{-1}\mathscr{F})_{\varphi_y}$ *is equal to* $\sum_{\psi_y \leqslant_y \varphi_y} \mathscr{F}'_{\psi_y}$ *(the sum is taken in* $\mathscr{F}_y$*, due to the canonical identification of* $(\mu^{-1}\mathscr{F})_{\psi_y}$ *with* $\mathscr{F}_y$*).*

*Proof.* — The inclusion $\subset$ is clear. Let $f_{\varphi_y} \in \sum_{\psi_y \leqslant_y \varphi_y} \mathscr{F}'_{\psi_y}$. There exists a finite subset $\Phi_y \in \mu^{-1}(y)$ such that $\psi_y \leqslant \varphi_y$ for any $\psi_y \in \Phi_y$ and $f_{\varphi_y} = \sum_{\psi_y\in\Phi_y} f'_{\psi_y}$, with $f'_{\psi_y} \in \mathscr{F}'_{\psi_y}$, and the sum is taken in $\mathscr{F}_y$. Choose an open neighbourhood $U$ of $y$ in $Y$ such that the germs $\varphi_y$ and $\psi_y \in \Phi_y$ are defined on $U$. In particular, $\mu : \psi(U) \to U$ is a homeomorphism for any $\psi$. Up to shrinking $U$, we can also assume that each germ $f'_{\psi_y}$ is defined on $\psi(U)$ as a section $f'_\psi \in \Gamma(\psi(U), \mathscr{F}')$. Via $\mu$, we consider $\mathscr{F}'_{|\psi(U)}$ as a subsheaf of $\mathscr{F}_{|U}$ and we set $f_\varphi = \sum_{\psi_y\in\Phi_y} f'_\psi \in \Gamma(U, \mathscr{F}) = \Gamma(\varphi(U), \mu^{-1}(\mathscr{F}))$.

If $U$ is small enough, for any $z \in U$ and any $\psi_y \in \Phi_y$, we have $\psi_z \leqslant_z \varphi_z$, by the openness of $\leqslant$ and because $\Phi_y$ is finite. This implies that the germ of $f_\varphi$ at any such $z$ belongs to $\sum_{\eta_z \leqslant_y \varphi_z} \mathscr{F}'_{\eta_z}$, as was to be shown. $\square$

***Remark 1.27***. — The correspondence $(\mathscr{F}, \mathscr{F}') \mapsto (\mathscr{F}, \mathrm{Tr}_{\leqslant}(\mathscr{F}', \mu^{-1}\mathscr{F}))$ from the category of pairs $(\mathscr{F}, \mathscr{F}')$ with $\mathscr{F}' \subset \mu^{-1}\mathscr{F}$ (and morphisms being morphisms $\lambda : \mathscr{F}_1 \to \mathscr{F}_2$ such that $\mu^{-1}\lambda$ sends $\mathscr{F}'_1$ to $\mathscr{F}'_2$) to the category of pre-$\mathcal{I}$-filtered sheaves is functorial. Indeed, if $\mu^{-1}\lambda(\mathscr{F}'_1) \subset \mathscr{F}'_2$, then $\mu^{-1}\lambda\big[\mathrm{Tr}_{\leqslant}(\mathscr{F}'_2, \mu^{-1}\mathscr{F}_2))\big] \subset \mathrm{Tr}_{\leqslant}(\mathscr{F}'_2, \mu^{-1}\mathscr{F}_2))$.

**1.e. Pre-$\mathcal{I}$-filtrations of sheaves.** — We assume that $\mathcal{I}$ is as in §1.b. As above, $\mathscr{F}$ denotes a sheaf of $\boldsymbol{k}$-vector spaces on $Y$.

***Definition 1.28* (Pre-$\mathcal{I}$-filtration of a sheaf).** — We say that a subsheaf $\mathscr{F}_{\leqslant}$ of $\mu^{-1}\mathscr{F}$ is a *pre-$\mathcal{I}$-filtration* of $\mathscr{F}$ if, for any $y \in Y$ and any $\varphi_y, \psi_y \in \mu^{-1}(y)$, $\psi_y \leqslant_y \varphi_y$ implies $\mathscr{F}_{\leqslant \psi_y} \subset \mathscr{F}_{\leqslant \varphi_y}$ or, equivalently, if $\mathscr{F}_{\leqslant} = \mathrm{Tr}_{\leqslant}(\mathscr{F}_{\leqslant}, \mu^{-1}\mathscr{F})$. We will denote by $(\mathscr{F}, \mathscr{F}_\bullet)$ the data of a sheaf and a pre-$\mathcal{I}$-filtration on it.

In other words, a pre-$\mathcal{I}$-filtration of $\mathscr{F}$ is an object $\mathscr{F}_{\leqslant}$ of $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ together with an inclusion into the constant pre-$\mathcal{I}$-filtration $\mu^{-1}\mathscr{F}$: indeed, for any $\varphi_y \leqslant_y \psi_y$, the morphism $\mathscr{F}_{\leqslant \varphi_y} \to \mathscr{F}_{\leqslant \psi_y}$ given by the pre-$\mathcal{I}$-filtration is then an inclusion, since both $\mathscr{F}_{\leqslant \varphi_y} \to \mathscr{F}_y$ and $\mathscr{F}_{\leqslant \psi_y} \to \mathscr{F}_y$ are inclusions.

***Definition 1.29***. — Let $(\mathscr{F}, \mathscr{F}_\bullet), (\mathscr{F}', \mathscr{F}'_\bullet)$ be pre-$\mathcal{I}$-filtered sheaves and let $f : \mathscr{F} \to \mathscr{F}'$ be a morphism of sheaves. We say that it is a morphism of pre-$\mathcal{I}$-filtered sheaves if $\mu^{-1}f$ sends $\mathscr{F}_{\leqslant}$ to $\mathscr{F}'_{\leqslant}$.

We therefore get the (a priori non abelian) category of pre-$\mathcal{I}$-filtered sheaves of $\boldsymbol{k}$-vector spaces. Clearly, $\mu^{-1}f$ induces a morphism $\mathscr{F}_{\leqslant} \to \mathscr{F}'_{\leqslant}$ in the category $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. The notion of twist (Definition 1.10) applies to pre-$\mathcal{I}$-filtered sheaves.

***Definition 1.30* (Exhaustivity).** — We say that a pre-$\mathcal{I}$-filtration $\mathscr{F}_{\leqslant}$ of $\mathscr{F}$ is *exhaustive* if for any $y \in Y$ there exists $\varphi_y, \psi_y \in \mathcal{I}_y$ such that $\mathscr{F}_{\leqslant \varphi_y} = \mathscr{F}_y$ and $\mathscr{F}_{\leqslant \psi_y} = 0$.

We now introduce some operations on pre-$\mathcal{I}$-filtered sheaves.

*Extending the sheaf* $\mathcal{I}$. — Let $q : \widetilde{\mathcal{I}} \to \mathcal{I}'$ be a morphism of sheaves of ordered abelian groups on $Y'$. We also denote by $q : \widetilde{\mathcal{I}^{\text{ét}}} \to \mathcal{I}'^{\text{ét}}$ the corresponding map between étale spaces, so that we have a commutative diagram of maps:

$$\begin{tikzcd} \widetilde{\mathcal{I}^{\text{ét}}} \arrow[r, "q"] \arrow[dr, "\widetilde{\mu}"'] & \mathcal{I}'^{\text{ét}} \arrow[d, "\mu'"] \\ & Y' \end{tikzcd} \tag{1.31}$$

Let $(\mathscr{F}', \widetilde{\mathscr{F}}_\bullet)$ be a pre-$\widetilde{\mathcal{I}}$-filtered sheaf. We will define a pre-$\mathcal{I}'$-filtered sheaf $q_*(\mathscr{F}', \widetilde{\mathscr{F}}_\bullet) = (\mathscr{F}', (q\widetilde{\mathscr{F}})_\bullet)$. When we consider the diagram (1.31), we can introduce an ordered trace map as in Lemma 1.26.

***Lemma 1.32***. — *Let $(\mathscr{F}', \widetilde{\mathscr{F}}_\bullet)$ be a pre-$\widetilde{\mathcal{I}}$-filtered sheaf. Then there is a unique subsheaf $\operatorname{Tr}_{\leqslant q}(\widetilde{\mathscr{F}}_\leqslant, \mu'^{-1}\mathscr{F}')$ of $\mu'^{-1}\mathscr{F}'$ defined by the property that, for any $y' \in Y'$ and any $\varphi'_{y'} \in \mu'^{-1}(y')$,*

$$\operatorname{Tr}_{\leqslant q}(\widetilde{\mathscr{F}}_\leqslant, \mu'^{-1}\mathscr{F}')_{\varphi'_{y'}} = \sum_{\substack{\psi_{y'} \in \widetilde{\mathcal{I}}_{y'} \\ q(\psi_{y'}) \leqslant_{y'} \varphi'_{y'}}} \widetilde{\mathscr{F}}_{\leqslant \psi_{y'}}$$

*(where the sum is taken in $\mathscr{F}'_{y'}$) is a pre-$\mathcal{I}'$-filtration of $\mathscr{F}'$.*

*Proof*. — We note that $\operatorname{Tr}_{\leqslant q}(\widetilde{\mathscr{F}}_\leqslant, \mu'^{-1}\mathscr{F}') = \operatorname{Tr}_\leqslant\big(\operatorname{Tr}_q(\widetilde{\mathscr{F}}_\leqslant, \mu'^{-1}\mathscr{F}'), \mu'^{-1}\mathscr{F}'\big)$, so the lemma is a consequence of Lemmas 1.25 and 1.26. ☐

***Definition 1.33***. — Given a pre-$\widetilde{\mathcal{I}}$-filtration $\widetilde{\mathscr{F}}_\leqslant$ of $\mathscr{F}'$, we set

$$(q\widetilde{\mathscr{F}})_\leqslant := \operatorname{Tr}_{\leqslant q}(\widetilde{\mathscr{F}}_\leqslant, \mu'^{-1}\mathscr{F}').$$

*Restricting the sheaf* $\mathcal{I}$. — In the previous situation, let $(\mathscr{F}', \mathscr{F}'_\bullet)$ be a pre-$\mathcal{I}'$-filtered sheaf on $Y$. Then $\mathscr{F}'$ becomes pre-$\widetilde{\mathcal{I}}$-filtered by $q^{-1}\mathscr{F}'_\leqslant$.

*Pull-back*. — Let $f : Y' \to Y$ be a continuous map between locally compact spaces. Assume we are sheaves $\mathcal{I}$ on $Y$ and $\mathcal{I}'$ on $Y'$, and a morphism $q_f : \widetilde{\mathcal{I}} := f^{-1}\mathcal{I} \to \mathcal{I}'$ compatible with the order, where $\widetilde{\mathcal{I}}$ is equipped with the pull-back order (see (1.13)). Let $(\mathscr{F}, \mathscr{F}_\leqslant)$ be a $\mathcal{I}$-filtered sheaf on $Y$. We will define a pre-$\mathcal{I}'$-filtered sheaf $f^+(\mathscr{F}, \mathscr{F}_\leqslant) = (f^{-1}\mathscr{F}, f^+\mathscr{F}_\leqslant)$ on $Y'$.

We have a continuous map $\widetilde{f} : \widetilde{\mathcal{I}^{\text{ét}}} \to \mathcal{I}^{\text{ét}}$ over $f : Y' \to Y$. Then $f^{-1}(\mathscr{F}, \mathscr{F}_\leqslant) := (f^{-1}\mathscr{F}, \widetilde{f}^{-1}\mathscr{F}_\leqslant)$ is a pre-$\widetilde{\mathcal{I}}$-filtered sheaf of $\boldsymbol{k}$-vector spaces on $Y'$.

***Definition 1.34* (Pull-back).** — The pull-back $f^+(\mathscr{F}, \mathscr{F}_\leqslant)$ is the sheaf $\mathscr{F}' = f^{-1}\mathscr{F}$ equipped with the pre-$\mathcal{I}'$-filtration $f^+\mathscr{F}_\leqslant := (q_f f^{-1}\mathscr{F})_\leqslant = \operatorname{Tr}_{\leqslant q_f}(\widetilde{f}^{-1}\mathscr{F}_\leqslant, \mu'^{-1}f^{-1}\mathscr{F})$.

*Push-forward*. — The push-forward is a priori not defined at the level of pre-$\mathcal{I}$-filtrations, since the condition that $\mathscr{F}_\leqslant \to \mu^{-1}\mathscr{F}$ is injective may be not satisfied after the push-forward of Definition 1.23.

*Grading*. — Let us now assume that $\mathcal{I}^{\text{ét}}$ is Hausdorff. The relation $<_y$ is then also open and we can work with it as with $\leqslant_y$.

***Definition 1.35* (Grading a pre-$\mathcal{I}$-filtration (the Hausdorff case))**

Assume that $\mathcal{I}^{\text{ét}}$ is Hausdorff. Given a pre-$\mathcal{I}$-filtration $\mathscr{F}_\leqslant$ of $\mathscr{F}$, there is, due to the openness of $<$, a unique subsheaf $\mathscr{F}_<$ of $\mathscr{F}_\leqslant$ such that, for any $y \in Y$ and any $\varphi_y \in \mathcal{I}_y$, $\mathscr{F}_{<\varphi_y} = \sum_{\psi_y <_y \varphi_y} \mathscr{F}_{\leqslant\psi_y}$. We denote by $\operatorname{gr}\mathscr{F}$ the quotient sheaf $\mathscr{F}_\leqslant / \mathscr{F}_<$.

Notice that $\mathscr{F}_<$ is also a pre-$\mathcal{I}$-filtration of $\mathscr{F}$. Any morphism $(\mathscr{F},\mathscr{F}_\bullet)\to(\mathscr{F}',\mathscr{F}'_\bullet)$ sends $\mathscr{F}_<$ to $\mathscr{F}'_<$, hence defines a morphism $\operatorname{gr} f:\operatorname{gr}\mathscr{F}\to\operatorname{gr}\mathscr{F}'$.

**1.f. $\mathcal{I}$-filtrations of sheaves (the Hausdorff case).** — In order to define the notion of a $\mathcal{I}$-filtration, we will make the assumption that $\mathcal{I}^{\text{ét}}$ is Hausdorff. A more general definition will be given in §1.h. This assumption is now implicitly understood. The point is that, if $\mathcal{I}^{\text{ét}}$ is Hausdorff, it is locally compact (because $Y$ is so), and we make use of the direct image with proper supports in the usual way.

***Example 1.36* (Graded $\mathcal{I}$-filtrations).** — Let $\mathscr{G}$ be a sheaf of $\boldsymbol{k}$-vector spaces on $\mathcal{I}^{\text{ét}}$. We note that, for any $y\in Y$,

$$i_y^{-1}\mu_!\mathscr{G}=\mu_! i_{\mu^{-1}(y)}^{-1}\mathscr{G}=\bigoplus_{\varphi_y\in\mu^{-1}(y)}\mathscr{G}_{\varphi_y}. \tag{1.36 $*$}$$

Similarly, we have a commutative diagram of étale maps

$$\begin{tikzcd}
\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}} \arrow[r, "p_1"] \arrow[d, "p_2"'] & \mathcal{I}^{\text{ét}} \arrow[d, "\mu"] \\
\mathcal{I}^{\text{ét}} \arrow[r, "\mu"] & Y
\end{tikzcd}$$

and base change isomorphism $p_{2,!}p_1^{-1}\mathscr{G}=\mu^{-1}\mu_!\mathscr{G}$ (see e.g. [**35**, Prop. 2.5.11]). We denote by $(p_1^{-1}\mathscr{G})_\leqslant$ the extension by 0 of the restriction of $p_1^{-1}\mathscr{G}$ to $(\mathcal{I}^{\text{ét}}\times_Y\mathcal{I}^{\text{ét}})_\leqslant$, that is, $(p_1^{-1}\mathscr{G})_\leqslant=j_{\leqslant,!}j_\leqslant^{-1}p_1^{-1}\mathscr{G}$, and by $(\mu_!\mathscr{G})_\leqslant$ the subsheaf $p_{2,!}[(p_1^{-1}\mathscr{G})_\leqslant]$ of $p_{2,!}p_1^{-1}\mathscr{G}=\mu^{-1}\mu_!\mathscr{G}$.

For any $\varphi_y\in\mu^{-1}(y)$, one has $(\mu_!\mathscr{G})_{\leqslant\varphi_y}=\bigoplus_{\psi_y\leqslant_y\varphi_y}\mathscr{G}_{\psi_y}$, and $(\mu_!\mathscr{G})_\leqslant$ is a pre-$\mathcal{I}$-filtration on $\mu_!\mathscr{G}$, that we call the *graded $\mathcal{I}$-filtration on $\mu_!\mathscr{G}$*.

Note that, if $\mathcal{I}$ satisfies the exhaustivity property (Definition 1.3) and if $\mu$ is proper on $\operatorname{Supp}\mathscr{G}$ then, as the sum in (1.36 $*$) is finite, any graded $\mathcal{I}$-filtration is exhaustive.

***Definition 1.37* ($\mathcal{I}$-filtrations).** — Let $\mathscr{F}_\leqslant$ be a pre-$\mathcal{I}$-filtration of $\mathscr{F}$. We say that it is a $\mathcal{I}$-filtration of $\mathscr{F}$ if, *locally on $Y$*, $\mathscr{F}\simeq\mu_!\mathscr{G}$ and the inclusion $\mathscr{F}_\leqslant\subset\mu^{-1}(\mathscr{F})$ is isomorphic to the corresponding inclusion for the graded $\mathcal{I}$-filtration on $\mu_!\mathscr{G}$ for some sheaf $\mathscr{G}$ (a priori only locally (w.r.t. $Y$) defined on $\mathcal{I}^{\text{ét}}$).

The category of $\mathcal{I}$-filtered sheaves is the full subcategory of that of pre-$\mathcal{I}$-filtered sheaves whose objects are $\mathcal{I}$-filtered sheaves. In other words, a morphism between sheaves underlying $\mathcal{I}$-filtered sheaves is a morphism of $\mathcal{I}$-filtered sheaves iff it is compatible with order. Note that it is *not* required that $\mu^{-1}f$ is locally decomposed in the local graded models $\mu_!\mathscr{G},\mu_!\mathscr{G}'$ (i.e., $\mu^{-1}f$ is "filtered", but possibly not "graded").

***Example 1.38* (Grading a graded $\mathcal{I}$-filtration).** — For the graded $\mathcal{I}$-filtration on $\mu_!\mathscr{G}$, we have $(\mu_!\mathscr{G})_{<\varphi_y}=\bigoplus_{\psi_y<_y\varphi_y}\mathscr{G}_{\psi_y}$ and $\operatorname{gr}(\mu_!\mathscr{G})=\mathscr{G}$, so $\mathscr{G}$ can be recovered from $(\mu_!\mathscr{G})_\leqslant$.

Therefore, a $\mathcal{I}$-filtration $(\mathscr{F},\mathscr{F}_\bullet)$ is, locally (on $Y$), isomorphic to the graded $\mathcal{I}$-filtration on $\mu_!\operatorname{gr}\mathscr{F}$. Notice that the model graded $\mathcal{I}$-filtration exists globally on $Y$. In particular, for any $y\in Y$ and any $\varphi_y\in\mathcal{I}_y$, we have

$$\begin{aligned}\mathscr{F}_{<\varphi_y,y}&\simeq\bigoplus_{\substack{\psi_y\in\mathcal{I}_y\\ \psi_y<_y\varphi_y}}\operatorname{gr}_{\psi_y}\mathscr{F}_y,\\ \mathscr{F}_{\leqslant\varphi_y,y}&\simeq\bigoplus_{\substack{\psi_y\in\mathcal{I}_y\\ \psi_y\leqslant_y\varphi_y}}\operatorname{gr}_{\psi_y}\mathscr{F}_y=\mathscr{F}_{<\varphi_y,y}\oplus\operatorname{gr}_{\varphi_y}\mathscr{F}_y,\\ \mathscr{F}_y&\simeq\bigoplus_{\psi_y\in\mathcal{I}_y}\operatorname{gr}_{\psi_y}\mathscr{F}_y,\end{aligned}\tag{1.39}$$

in a way compatible with the inclusion $\mathscr{F}_{<\varphi_y,y}\subset\mathscr{F}_{\leqslant\varphi_y,y}\subset\mathscr{F}_y$. Recall also that a morphism may not be graded with respect to such a gradation.

***Remark 1.40***. — If $\mathcal{I}$ satisfies the exhaustivity property (Definition 1.3) and if $\mu$ is proper on $\operatorname{Supp}\operatorname{gr}\mathscr{F}$, then the $\mathcal{I}$-filtration is exhaustive.

***Lemma 1.41* (Stability by pull-back).** — *In the setting of Definition* 1.34*, if $\mathscr{F}_{\leqslant}$ is a $\mathcal{I}$-filtration of $\mathscr{F}$, then $f^+(\mathscr{F},\mathscr{F}_{\leqslant})$ is $\mathcal{I}'$-filtered sheaf on $Y'$.*

*Proof*. — Since the assertion is local, it is enough to start from a graded $\mathcal{I}$-filtration. Let us set $\mathscr{F}'_{\leqslant}=f^+\mathscr{F}_{\leqslant}$ and $\mathscr{F}_{\leqslant}=(\mu_!\mathscr{G})_{\leqslant}$. Then, for $y'\in Y'$ and $y=f(y')$, we have

$$\mathscr{F}'_{\leqslant_{y'}\varphi'_{y'}}=\sum_{\psi_y\leqslant_{y'}\varphi'_{y'}}\mathscr{F}_{\leqslant\psi_y}=\sum_{\psi_y\leqslant_{y'}\varphi'_{y'}}\bigoplus_{\varphi_y\leqslant_y\psi_y}\mathscr{G}_{\varphi_y}=\bigoplus_{\varphi_y\leqslant_{y'}\varphi'_{y'}}\mathscr{G}_{\varphi_y}.\qquad\square$$

***Remark 1.42* (co-$\mathcal{I}$-filtration).** — There is a dual notion of $\mathcal{I}$-filtration, that one can call a *co-$\mathcal{I}$-filtration*. It consists of a pair $(\mathscr{F},\mathscr{F}_<)$, where $\mathscr{F}_<$ is a pre-$\mathcal{I}$-filtration, and an injective morphism $\mathscr{F}_<\hookrightarrow\mu^{-1}\mathscr{F}$, which satisfy the local co-gradedness property, i.e.,

$$\mathscr{F}_y/\mathscr{F}_{<\varphi_y}=\bigoplus_{\substack{\psi_y\in\mathcal{I}_y\\ \varphi_y\leqslant_y\psi_y}}\operatorname{gr}_\psi\mathscr{F}_y.$$

From a $\mathcal{I}$-filtration $\mathscr{F}_{\leqslant}$, one defines a co-$\mathcal{I}$-filtration by using $\mathscr{F}_<$ of Definition 1.35. Conversely, given a co-$\mathcal{I}$-filtration $\mathscr{F}_<$ of $\mathscr{F}$, we define $\mathscr{F}_{\leqslant}$ by the formula

$$\mathscr{F}_{\leqslant\varphi_y}=\bigcap_{\substack{\psi_y\in\mathcal{I}_y\\ \varphi_y<_y\psi_y}}\mathscr{F}_{<\psi_y}.$$

Then $\mathscr{F}_{\leqslant}$ is a $\mathcal{I}$-filtration of $\mathscr{F}$.

*Classification of $\mathcal{I}$-filtered sheaves*. — Let $(\mathscr{F},\mathscr{F}_\bullet)$ and $(\mathscr{F}',\mathscr{F}'_\bullet)$ be two pre-$\mathcal{I}$-filtered sheaves. Then $\mathscr{H}\!om_{\boldsymbol{k}}(\mathscr{F},\mathscr{F}')$ is equipped with a natural pre-$\mathcal{I}$-filtration: for any $\varphi\in\Gamma(U,\mathcal{I})$, $\mathscr{H}\!om_{\boldsymbol{k}}(\mathscr{F},\mathscr{F}')_{\leqslant\varphi}$ is the subsheaf of $\mathscr{H}\!om_{\boldsymbol{k}}(\mathscr{F},\mathscr{F}')_{|U}$ such that, for any open set $V\subset U$, $f\in\operatorname{Hom}_{\boldsymbol{k}}(\mathscr{F}_{|V},\mathscr{F}'_{|V})$ belongs to $\operatorname{Hom}_{\boldsymbol{k}}(\mathscr{F}_{|V},\mathscr{F}'_{|V})_{\leqslant\varphi}$ iff $f(\mathscr{F}_{\leqslant\eta|V})\subset\mathscr{F}'_{\leqslant\eta+\varphi|V}$ for any $\eta\in\Gamma(V,\mathcal{I})$. In case $(\mathscr{F},\mathscr{F}_\bullet)$ and $(\mathscr{F}',\mathscr{F}'_\bullet)$ are $\mathcal{I}$-filtered sheaves, then the local decomposition of $\mathscr{F},\mathscr{F}'$ induces a local decomposition of $\mathscr{H}\!om_{\boldsymbol{k}}(\mathscr{F},\mathscr{F}')$.

We denote by $\mathscr{A}ut_{\boldsymbol{k}}^{<0}(\mathscr{F})$ the subsheaf $\mathrm{Id}+\mathscr{H}om_{\boldsymbol{k}}(\mathscr{F},\mathscr{F})_{<0}$ of $\mathscr{H}om_{\boldsymbol{k}}(\mathscr{F},\mathscr{F})_{\leqslant 0}$.

***Proposition 1.43.*** — *The set of isomorphism classes of $\mathcal{I}$-filtration $(\mathscr{F},\mathscr{F}_\bullet)$ with $\mathrm{gr}\,\mathscr{F}=\mathscr{G}$ is equal to $H^1\big(Y,\mathscr{A}ut_{\boldsymbol{k}}^{<0}(\mu_!\mathscr{G})\big)$, where $\mu_!\mathscr{G}$ is equipped with the graded $\mathcal{I}$-filtration.*

*Proof.* — This is standard (see e.g. the proof of Prop. II.1.2.2, p. 112 in [**2**]). ☐

**1.g. $\mathcal{I}$-filtered local systems (the Hausdorff case).** — The case where $\mathscr{F}$ is a *locally constant sheaf of finite dimensional $\boldsymbol{k}$-vector spaces on $Y$* (that we call a local system) will be the most important for us. Let us assume that $\mathscr{F}$ is a locally constant sheaf. As locally constant sheaves are stable by direct summand, it follows from the local isomorphism $\mathscr{F}\simeq\mu_!\,\mathrm{gr}\,\mathscr{F}$ that $\mu$ is proper on $\Sigma=\mathrm{Supp}\,\mathrm{gr}\,\mathscr{F}$, that $\Sigma$ is a finite covering of $Y$, and $\mathrm{gr}\,\mathscr{F}$ is a locally constant sheaf on its support. Moreover, $\mu_*\,\mathrm{gr}\,\mathscr{F}=\mu_!\,\mathrm{gr}\,\mathscr{F}$ is a local system on $Y$ locally isomorphic to $\mathscr{F}$. We call $\Sigma$ a $\mathcal{I}$-covering of $Y$.

We note that $\mu:\Sigma\to Y$, being a finite covering, is a local homeomorphism and therefore defines a subsheaf $\Sigma^{\mathrm{sh}}$ of $\mathcal{I}$. This subsheaf is a sheaf of ordered abelian sets (in general not abelian groups) with the ordered induced from that of $\mathcal{I}$. Restricting $\mathscr{F}_{\leqslant}$ to $\Sigma$ defines then a $\Sigma^{\mathrm{sh}}$-filtration of $\mathscr{F}$. Due to the local grading property of $\mathscr{F}_{\leqslant}$, it is not difficult to check that $\mathscr{F}_{\leqslant}$ is determined by $\mathscr{F}_{\leqslant|\Sigma}$ and that both $\mathscr{F}_{\leqslant}$ and $\mathscr{F}_{\leqslant|\Sigma}$ have the same graded sheaves. In other words, if we know $\mathscr{F}_{\leqslant\varphi_y,y}\subset\mathscr{F}_y$ for any $\varphi_y\in\Sigma_y$, we can reconstruct $\mathscr{F}_{\leqslant\psi_y,y}\subset\mathscr{F}_y$ for any $\psi_y\in\mathcal{I}_y$ by setting $\mathscr{F}_{\leqslant\psi_y,y}=\sum_{\varphi_y\leqslant\psi_y,\,\varphi_y\in\Sigma_y}\mathscr{F}_{\leqslant\varphi_y,y}$, where the sum is taken in $\mathscr{F}_y$.

Let us make precise Lemma 1.41 in the case where $\mathscr{F}$ is a local system. We consider the setting of Definition 1.34. Firstly, the pre-$\widetilde{\mathcal{I}}$-filtration $\widetilde{f}^{-1}\mathscr{F}_{\leqslant}$ of $f^{-1}\mathscr{F}$ is a $\widetilde{\mathcal{I}}$-filtration and we have $\mathrm{gr}\,f^{-1}\mathscr{F}=\widetilde{f}^{-1}\,\mathrm{gr}\,\mathscr{F}$, whose support $\widetilde{\Sigma}$ is nothing but $\widetilde{f}^{-1}\Sigma$, and is a finite covering of $Y'$. Since $q_f:\widetilde{\mathcal{I}^{\text{ét}}}\to\mathcal{I}'^{\text{ét}}$ is étale, being induced by a morphism of sheaves, the image $\Sigma'=q_f(\widetilde{\Sigma})$ is a finite covering of $Y'$ and $q_f:\widetilde{\Sigma}\to\Sigma'$ is also a finite covering. The proof of Lemma 1.41 shows that $\mathrm{gr}\,\mathscr{F}'=q_{f,*}\widetilde{f}^{-1}\,\mathrm{gr}\,\mathscr{F}$.

**1.h. $\mathcal{I}$-filtered local systems (the stratified Hausdorff case).** — We will now extend the notion of $\mathcal{I}$-filtered local system in the case $\mathcal{I}$ satisfies a property weaker than Hausdorff, that we call the *stratified Hausdorff property.* We restrict to local systems, as this will be the only case of interest for us, and we will thus avoid general definitions as for the Hausdorff case.

Let $\mathscr{Y}=(Y_\alpha)_{\alpha\in A}$ be a stratification of $Y$ (that is, a locally finite partition of $Y$ into locally closed sets, called strata, such that the closure of every stratum is a union of strata). For $\alpha,\beta\in A$, we will denote by $\alpha\preccurlyeq\beta$ the relation $Y_\alpha\subset\overline{Y}_\beta$. In the following, we will always assume that the stratification satisfies the following property:

(1.44) For each $\alpha\in A$, each point $y\in Y_\alpha$ has a fundamental system of open neighbourhoods $\mathrm{nb}(y)$ such that, for any $\beta\succcurlyeq\alpha$, $\mathrm{nb}(y)\cap\overline{Y}_\beta$ is contractible.

***Definition 1.45* (The stratified Hausdorff property).** — Let $\mathcal{I}$ be a sheaf on $Y$ (in the setting of §1.a). We say that $\mathcal{I}$ satisfies the *Hausdorff property with respect to* $\mathscr{Y}$ if for any $Y_\alpha \in \mathscr{Y}$, $\mu^{-1}(Y_\alpha)$ is Hausdorff, that is, the sheaf-theoretic restriction of $\mathcal{I}$ to every $Y_\alpha$ satisfies the Hausdorff property.

***Examples 1.46***

(1) If $\mathcal{I}$ is a constructible sheaf on $Y$ with respect to a stratification $\mathscr{Y}$ for which the strata are locally connected, then $\mathcal{I}$ is Hausdorff with respect to $\mathscr{Y}$.

(2) Let $X$ be a complex manifold, let $D$ be a reduced divisor in $X$ and let $\mathscr{Y}$ be a stratification of $D$ by complex locally closed submanifolds such that $(X \smallsetminus D, \mathscr{Y})$ is a Whitney stratification of $X$. Then $\mathcal{I} = \mathscr{O}_X(*D)/\mathscr{O}_X$ has the Hausdorff property with respect to $\mathscr{Y}$. Indeed, if $x$ belongs to a connected stratum $Y_\alpha$ and if a germ $\varphi_x \in \mathscr{O}_{X,x}(*D)$, with representative $\varphi$, is such that $\varphi_y$ is holomorphic for $y \in Y_\alpha$ close enough to $x$, then $\varphi$ is holomorphic on each maximal dimensional connected stratum of $D$ which contains $Y_\alpha$ in its closure, so, by the local product property, $\varphi$ is holomorphic on the smooth part of $D$ in the neighbourhood of $x$. By Hartogs, $\varphi$ is holomorphic near $x$.

(3) The same argument holds if $X$ is only assumed to be a normal complex space and $D$ is a locally principal divisor in $X$.

We have seen in §1.g that the support of $\operatorname{gr}\mathscr{F}$ is a finite covering of $Y$ contained in $\mathcal{I}^{\text{ét}}$. We now introduce the notion of stratified $\mathcal{I}$-covering (with respect to $\mathscr{Y}$).

***Definition 1.47* (stratified $\mathcal{I}$-covering).** — Let $\Sigma \subset \mathcal{I}^{\text{ét}}$ be a closed subset. We will say that $\mu_{|\Sigma} : \Sigma \to Y$ is a stratified $\mathcal{I}$-covering of $Y$ if

(1) $\mu_{|\Sigma} : \Sigma \to Y$ is a local homeomorphism, that is, $\Sigma$ is the étale space of a subsheaf $\Sigma^{\text{sh}}$ of $\mathcal{I}$,

(2) for each $\alpha \in A$, setting $\Sigma_\alpha = \Sigma \cap \mu^{-1}(Y_\alpha)$, the map $\mu_{|\Sigma_\alpha} : \Sigma_\alpha \to Y_\alpha$ is a finite covering,

Let us explain this definition. The first property implies that $\mu_{|\Sigma_\alpha}$ is a local homeomorphism on $Y_\alpha$. Since $\mathcal{I}^{\text{ét}}_{|Y_\alpha}$ is Hausdorff, so is $\Sigma_\alpha$, and the second property is then equivalent to $\mu_{|\Sigma_\alpha}$ being proper.

What does $\Sigma$ look like near a point of $Y_\alpha$? For each $\alpha \in A$, each $y \in Y_\alpha$ has an open neighbourhood $\operatorname{nb}(y) \subset Y$ such that, denoting $\varphi_y^{(i)}$ the points in $\mu_{|\Sigma_\alpha}^{-1}(y)$, the germs $\varphi_y^{(i)}$ are induced by sections $\varphi^{(i)} \in \Gamma(\operatorname{nb}(y), \mathcal{I})$. Then we have $\Sigma \cap \mu^{-1}(\operatorname{nb}(y)) = \bigcup_i \varphi^{(i)}(\operatorname{nb}(y))$. We note however that, while the germs $\varphi_y^{(i)}$ (or more generally the germs $\varphi_{y'}^{(i)}$ for $y' \in \operatorname{nb}(y) \cap Y_\alpha$) are distinct, such may not be the case when $y' \notin Y_\alpha$ and, for $\beta \succcurlyeq \alpha$, $\Sigma_\beta \cap \mu^{-1}(\operatorname{nb}(y))$ may have less connected components than $\Sigma_\alpha \cap \mu^{-1}(\operatorname{nb}(y))$.

***Definition 1.48* ($\mathcal{I}$-filtration (the stratified Hausdorff case)).** — Let $\mathcal{I}$ be stratified Hausdorff with respect to $\mathscr{Y}$, let $\mathscr{F}$ be a locally constant sheaf of finite dimensional

$\boldsymbol{k}$-vector spaces on $Y$ and let $\mathscr{F}_{\leqslant}$ be a pre-$\mathcal{I}$-filtration of $\mathscr{F}$. We say that $\mathscr{F}_{\leqslant}$ is a $\mathcal{I}$-filtration of $\mathscr{F}$ if

(1) for each $\alpha \in A$, the restriction $\mathscr{F}_{\leqslant|\mu^{-1}(Y_\alpha)}$ is a $\mathcal{I}_{|\mu^{-1}(Y_\alpha)}$-filtration of $\mathscr{F}_{|Y_\alpha}$ in the sense of Definition 1.37, with associated covering $\Sigma_\alpha \subset \mathcal{I}^{\text{ét}}_{|\mu^{-1}(Y_\alpha)}$ and associated local system $\mathscr{G}_\alpha = \operatorname{gr}\mathscr{F}_{|\mu^{-1}(Y_\alpha)}$,

(2) the set $\Sigma = \bigcup_\alpha \Sigma_\alpha$ is a stratified $\mathcal{I}$-covering of $Y$,

(3) for each $\alpha \in A$ and each $y \in Y_\alpha$, let us set $\mu^{-1}(y) \cap \Sigma_\alpha = \{\varphi_y^{(i)} \mid i \in I_\alpha\}$, and let $\operatorname{nb}(y)$ be a small open neighbourhood on which all $\varphi^{(i)}$ are defined as sections of $\mathcal{I}$; for each $\beta \succcurlyeq \alpha$, let $\Sigma_\beta(y,j)$ be the connected components of $\mu^{-1}(\operatorname{nb}(y) \cap \Sigma_\beta) = \bigcup_i \varphi^{(i)}(\operatorname{nb}(y) \cap Y_\beta)$; then, for each $j$, $\mathscr{G}_{\beta|\Sigma_\beta(y,j)}$ is a trivial local system and

$$\operatorname{rk}\mathscr{G}_{\beta|\Sigma_\beta(y,j)} = \sum_{i|\varphi^{(i)}(\operatorname{nb}(y)\cap Y_\beta)=\Sigma_\beta(y,j)} \operatorname{rk}\mathscr{G}_{\alpha|\varphi^{(i)}(\operatorname{nb}(y)\cap Y_\alpha)}.$$

The last condition is a gluing condition near a stratum $Y_\alpha$. If the stratified space is reasonable, each $y \in Y_\alpha$ has a fundamental system of simply connected neighbourhood such that $\operatorname{nb}(y) \cap Y_\alpha$ is also simply connected. Then the local system $\mathscr{G}_\alpha$, restricted to $\varphi^{(i)}(\operatorname{nb}(y) \cap Y_\alpha)$ extends in a unique way to a local system $\mathscr{G}_\alpha^{(i)}$ on $\varphi^{(i)}(\operatorname{nb}(y))$ (both local systems are trivial, according to our assumptions on $\operatorname{nb}(y)$). The second condition means that $\mathscr{G}_{\beta|\Sigma_\beta(y,j)}$ is isomorphic to the direct sum of the $\mathscr{G}_\alpha^{(i)}$ restricted to $\operatorname{nb}(y) \cap Y_\beta$.

As in the Hausdorff case, we also note that the restriction of $\mathscr{F}_{\leqslant}$ to $\Sigma$ defines a $\Sigma^{\text{sh}}$-filtration of $\mathscr{F}$, and that $\mathscr{F}_{\leqslant}$ is determined by $\mathscr{F}_{\leqslant|\Sigma}$.

***Lemma 1.49* (Stability by pull-back).** — *Let $f : (Y', \mathscr{Y}') \to (Y, \mathscr{Y})$ be a stratified continuous map, let $\mathcal{I}$ be stratified Hausdorff with respect to $\mathscr{Y}$ and let $(\mathscr{F}, \mathscr{F}_{\leqslant})$ be a $\mathcal{I}$-filtered local system. Assume that we are in the setting of Definition* 1.34. *Then $f^+(\mathscr{F}, \mathscr{F}_{\leqslant})$ is $\mathcal{I}'$-filtered local system with associated $\mathcal{I}'$ stratified covering $\Sigma'$ given by $q_f(\widetilde{f}^{-1}(\Sigma))$.*

*Proof.* — The result on each stratum follows from Lemma 1.41 and the details given in §1.g. The point is to check the gluing properties 1.48(2) and 1.48(3). Firstly, these properties are easily checked for $\widetilde{f}^{-1}\mathscr{F}_{\leqslant}$ as a $\widetilde{\mathcal{I}}$-filtration, and for $\widetilde{\Sigma} = \widetilde{f}^{-1}\Sigma$. One then shows that $q_f : \widetilde{\Sigma} \to \Sigma' := q_f(\Sigma)$ is a stratified covering with respect to the stratification $(\Sigma'_{\alpha'})_{\alpha' \in A'}$. The desired properties follow. ☐

**1.i. Comments.** — The first definition of a $\mathcal{I}$-filtered local system appears in [**15**] in the following way (taking the notation of the present notes):

Given the local system $\mathcal{I}$ of multi-valued differential forms on the circle $S^1$, equipped with the order as defined in Example 1.6, a $\mathcal{I}$-filtration on a local system $\mathscr{L}$ on $S^1$ consists of the data of a family of subsheaves $\mathscr{L}^\varphi$ of $\mathscr{L}$ indexed by $\mathcal{I}$ (be careful that $\mathcal{I}$ is only a local system, but this is enough) satisfying the local grading property like (1.39).

This definition is taken up in [**46**, §4, Def. 4.1], and later in [**49**, Def. 2.1, p. 57] in a similar way, but B. Malgrange adds between brackets:

“Be careful that the $\mathscr{L}^\varphi$ are not subsheaves in the usual sense, because they are indexed by a local system and not a set.”

Later, the definition is once more taken up in [**2**, p. 74–77], starting by defining a $\mathcal{I}$-graded local system (this is similar to Example 1.36) and then a $\mathcal{I}$-filtered local system by gluing local $\mathcal{I}$-graded local systems in a way which preserves the filtration (but possibly not the gradation).

In this lecture we have tried to give a precise intrinsic answer to the question:

what *is* a $\mathcal{I}$-filtered local system?

in order to use it in higher dimensions.

Another approach is due to T. Mochizuki in [**65**, §2.1], starting with a family of filtrations indexed by ordered sets, and adding compatibility conditions (corresponding to compatibility with respect to restriction morphisms in the sheaf-theoretic approach of the present notes).

# PART I

# DIMENSION ONE

# LECTURE 2

# STOKES-FILTERED LOCAL SYSTEMS IN DIMENSION ONE

**Summary.** We consider Stokes filtrations on local systems on $S^1$. We review some of the definitions of the previous lecture in this case and make explicit the supplementary properties coming from this particular case. This lecture can be read independently of Lecture 1. References are [**15**], [**46**], [**2**] and [**49**, Chap. IV].

**2.a. Non-ramified Stokes-filtered local systems.** — Let $\boldsymbol{k}$ be a field. In this section, we consider local systems of finite dimensional $\boldsymbol{k}$-vector spaces on $S^1$. The indexing sheaf will be the constant sheaf $\mathcal{I}$ with fibre $\mathcal{P}$, ordered as as in Example 1.6, namely, if $\eta \neq 0$, setting $\eta = u_n(x)x^{-n}$ with $n \geqslant 1$ and $u_n(0) \neq 0$,

$$\eta \leqslant_{\theta} 0 \iff \eta = 0 \text{ or } \arg u_n(0) - n\theta \in (\pi/2, 3\pi/2) \mod 2\pi.$$

Let us rephrase Definition 1.28 in this setting.

***Definition 2.1.*** — A non-ramified pre-Stokes filtration on a local system $\mathscr{L}$ of finite dimensional $\boldsymbol{k}$-vector spaces on $S^1$ consists of the data of a family of subsheaves $\mathscr{L}_{\leqslant\varphi}$ indexed by $\mathcal{P}$ such that, for any $\theta \in S^1$, $\varphi \leqslant_{\theta} \psi \Rightarrow \mathscr{L}_{\leqslant\varphi,\theta} \subset \mathscr{L}_{\leqslant\psi,\theta}$.

The étale space $\mathcal{I}^{\text{ét}}$ is Hausdorff (see Example 1.1(1)). Let us set, for any $\varphi \in \mathcal{P}$ and any $\theta \in S^1$,

$$\mathscr{L}_{<\varphi,\theta} = \sum_{\psi <_{\theta} \varphi} \mathscr{L}_{\leqslant\psi,\theta}. \tag{2.2}$$

This defines a subsheaf $\mathscr{L}_{<\varphi}$ of $\mathscr{L}_{\leqslant\varphi}$, and we set $\operatorname{gr}_{\varphi} \mathscr{L} = \mathscr{L}_{\leqslant\varphi}/\mathscr{L}_{<\varphi}$ (see Definition 1.35).

***Notation 2.3.*** — We rephrase here Notation 1.5 in the present setting. Let $\varphi, \psi \in \mathcal{P}$. Recall that we denote by $S^1_{\psi\leqslant\varphi} \subset S^1$ is the subset of $S^1$ consisting of the $\theta$ for which $\psi \leqslant_{\theta} \varphi$. Similarly, $S^1_{\psi<\varphi} \subset S^1$ is the subset of $S^1$ consisting of the $\theta$ for which $\psi <_{\theta} \varphi$.

Both subsets are a finite union of open intervals. They are equal if $\varphi \neq \psi$. Otherwise, $S^1_{\varphi\leqslant\varphi} = S^1$ and $S^1_{\varphi<\varphi} = \varnothing$.

Given a sheaf $\mathscr{F}$ on $S^1$, we will denote by $\beta_{\psi\leqslant\varphi}\mathscr{F}$ the sheaf obtained by restricting $\mathscr{F}$ to the open set $S^1_{\psi\leqslant\varphi}$ and extending it by 0 as a sheaf on $S^1$ (for any open set $Z \subset S^1$, this operation is denoted $\mathscr{F}_Z$ in [**35**]). A similar definition holds for $\beta_{\psi<\varphi}\mathscr{F}$.

***Definition 2.4*** **((Graded) Stokes filtration).** — Given a finite set $\Phi \subset \mathcal{P}$, a Stokes-graded local system with $\Phi$ as set of exponential factors consists of the data of local systems (that we denote by) $\operatorname{gr}_\varphi \mathscr{L}$ on $S^1$ ($\varphi \in \Phi$). Then the graded non-ramified Stokes filtration on $\operatorname{gr}\mathscr{L} := \bigoplus_{\psi\in\Phi}\operatorname{gr}_\psi\mathscr{L}$ is given by

$$(\operatorname{gr}\mathscr{L})_{\leqslant\varphi} = \bigoplus_{\psi\in\Phi}\beta_{\psi\leqslant\varphi}\operatorname{gr}_\psi\mathscr{L}.$$

We the also have

$$(\operatorname{gr}\mathscr{L})_{<\varphi} = \bigoplus_{\psi\in\Phi}\beta_{\psi<\varphi}\operatorname{gr}_\psi\mathscr{L}.$$

A *non-ramified* $\boldsymbol{k}$*-Stokes filtration* on $\mathscr{L}$ is a pre-Stokes filtration which is locally on $S^1$ isomorphic to a graded Stokes filtration. It is denoted by $\mathscr{L}_\bullet$.

For a Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$, each sheaf $\operatorname{gr}_\varphi\mathscr{L}$ is a (possibly zero) local system on $S^1$. By definition, for every $\varphi$ and every $\theta_o \in S^1$, we have on some neighbourhood $\operatorname{nb}(\theta_o)$ of $\theta_o$,

$$\begin{aligned}
&\mathscr{L}_{<\varphi|\operatorname{nb}(\theta_o)} \simeq \bigoplus_{\psi\in\Phi}\beta_{\psi<\varphi}\operatorname{gr}_\psi\mathscr{L}_{|\operatorname{nb}(\theta_o)},\\
&\mathscr{L}_{\leqslant\varphi|\operatorname{nb}(\theta_o)} \simeq \mathscr{L}_{<\varphi|\operatorname{nb}(\theta_o)} \oplus \operatorname{gr}_\varphi\mathscr{L}_{|\operatorname{nb}(\theta_o)} = \bigoplus_{\psi\in\Phi}\beta_{\psi\leqslant\varphi}\operatorname{gr}_\psi\mathscr{L}_{|\operatorname{nb}(\theta_o)},\\
&\mathscr{L}_{|\operatorname{nb}(\theta_o} \simeq \bigoplus_{\psi\in\Phi}\operatorname{gr}_\psi\mathscr{L}_{|\operatorname{nb}(\theta_o)}
\end{aligned}\tag{2.5}$$

in a way compatible with the natural inclusions.

***Exercise 2.6.*** — Show that the category of Stokes-filtered local systems has direct sums, and that any Stokes-graded local system is the direct sum of Stokes-graded local systems, each of which has exactly one exponential factor.

One can make more explicit the definition of a non-ramified Stokes-filtered local system.

***Lemma 2.7.*** — *Giving a non-ramified Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$ is equivalent to giving, for each $\varphi \in \mathcal{P}$, a $\mathbb{R}$-constructible subsheaf $\mathscr{L}_{\leqslant\varphi} \subset \mathscr{L}$ subject to the following conditions:*

(1) *for any $\theta \in S^1$, the germs $\mathscr{L}_{\leqslant\varphi,\theta}$ form an exhaustive increasing filtration of $\mathscr{L}_\theta$;*

(2) *defining $\mathscr{L}_{<\varphi}$, and therefore $\operatorname{gr}_\varphi\mathscr{L}$, from the family $\mathscr{L}_{\leqslant\psi}$ as in* (2.2)*, the sheaf $\operatorname{gr}_\varphi\mathscr{L}$ is a local system of finite dimensional $\boldsymbol{k}$-vector spaces on $S^1$;*

(3) *for any $\theta \in S^1$ and any $\varphi \in \mathcal{P}$, $\dim\mathscr{L}_{\leqslant\varphi,\theta} = \sum_{\psi\leqslant_\theta\varphi}\dim\operatorname{gr}_\psi\mathscr{L}_\theta$.*

We note that when 2.7(2) is satisfied, 2.7(3) is equivalent to

($3'$) For any $\theta \in S^1$ and any $\varphi \in \mathcal{P}$, $\dim \mathscr{L}_{<\varphi,\theta} = \sum_{\psi <_\theta \varphi} \dim \operatorname{gr}_\psi \mathscr{L}_\theta$.

*Proof of Lemma* 2.7. — The point is to get the local gradedness property from the dimension property of 2.7(3).

***Lemma 2.8.*** — *Let $\mathscr{F}$ be a $\mathbb{R}$-constructible sheaf of $\boldsymbol{k}$-vector spaces on $S^1$. For any $\theta_o \in S^1$, let $I$ be an open interval containing $\theta_o$ such that $\mathscr{F}_{|I\smallsetminus\{\theta_o\}}$ is a local system of finite dimensional $\boldsymbol{k}$-vector spaces. Then $H^1(I, \mathscr{F}) = 0$.*

*Proof.* — Let $\iota : I \smallsetminus \{\theta_o\} \hookrightarrow I$ be the inclusion. We have an exact sequence $0 \to \iota_!\iota^{-1}\mathscr{F} \to \mathscr{F} \to \mathscr{G} \to 0$, where $\mathscr{G}$ is supported at $\theta_o$. It is therefore enough to prove the result for $\iota_!\iota^{-1}\mathscr{F}$. This reduces to the property that, if $i : (a, b) \hookrightarrow (a, b]$ is the inclusion (with $a, b \in \mathbb{R}$, $a < b$), then $H^1((a, b], i_!\boldsymbol{k}) = 0$, which is clear by Poincaré duality. ☐

Let us fix $\theta_o \in S^1$ and let $\mathrm{nb}(\theta_o)$ be an open interval of $S^1$ containing $\theta_o$ on which $(\mathscr{L}, \mathscr{L}_\bullet)$ is graded with (finite) set of exponential factors $\Phi_{\theta_o}$, and such that, for any $\psi \in \Phi_{\theta_o}$, $\mathscr{L}_{<\psi}$ is a local system on $\mathrm{nb}(\theta_o) \smallsetminus \{\theta_o\}$. Then, for any $\psi \in \Phi_{\theta_o}$, $H^1(\mathrm{nb}(\theta_o), \mathscr{L}_{<\psi}) = 0$ and, as $\operatorname{gr}_\psi \mathscr{L}$ is a constant local system on $\mathrm{nb}(\theta_o)$, we can lift a basis of global sections of $\operatorname{gr}_\psi \mathscr{L}_{|\mathrm{nb}(\theta_o)}$ as a family of sections of $\mathscr{L}_{\leqslant\psi,|\mathrm{nb}(\theta_o)}$. This defines a morphism $\bigoplus_\psi \operatorname{gr}_\psi \mathscr{L}_{|\mathrm{nb}(\theta_o)} \to \mathscr{L}_{|\mathrm{nb}(\theta_o)}$ sending $\bigoplus_{\psi\leqslant_\theta\varphi} \operatorname{gr}_\psi \mathscr{L}_\theta$ to $\mathscr{L}_{\leqslant\varphi,\theta}$ for any $\theta \in \mathrm{nb}(\theta_o)$ and any $\varphi$. By induction on $\#\{\psi \in \Phi \mid \psi \leqslant_\theta \varphi\}$, and using (2.2), it sends $\bigoplus_{\psi\leqslant_\theta\varphi} \operatorname{gr}_\psi \mathscr{L}_\theta$ *onto* $\mathscr{L}_{\leqslant\varphi,\theta}$ for any $\theta \in \mathrm{nb}(\theta_o)$ and any $\varphi$. As both spaces have the same dimension, due to 2.7(3), this is an isomorphism. ☐

Since $\operatorname{gr}_\varphi \mathscr{L}$ is a local system, it is zero if and only if it is zero near some $\theta_o$. By the local grading property, we conclude that the sets $\Phi_{\theta_o} \subset \mathcal{P}$ such that $\operatorname{gr}_\varphi \mathscr{L}_{\theta_o} \neq 0$ is finite and independent of $\theta_o$, and we simply denote it by $\Phi$. It is called the set of exponential factors of the non-ramified Stokes filtration. The following proposition is easily checked, showing more precisely exhaustivity.

***Proposition 2.9.*** — *Let $\mathscr{L}_\bullet$ be a non-ramified $\boldsymbol{k}$-Stokes filtration on $\mathscr{L}$. Then, for any $\theta \in S^1$, and any $\varphi \in \mathcal{P}$,*

- *if $\varphi <_\theta \Phi$, then $\mathscr{L}_{\leqslant\varphi,\theta} = 0$,*
- *if $\Phi <_\theta \varphi$, then $\mathscr{L}_{<\varphi,\theta} = \mathscr{L}_{\leqslant\varphi,\theta} = \mathscr{L}_\theta$.* ☐

***Remark 2.10.*** — One can also remark that the category of Stokes-filtered local systems with set of exponential factors contained in $\Phi$ is equivalent to the category of $\Phi$-filtered local systems, where we regard $\Phi$ as a constant sheaf on $S^1$, equipped with the ordered induced by that of $\mathcal{P}$.

***Examples 2.11***

(1) The graded Stokes filtration with $\Phi = \{0\}$ (see Example 1.36) on the constant sheaf $\boldsymbol{k}_{S^1}$ is defined by $\boldsymbol{k}_{S^1,\leqslant\varphi} := \beta_{0\leqslant\varphi}\boldsymbol{k}_{S^1}$ for any $\varphi$, so that $\boldsymbol{k}_{S^1,\leqslant 0} = \boldsymbol{k}_{S^1}$, $\boldsymbol{k}_{S^1,<0} = 0$, and, for any $\varphi \neq 0$, $\boldsymbol{k}_{S^1,\leqslant\varphi} = \boldsymbol{k}_{S^1,<\varphi}$ has germ equal to $\boldsymbol{k}_\theta$ at $\theta \in S^1$ iff $0 \leqslant_\theta \varphi$, and has germ equal to 0 otherwise.

(2) Let $\mathscr{L}_\bullet$ be a non-ramified Stokes filtration with set of exponential factors $\Phi$ reduced to one element $\eta$. According to Proposition 2.9, we have $\mathscr{L}_{<\eta} = 0$, and $\mathscr{L}_{\leqslant\eta} = \operatorname{gr}_\eta \mathscr{L} = \mathscr{L}$ is a local system on $S^1$. The non-ramified Stokes filtration is then described as in 2.11(1) above, that is, $\mathscr{L}_{\leqslant\varphi} = \beta_{\eta\leqslant\varphi}\mathscr{L}$. If we denote by $\mathscr{L}_\bullet$ this Stokes filtration of $\mathscr{L}$ then, using the twist operation of Definition 1.10, the twisted Stokes filtration $\mathscr{L}[-\eta]_\bullet$ is nothing but the graded Stokes filtration on $\mathscr{L}$, defined as in 2.11(1).

(3) Assume that $\#\Phi = 2$ or, equivalently (by twisting, see above), that $\Phi = \{0, \varphi_o\}$ with $\varphi_o \neq 0$. If the order of the pole of $\varphi_o$ is $n$, then there are $2n$ Stokes directions (see Example 1.6) dividing the circle in $2n$ intervals. Given such an open interval, then 0 and $\varphi_o$ are comparable (in the same way) at any $\theta$ in the interval, and the comparison changes alternatively on the intervals. Assume that $0 <_\theta \varphi_o$. Then, according to Proposition 2.9 and to 2.7(3), $\mathscr{L}_{\leqslant\varphi_o,\theta} = \mathscr{L}_\theta$ and $\mathscr{L}_{<0,\theta} = 0$. Moreover, when restricted to the open interval containing $\theta$, $\mathscr{L}_{\leqslant 0} = \mathscr{L}_{<\varphi_o}$ is a local system of rank equal to $\operatorname{rk}\operatorname{gr}_0 \mathscr{L}$. On the other intervals, the roles of 0 and $\varphi_o$ are exchanged.

Let now $\theta$ be a Stokes direction for $(0, \varphi_o)$. As $\varphi_o$ and 0 are not comparable at $\theta$, 2.7(3′) implies that $\mathscr{L}_{<\varphi_o,\theta} = \mathscr{L}_{<0,\theta} = 0$ and, using $\varphi$ such that $0, \varphi_o <_\theta \varphi$, we find, by exhaustivity, $\mathscr{L}_\theta = \mathscr{L}_{\leqslant\varphi_o,\theta} \oplus \mathscr{L}_{\leqslant 0,\theta}$. This decomposition reads as an isomorphism $\mathscr{L}_\theta \simeq \operatorname{gr}_{\varphi_o} \mathscr{L}_\theta \oplus \operatorname{gr}_0 \mathscr{L}_\theta$. It extends on a neighbourhood $\operatorname{nb}(\theta)$ of $\theta$ in $S^1$ (we can take for $\operatorname{nb}(\theta)$ the union of the two adjacent intervals ending at $\theta$) in a unique way as an isomorphism of local systems $\mathscr{L}_{|\operatorname{nb}(\theta)} \simeq (\operatorname{gr}_{\varphi_o} \mathscr{L} \oplus \operatorname{gr}_0 \mathscr{L})_{|\operatorname{nb}(\theta)}$.

In order to end the description, we will show that $\mathscr{L}_{\leqslant\varphi} = \mathscr{L}_{<\varphi}$ for $\varphi \notin \{0, \varphi_o\}$ can be deduced from the data of the corresponding sheaves for $\varphi \in \{0, \varphi_o\}$. Let us fix $\theta \in S^1$. Assume first $0 <_\theta \varphi_o$ (and argue similarly if $\varphi_o <_\theta 0$).

- If $\varphi$ is neither comparable to $\varphi_o$ nor to 0, then 2.7(3) shows that $\mathscr{L}_{\leqslant\varphi,\theta} = 0$.
- If $\varphi$ is comparable to $\varphi_o$ but not to 0,
  - if $\varphi_o <_\theta \varphi$, then $\mathscr{L}_{\leqslant\varphi,\theta} = \mathscr{L}_\theta$,
  - if $\varphi <_\theta \varphi_o$, then $\mathscr{L}_{\leqslant\varphi,\theta} \subset \mathscr{L}_{<\varphi_o,\theta} = \mathscr{L}_{\leqslant 0,\theta}$, hence 2.7(3) implies $\mathscr{L}_{\leqslant\varphi,\theta} = 0$.
- If $\varphi$ is comparable to 0 but not to $\varphi_o$, the result is similar.
- If $\varphi$ is comparable to both $\varphi_o$ and 0, then,
  - if $0 <_\theta \varphi_o <_\theta \varphi$, $\mathscr{L}_{\leqslant\varphi,\theta} = \mathscr{L}_\theta$,
  - if $0 <_\theta \varphi <_\theta \varphi_o$, then $\mathscr{L}_{\leqslant\varphi,\theta} = \mathscr{L}_{\leqslant 0,\theta}$,
  - if $\varphi <_\theta 0 <_\theta \varphi_o$, then $\mathscr{L}_{\leqslant\varphi,\theta} = 0$.

If $\varphi_o$ and 0 are not comparable at $\theta$, then one argues similarly to determine $\mathscr{L}_{\leqslant\varphi,\theta}$.

***Definition 2.12*.** — A *morphism* $\lambda : (\mathscr{L}, \mathscr{L}_\bullet) \to (\mathscr{L}', \mathscr{L}'_\bullet)$ of non-ramified $\boldsymbol{k}$-Stokes-filtered local systems is a morphism of local systems $\mathscr{L} \to \mathscr{L}'$ on $S^1$ such that, for any $\varphi \in \mathcal{P}$, $\lambda(\mathscr{L}_{\leqslant\varphi}) \subset \mathscr{L}'_{\leqslant\varphi}$, hence, according to (2.2), a morphism also satisfies $\lambda(\mathscr{L}_{<\varphi}) \subset \mathscr{L}'_{<\varphi}$. A morphism $\lambda$ is said to be *strict* if, for any $\varphi$, $\lambda(\mathscr{L}_{\leqslant\varphi}) = \lambda(\mathscr{L}) \cap \mathscr{L}'_{\leqslant\varphi}$.

***Definition 2.13*.** — Given two non-ramified $\boldsymbol{k}$-Stokes-filtered local systems $(\mathscr{L}, \mathscr{L}_\bullet)$ and $(\mathscr{L}', \mathscr{L}'_\bullet)$,

- the direct sum $(\mathscr{L}, \mathscr{L}_\bullet) \oplus (\mathscr{L}', \mathscr{L}'_\bullet)$ has local system $\mathscr{L} \oplus \mathscr{L}'$ and filtration $(\mathscr{L} \oplus \mathscr{L}')_{\leqslant\varphi} = \mathscr{L}_{\leqslant\varphi} \oplus \mathscr{L}'_{\leqslant\varphi}$,
- $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_{\leqslant\eta}$ is the subsheaf of $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')$ consisting of local morphisms $\mathscr{L} \to \mathscr{L}'$ sending $\mathscr{L}_{\leqslant\varphi}$ into $\mathscr{L}'_{\leqslant\varphi+\eta}$ fo any $\varphi$;
- in particular, $(\mathscr{L}^\vee, \mathscr{L}^\vee_\bullet) = (\mathscr{H}\!om(\mathscr{L}, \boldsymbol{k}_{S^1}), \mathscr{H}\!om(\mathscr{L}, \boldsymbol{k}_{S^1})_\bullet)$, where $\boldsymbol{k}_{S^1}$ is equipped with the graded Stokes filtration of Example 2.11(1).
- $(\mathscr{L} \otimes \mathscr{L}')_{\leqslant\eta} = \sum_\varphi \mathscr{L}_{\leqslant\varphi} \otimes \mathscr{L}'_{\leqslant\eta-\varphi} \subset \mathscr{L} \otimes \mathscr{L}'$.

In particular, a morphism of non-ramified Stokes-filtered local systems is a global section of $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_{\leqslant 0}$.

***Example 2.14* (Twist).** — Let $\boldsymbol{k}_{S^1}[\eta]_\bullet$ be the graded Stokes filtration twisted by $\eta$ (see Example 2.11(2) with $\mathscr{L} = \boldsymbol{k}_{S^1}$). Then we have $(\mathscr{L} \otimes \boldsymbol{k}_{S^1}[\eta])_\bullet = \mathscr{L}[\eta]_\bullet$.

***Proposition 2.15*.** — *Given two non-ramified $\boldsymbol{k}$-Stokes filtrations $\mathscr{L}_\bullet, \mathscr{L}'_\bullet$ of $\mathscr{L}, \mathscr{L}'$, $\mathscr{L}_\bullet \oplus \mathscr{L}'_\bullet$, $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_\bullet$, $(\mathscr{L}^\vee)_\bullet$ and $(\mathscr{L} \otimes \mathscr{L}')_\bullet$ are non-ramified $\boldsymbol{k}$-Stokes filtrations of the corresponding local systems and $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_\bullet \simeq (\mathscr{L}'^\vee \otimes \mathscr{L})_\bullet$. Moreover,*

(1) *$\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_{<\eta}$ is the subsheaf of $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')$ consisting of local morphisms $\mathscr{L} \to \mathscr{L}'$ sending $\mathscr{L}_{\leqslant\varphi}$ into $\mathscr{L}'_{<\varphi+\eta}$ fo any $\varphi$;*

(2) *$(\mathscr{L}^\vee)_{\leqslant\varphi} = (\mathscr{L}_{<-\varphi})^\perp$ and $(\mathscr{L}^\vee)_{<\varphi} = (\mathscr{L}_{\leqslant-\varphi})^\perp$ for any $\varphi$, so that $\mathrm{gr}_\varphi \mathscr{L}^\vee = (\mathrm{gr}_{-\varphi} \mathscr{L})^\vee$ (here, $(\mathscr{L}_{<-\varphi})^\perp$, resp. $(\mathscr{L}_{\leqslant-\varphi})^\perp$, consists of local morphisms $\mathscr{L} \to \boldsymbol{k}_{S^1}$ sending $\mathscr{L}_{<-\varphi}$, resp. $\mathscr{L}_{\leqslant-\varphi}$, to zero);*

(3) *$(\mathscr{L} \otimes \mathscr{L}')_{<\eta} = \sum_\varphi \mathscr{L}_{\leqslant\varphi} \otimes \mathscr{L}'_{<\eta-\varphi} = \sum_\varphi \mathscr{L}_{<\varphi} \otimes \mathscr{L}'_{\leqslant\eta-\varphi}$.*

*Proof*. — For the first assertion, let us consider the case of $\mathscr{H}\!om$ for instance. Using a local decomposition of $\mathscr{L}, \mathscr{L}'$ given by the $\mathcal{I}$-filtration condition, we find that a section of $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_\theta$ is decomposed as a section of $\bigoplus_{\varphi,\psi} \mathscr{H}\!om(\mathrm{gr}_\varphi \mathscr{L}, \mathrm{gr}_\psi \mathscr{L}')_\theta$, and that it belongs to $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_{\leqslant\eta,\theta}$ if and only if its components $(\varphi, \psi)$ are zero whenever $\psi - \varphi \not\leqslant_\theta \eta$. The assertion is then clear, as well as the characterization of $\mathscr{H}\!om(\mathscr{L}, \mathscr{L}')_{<\eta}$.

As a consequence, a local section of $(\mathscr{L}^\vee)_{\leqslant\varphi}$ has to send $\mathscr{L}_{<-\varphi}$ to zero for any $\varphi$. The converse is also clear by using the local decomposition of $(\mathscr{L}, \mathscr{L}_\bullet)$, as well as the other assertions for $\mathscr{L}^\vee$.

The assertion on the tensor product is then routine. ☐

***Remark 2.16.*** — One easily gets the behaviour of the set of exponential factors with respect to such operations. For instance, the direct sum corresponds to $(\Phi,\Phi')\mapsto\Phi\cup\Phi'$, the dual to $\Phi\mapsto-\Phi$ and the tensor product to $(\Phi,\Phi')\mapsto\Phi+\Phi'$.

*Poincaré-Verdier duality.* — For a sheaf $\mathscr{F}$ on $S^1$, its Poincaré-Verdier dual $\boldsymbol{D}\mathscr{F}$ is $\boldsymbol{R}\mathscr{H}\!om_{\mathbb{C}}(\mathscr{F},\boldsymbol{k}_{S^1}[1])$ and we denote by $\boldsymbol{D}'\mathscr{F}=\boldsymbol{R}\mathscr{H}\!om_{\mathbb{C}}(\mathscr{F},\boldsymbol{k}_{S^1})$ the shifted complex. We clearly have $\boldsymbol{D}'\mathscr{L}=\mathscr{H}\!om_{\mathbb{C}}(\mathscr{L},\boldsymbol{k}_{S^1})=:\mathscr{L}^\vee$.

***Lemma 2.17* (Poincaré duality).** — *For any $\varphi\in\mathcal{P}$, the complexes $\boldsymbol{D}'(\mathscr{L}_{\leqslant\varphi})$ and $\boldsymbol{D}'(\mathscr{L}/\mathscr{L}_{\leqslant\varphi})$ are sheaves. Moreover, the two subsheaves $(\mathscr{L}^\vee)_{\leqslant\varphi}$ and $\boldsymbol{D}'(\mathscr{L}/\mathscr{L}_{<-\varphi})$ of $\mathscr{L}^\vee$ coincide.*

*Proof.* — The assertions are local on $S^1$, so we can assume that $(\mathscr{L},\mathscr{L}_\bullet)$ is split with respect to the Stokes filtration, and therefore that $(\mathscr{L},\mathscr{L}_\bullet)$ has only one exponential factor $\eta$, that is, $\mathscr{L}_{\leqslant\varphi}=\beta_{\eta\leqslant\varphi}\mathscr{L}$. Let us denote by $\alpha_{\psi<\varphi}$ the functor composed of the restriction to the open set $S^1_{\psi<\varphi}$ (see Notation 2.3) and the maximal extension to $S^1$, and similarly with $\leqslant$. We have an exact sequence

$$0\longrightarrow\beta_{\eta<\varphi}\mathscr{L}\longrightarrow\mathscr{L}\longrightarrow\alpha_{\varphi\leqslant\eta}\mathscr{L}\longrightarrow0$$

which identifies $\alpha_{\varphi\leqslant\eta}\mathscr{L}$ to $\mathscr{L}/\mathscr{L}_{<\varphi}$, and a similar one with $\beta_{\eta\leqslant\varphi}$ and $\alpha_{\varphi<\eta}$. On the other hand, $\boldsymbol{D}'(\beta_{\eta\leqslant\varphi}\mathscr{L})=\alpha_{\eta\leqslant\varphi}\mathscr{L}^\vee$ and $\boldsymbol{D}'(\alpha_{\varphi\leqslant\eta}\mathscr{L})=\beta_{\varphi\leqslant\eta}\mathscr{L}^\vee$. The dual of the previous exact sequence, when we replace $\varphi$ with $-\varphi$, is then

$$0\longrightarrow\beta_{-\varphi\leqslant\eta}\mathscr{L}^\vee\longrightarrow\mathscr{L}^\vee\longrightarrow\alpha_{\eta<-\varphi}\mathscr{L}^\vee\longrightarrow0,$$

also written as

$$0\longrightarrow\beta_{-\eta\leqslant\varphi}\mathscr{L}^\vee\longrightarrow\mathscr{L}^\vee\longrightarrow\alpha_{\varphi<-\eta}\mathscr{L}^\vee\longrightarrow0,$$

showing that $\boldsymbol{D}'(\mathscr{L}/\mathscr{L}_{<-\varphi})=(\mathscr{L}^\vee)_{\leqslant\varphi}$. ☐

**2.b. Pull-back and push-forward.** — Let $f:X'\to X$ be a holomorphic map from the disc $X'$ (with coordinate $x'$) to the disc $X$. We assume that both discs are small enough so that $f$ is ramified at $x'=0$ only. We now denote by $S^1_{x'}$ and $S^1_x$ the circles of directions in $\widetilde{X}'$ and $\widetilde{X}$ respectively. Then $f$ induces $\widetilde{f}:S^1_{x'}\to S^1_x$, which is the composition of the multiplication by $N$ (the index of ramification of $f$) and a translation (the argument of $f^{(N)}(0)$). Similarly, $\mathcal{P}_x$ and $\mathcal{P}_{x'}$ denote the polar parts in the variables $x$ and $x'$ respectively.

***Remark 2.18.*** — Let $\eta\in\mathcal{P}_x$ and set $f^*\eta=\eta\circ f\in\mathcal{P}_{x'}$. For any $\theta'\in S^1_{x'}$, set $\theta=\widetilde{f}(\theta')$. Then we have

$$f^*\eta\leqslant_{\theta'}0\Longleftrightarrow\eta\leqslant_\theta0\quad\text{and}\quad f^*\eta<_{\theta'}0\Longleftrightarrow\eta<_\theta0.$$

(This is easily seen using the definition in terms of moderate growth, since $\widetilde{f}:\widetilde{X}'\to\widetilde{X}$ is a finite covering.)

***Definition 2.19* (Pull-back).** — Let $\mathscr{L}$ be a local system on $S^1_x$ and let $\mathscr{L}_\bullet$ be a non-ramified $\boldsymbol{k}$-pre-Stokes filtration on $\mathscr{L}$. For any $\varphi' \in \mathcal{P}_{x'}$ and any $\theta' \in S^1_{x'}$, let us set

$$(\widetilde{f}^{-1}\mathscr{L})_{\leqslant\varphi',\theta'} := \sum_{\substack{\psi\in\mathcal{P}_x\\ f^*\psi\leqslant_{\theta'}\varphi'}} \mathscr{L}_{\leqslant\psi,\widetilde{f}(\theta')} \subset \mathscr{L}_{\widetilde{f}(\theta')} = (\widetilde{f}^{-1}\mathscr{L})_{\theta'}.$$

Then $(\widetilde{f}^{-1}\mathscr{L})_\bullet$ is a non-ramified pre-Stokes filtration on $\widetilde{f}^{-1}\mathscr{L}$, called the *pull-back* of $\mathscr{L}_\bullet$ by $f$.

***Proposition 2.20* (Pull-back).** — *The pull-back has the following properties:*

(1) *For any $\varphi' \in \mathcal{P}_{x'}$ and any $\theta' \in S^1_{x'}$,*

$$(\widetilde{f}^{-1}\mathscr{L})_{<\varphi',\theta'} = \sum_{\substack{\psi\in\mathcal{P}_x\\ f^*\psi<_{\theta'}\varphi'}} \widetilde{f}^{-1}(\mathscr{L}_{\leqslant\psi,\widetilde{f}(\theta')}).$$

(2) *For any $\varphi \in \mathcal{P}_x$,*

$$\begin{aligned}(\widetilde{f}^{-1}\mathscr{L})_{\leqslant f^*\varphi} &= \widetilde{f}^{-1}(\mathscr{L}_{\leqslant\varphi}),\\ (\widetilde{f}^{-1}\mathscr{L})_{<f^*\varphi} &= \widetilde{f}^{-1}(\mathscr{L}_{<\varphi})\end{aligned}$$

and

$$\mathrm{gr}_{f^*\varphi}(\widetilde{f}^{-1}\mathscr{L}) = \widetilde{f}^{-1}(\mathrm{gr}_\varphi\mathscr{L}).$$

(3) *In particular, if $(\widetilde{f}^{-1}\mathscr{L})_\bullet$ is a non-ramified Stokes filtration for some $f$, then for any $\varphi \in \mathcal{P}_x$, $\mathrm{gr}_\varphi\mathscr{L}$ is a local system on $S^1_x$.*

(4) *Let $\mathscr{L},\mathscr{L}'$ be two local systems on $S^1_x$ equipped with non-ramified pre-Stokes filtrations and let $\lambda : \mathscr{L} \to \mathscr{L}'$ be a morphism of local systems such that, for some $f$, $\widetilde{f}^{-1}\lambda : \widetilde{f}^{-1}\mathscr{L} \to \widetilde{f}^{-1}\mathscr{L}'$ is compatible with the non-ramified pre-Stokes filtrations $(\widetilde{f}^{-1}\mathscr{L})_\bullet, (\widetilde{f}^{-1}\mathscr{L}')_\bullet$. Then $\lambda$ is compatible with the non-ramified pre-Stokes filtrations $\mathscr{L}_\bullet, \mathscr{L}'_\bullet$.*

(5) *Assume now that $\mathscr{L}_\bullet$ is a non-ramified $\boldsymbol{k}$-Stokes filtration (i.e., is locally graded) and let $\Phi$ be its set of exponential factors. Then $(\widetilde{f}^{-1}\mathscr{L})_\bullet$ is a non-ramified $\boldsymbol{k}$-Stokes filtration on $\widetilde{f}^{-1}\mathscr{L}$ and, for any $\varphi' \in \mathcal{P}_{x'}$, $\mathrm{gr}_{\varphi'}\widetilde{f}^{-1}\mathscr{L} \neq 0 \Rightarrow \varphi' \in f^*\Phi$.*

(6) *The pull-back of non-ramified Stokes filtrations is compatible with $\mathscr{H}om$, duality and tensor product.*

*Proof.* — By definition,

$$(\widetilde{f}^{-1}\mathscr{L})_{<\varphi',\theta'} = \sum_{\psi'<_{\theta'}\varphi'} (\widetilde{f}^{-1}\mathscr{L})_{\leqslant\psi',\theta'} = \sum_{\psi'<_{\theta'}\varphi'} \sum_{\substack{\psi\in\mathcal{P}_x\\ f^*\psi\leqslant_{\theta'}\psi'}} \mathscr{L}_{\leqslant\psi,\widetilde{f}(\theta')},$$

and this is the RHS in 2.20(1).

The first two lines of 2.20(2) are a direct consequence of Remark 2.18, and the third one is a consequence of the previous ones. Then 2.20(3) follows, as each $\mathrm{gr}_{\varphi'}(\widetilde{f}^{-1}\mathscr{L})$ is a local system on $S^1_{x'}$.

2.20(5) follows from the third line in 2.20(2) and the local gradedness condition. Then, 2.20(6) is clear. ☐

***Remark 2.21.*** — In order to make clear the correspondence with the notion introduced in Definition 1.34 and considered in Lemma 1.41, note that the sheaf $\mathcal{I}$ is the constant sheaf on $S^1_x$ with fibre $\mathcal{P}_x$, $\widetilde{f}^{-1}\mathcal{I}$ is the constant sheaf on $S^1_{x'}$ with fibre $\mathcal{P}_x$ and $\mathcal{I}'$ is the constant sheaf on $S^1_{x'}$ with fibre $\mathcal{P}_{x'}$. The map $q_f$ is $f^*:\mathcal{P}_x\to\mathcal{P}_{x'}$.

***Exercise 2.22* (Push forward).** — Let $\mathscr{L}'$ be a local system on $S^1_{x'}$ equipped with a non-ramified pre-Stokes filtration $\mathscr{L}'_\bullet$. Show that

(1) $\widetilde{f}_*\mathscr{L}'$ is naturally equipped with a non-ramified Stokes filtration defined by $(\widetilde{f}_*\mathscr{L}')_{\leqslant\varphi}=\widetilde{f}_*(\mathscr{L}'_{\leqslant f^*\varphi})$.

(2) Let $\mathscr{L}'$ be a local system on $S^1_{x'}$ equipped with a non-ramified Stokes filtration $\mathscr{L}'_\bullet$ and let $\Phi'\subset\mathcal{P}_{x'}$ be its set of exponential factors. If there exists a finite subset $\Phi\subset\mathcal{P}_x$ such that $\Phi'=f^*\Phi$, then the push-forward pre-Stokes filtration $(\widetilde{f}_*\mathscr{L}')_\bullet$ is a Stokes filtration.

**2.c. Stokes filtrations on local systems.** — We now define the general notion of a (possibly ramified) Stokes filtration on a local system $\mathscr{L}$ on $S^1$.

Let $d$ be a nonzero integer and let $\rho_d:X_d\to X$ be a holomorphic function from a disc $X_d$ (coordinate $x'$) to the disc $X$ (coordinate $x$). For simplicity, we will assume that the coordinates are chosen so that $\rho_d(x')=x'^d$. We denote by $\mathcal{I}_d$ the local system on $S^1_x$ whose fibre at $\theta=0$ is $\mathcal{P}_{x'}$ and whose monodromy is given by $\mathcal{P}_{x'}\ni\varphi'(x')\mapsto\varphi'(e^{2\pi i/d}x')$. If we denote by $\widetilde{\rho}_d:S^1_{x'}\to S^1_x$, $\theta'\mapsto d\theta'$ the associated map, the sheaf $\widetilde{\rho}_d^{-1}\mathcal{I}_d$ is the constant sheaf on $S^1_{x'}$ with fibre $\mathcal{P}_{x'}$. In particular, $\mathcal{I}_d$ is a sheaf of ordered abelian groups on $S^1_x$, and the constant sheaf with fibre $\mathcal{P}_x$, that we now denote by $\mathcal{I}_1$, is a subsheaf of ordered abelian groups. We will then denote by $\mathcal{I}$ the sheaf $\bigcup_{d\geqslant1}\mathcal{I}_d$.

***Remark 2.23.*** — Let us give another description of the sheaf $\mathcal{I}_d$ which will be useful in higher dimensions. Let us denote by $j_\partial$ and $j_{\partial,d}$ the natural inclusions $X^*\hookrightarrow\widetilde{X}$ and $X^*\hookrightarrow\widetilde{X}_d$. The natural inclusion $\mathscr{O}_{X^*}\hookrightarrow\rho_{d,*}\mathscr{O}_{X_d^*}$ induces an injective morphism $j_{\partial,*}\mathscr{O}_{X^*}\hookrightarrow j_{\partial,*}\rho_{d,*}\mathscr{O}_{X_d^*}=\widetilde{\rho}_{d,*}j_{\partial,d,*}\mathscr{O}_{X_d^*}$, that we regard as the inclusion of a subsheaf.

Let us denote by $(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$ the subsheaf of $j_{\partial,*}\mathscr{O}_{X^*}$ consisting of functions which are locally bounded on $\widetilde{X}$. We have $(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}=\widetilde{\rho}_{d,*}(j_{\partial,d,*}\mathscr{O}_{X_d^*})^{\mathrm{lb}}\cap j_{\partial,*}\mathscr{O}_{X^*}$ since $\widetilde{\rho}_d$ is proper.

Let us set $\widetilde{\mathcal{I}}_1=\varpi^{-1}\mathscr{O}_X(*0)$, that we consider as a subsheaf of $j_{\partial,*}\mathscr{O}_{X^*}$. We have $\varpi^{-1}\mathscr{O}_X=\widetilde{\mathcal{I}}_1\cap(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$ in $j_{\partial,*}\mathscr{O}_{X^*}$ (a meromorphic function which is bounded in some sector is bounded everywhere, hence is holomorphic). Therefore, $\mathcal{I}_1:=\varpi^{-1}(\mathscr{O}_X(*0)/\mathscr{O}_X)$ is also equal to $\widetilde{\mathcal{I}}_1/\widetilde{\mathcal{I}}_1\cap(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$.

Similarly, we can first define $\widetilde{\mathcal{I}}_d$ as the subsheaf of $\mathbb{C}$-vector spaces of $j_{\partial,*}\mathscr{O}_{X^*}$ which is the intersection of $\widetilde{\rho}_{d,*}\varpi_d^{-1}\mathscr{O}_{X_d}(*0)$ and $j_{\partial,*}\mathscr{O}_{X^*}$ in $\widetilde{\rho}_{d,*}j_{\partial,d,*}\mathscr{O}_{X_d^*}$. We then set $\mathcal{I}_d = \widetilde{\mathcal{I}}_d/\widetilde{\mathcal{I}}_d \cap (j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$, which is a subsheaf of $j_{\partial,*}\mathscr{O}_{X^*}/(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$. We have $\mathcal{I}_d \subset \mathcal{I}_{d'}$ if $d$ divides $d'$.

As we already noticed, $\mathcal{I}_1 = \varpi^{-1}(\mathscr{O}_X(*0)/\mathscr{O}_X)$. More generally, let us show that $\widetilde{\rho}_d^{-1}\mathcal{I}_d = \varpi_d^{-1}(\mathscr{O}_{X_d}(*0)/\mathscr{O}_{X_d})$. We will start by showing that $\widetilde{\rho}_d^{-1}\widetilde{\mathcal{I}}_d = \varpi_d^{-1}\mathscr{O}_{X_d}(*0)$.

Let us first note that $\rho_d^{-1}\mathscr{O}_{X^*} = \mathscr{O}_{X_d^*}$ since $\rho_d$ is a covering, and $\widetilde{\rho}_d^{-1}j_{\partial,*}\mathscr{O}_{X^*} = j_{\partial,d,*}\rho_d^{-1}\mathscr{O}_{X^*}$ since $\widetilde{\rho}_d$ is a covering. Hence, $\widetilde{\rho}_d^{-1}j_{\partial,*}\mathscr{O}_{X^*} = j_{\partial,d,*}\mathscr{O}_{X_d^*}$.

It follows that $\widetilde{\rho}_d^{-1}\widetilde{\mathcal{I}}_d$ is equal to the intersection of $\widetilde{\rho}_d^{-1}\widetilde{\rho}_{d,*}[\varpi_d^{-1}\mathscr{O}_{X_d}(*0)]$ (since $\widetilde{\mathcal{I}}_{\widetilde{X}_d,1} = \varpi_d^{-1}\mathscr{O}_{X_d}(*0)$) and $j_{\partial,d,*}\mathscr{O}_{X_d^*}$ in $\widetilde{\rho}_d^{-1}\widetilde{\rho}_{d,*}[j_{\partial,d,*}\mathscr{O}_{X_d^*}]$. This is $\varpi_d^{-1}\mathscr{O}_{X_d}(*0)$. Indeed, a germ in $\widetilde{\rho}_d^{-1}\widetilde{\rho}_{d,*}[\varpi_d^{-1}\mathscr{O}_{X_d}(*0)]$ at $\theta'$ consists of a $d$-uple of germs in $\mathscr{O}_{X_d}(*0)$ at 0. This $d$-uple belongs to $j_{\partial,d,*}\mathscr{O}_{X_d^*,\theta'}$ iff the restrictions to $X_d^*$ of the terms of the $d$-uple coincide. Then all the terms of the $d$-uple are equal. The argument for $\mathcal{I}_d$ is similar.

Let us express these results in terms of étale spaces. We first note that, since $\mathcal{I}_1$ is a constant sheaf, the étale space $\mathcal{I}_1^{\text{ét}}$ is a trivial covering of $S^1_x$. The previous argument shows that the fibre product $S^1_{x'} \times_{S^1_x} \mathcal{I}_d^{\text{ét}}$ is identified with $\mathcal{I}^{\text{ét}}_{x',1}$, hence is a trivial covering of $S^1_{x'}$. It follows that, since $\widetilde{\rho}_d : S^1_{x'} \times_{S^1_x} \mathcal{I}_d^{\text{ét}} \to \mathcal{I}_d^{\text{ét}}$ is a finite covering of degree $d$, that $\mathcal{I}_d^{\text{ét}} \to S^1_x$ is a covering.

The following property will also be useful: there is a one-to-one correspondence between finite sets $\Phi_d$ of $\mathcal{P}_{x'}$ and finite coverings $\widetilde{\Sigma} \subset \mathcal{I}_d^{\text{ét}}$. Indeed, given such a $\widetilde{\Sigma}$, its pull-back $\widetilde{\Sigma}_d$ by $\widetilde{\rho}_d$ is a covering of $S^1_{x'}$ contained in the trivial covering $\widetilde{\rho}_d^{-1}\mathcal{I}_d^{\text{ét}}$, hence is trivial, and is determined by its fibre $\Phi_d \subset \mathcal{P}_{x'}$. Conversely, given such $\Phi_d$, it defines a trivial covering $\widetilde{\Sigma}_d$ of $S^1_{x'}$ contained in $\widetilde{\rho}_d^{-1}\mathcal{I}_d^{\text{ét}}$. Let $\widetilde{\Sigma}$ be its image in $\mathcal{I}_d^{\text{ét}}$. Because the composed map $\widetilde{\Sigma}_d \to S^1_{x'} \to S^1_x$ is a covering, so are both maps $\widetilde{\Sigma}_d \to \widetilde{\Sigma}$ and $\widetilde{\Sigma} \to S^1_x$. Moreover, the degree of $\widetilde{\Sigma} \to S^1_x$ is equal to that of $\widetilde{\Sigma}_d \to S^1_{x'}$, that is, $\#\Phi_d$. Lastly, the pull-back of $\widetilde{\Sigma}$ by $\widetilde{\rho}_d$ is a covering of $S^1_{x'}$ contained in $S^1_{x'} \times_{S^1_x} \mathcal{I}_d^{\text{ét}}$, hence is a trivial covering, of degree $\#\Phi_d$, and containing $\widetilde{\Sigma}_d$, so is equal to $\widetilde{\Sigma}_d$.

*Order.* — The sheaf $j_{\partial,*}\mathscr{O}_{X^*}$ is naturally ordered by defining $(j_{\partial,*}\mathscr{O}_{X^*})_{\leqslant 0}$ as the subsheaf of $j_{\partial,*}\mathscr{O}_{X^*}$ whose sections have an exponential with moderate growth along $S^1_x$. Similarly, $j_{\partial,d,*}\mathscr{O}_{X_d^*}$ is ordered. In this way, $\widetilde{\mathcal{I}}$ inherits an order: $\widetilde{\mathcal{I}}_{\leqslant 0} = \mathcal{I} \cap (j_{\partial,*}\mathscr{O}_{X^*})_{\leqslant 0}$. This order is not altered by adding a local section of $(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$, and thus defines an order on $\mathcal{I}$. For each $d$, we also have $\widetilde{\mathcal{I}}_{d,\leqslant 0} = \widetilde{\rho}_{d,*}\big((\varpi_d^{-1}\mathscr{O}_{X_d}(*0))_{\leqslant 0}\big) \cap j_{\partial,*}\mathscr{O}_{X^*}$ and we also conclude that $\widetilde{\rho}_d^{-1}(\mathcal{I}_{d,\leqslant 0}) = \big(\varpi_d^{-1}(\mathscr{O}_{X_d}(*0)/\mathscr{O}_{X_d})\big)_{\leqslant 0}$.

***Definition 2.24*** **((Pre-)Stokes filtration).** — Let $\mathscr{L}$ be a local system on $S^1_x$. A $\boldsymbol{k}$*-(pre-) Stokes filtration* on $\mathscr{L}$ is a (pre-)$\mathcal{I}$-filtration on $\mathscr{L}$. We say that it is ramified of order $\leqslant d$ if it is defined on $\mathcal{I}_d$.

***Remarks 2.25***

(1) One can rephrase the definition in the following way: a Stokes filtration (ramified of order $\leqslant d$) on $\mathscr{L}$ consists of a non-ramified Stokes filtration on $\mathscr{L}' := \rho_d^{-1}\mathscr{L}$

such that, for any automorphism $\sigma$

$$\begin{tikzcd} X_d \arrow[rr, "\sigma"] \arrow[dr, "\rho_d"'] & & X_d \arrow[dl, "\rho_d"] \\ & X & \end{tikzcd}$$

and any $\varphi' \in \mathcal{P}_{x'}$, we have $\mathscr{L}'_{\leqslant\sigma^*\varphi'} = \widetilde{\sigma}^{-1}\mathscr{L}'_{\leqslant\varphi'}$ in $\mathscr{L}' = \widetilde{\sigma}^{-1}\mathscr{L}'$. Similarly, a morphism of $\boldsymbol{k}$-Stokes-filtered local systems is a morphism of local systems which becomes a morphism of non-ramified Stokes-filtered local systems after ramification.

(2) The previous condition can be restated by saying that, for any $\sigma$, the Stokes-filtered local system $(\mathscr{L}', \mathscr{L}'_\bullet)$ and its pull-back by $\widetilde{\sigma}$ coincide (owing to the natural identification $\mathscr{L}' = \widetilde{\sigma}^{-1}\mathscr{L}'$).

(3) Restricting the Stokes-filtered local system to $\mathcal{I}_1^{\text{ét}}$ defines a non-ramified pre-Stokes filtration on $\mathscr{L}$. Note also that a morphism of Stokes-filtered local systems is compatible with this pre-Stokes filtration. Hence the category of Stokes filtrations on $\mathscr{L}$ is a subcategory of the category of pre-Stokes filtrations on $\mathscr{L}$.

(4) Notice however that, for $\varphi \in \mathcal{P}_x$, $\mathscr{L}_{<\varphi}$ is not computed by Formula (2.2) with $\psi \in \mathcal{P}_x$: one has to consider all $\psi' \in \mathcal{P}_{x'}$. Of course, for such a $\varphi$, $\operatorname{gr}_\varphi \mathscr{L}$ is a local system on $S^1_x$.

(5) The "set of exponential factors of the Stokes-filtered local system" is now replaced by a subset $\widetilde{\Sigma} \subset \mathcal{I}^{\text{ét}}$ such that the projection to $\partial\widetilde{X}$ is a finite covering. It corresponds to a finite subset $\Phi_d \subset \mathcal{P}_{x'}$ for a suitable ramified covering $\rho_d$ (see the last part of Remark 2.23), which is the set of exponential factors of the non-ramified Stokes filtration $\widetilde{\rho}_d^{-1}\mathscr{L}$ considered in (2).

(6) The category of Stokes-filtered local systems $(\mathscr{L}, \mathscr{L}_\bullet)$ with associated covering contained in $\widetilde{\Sigma}$ is equivalent to the category of $\widetilde{\Sigma}$-filtered local systems (see Remark 2.10).

(7) Proposition 2.15 holds for $\boldsymbol{k}$-Stokes filtrations.

(8) Lemma 2.17 holds for $\boldsymbol{k}$-Stokes filtrations, that is, the family $\boldsymbol{D}'(\mathscr{L}/\mathscr{L}_{\leqslant -\varphi})$ of local subsheaves of $\mathscr{L}^\vee$ indexed by local sections of $\mathcal{I}$ forms a Stokes filtration of $\mathscr{L}^\vee$.

(9) The category of non-ramified $\boldsymbol{k}$-Stokes-filtered local systems on $S^1_x$ is a full subcategory of that of $\boldsymbol{k}$-Stokes-filtered local systems. Indeed, given a non-ramified Stokes-filtered local system on $S^1_x$, one extends it as a ramified Stokes-filtered local system of order $d$ from $\mathcal{I}_1^{\text{ét}}$ to $\mathcal{I}_d^{\text{ét}}$ using a formula analogous to that of Proposition 2.20(1).

(10) If the set $\Phi_d$ of exponential factors of $\mathscr{L}_\bullet$ takes the form $\rho_d^*\Phi$ for some finite subset $\Phi \subset \mathcal{P}_x$ (equivalently, the finite covering $\widetilde{\Sigma}$ of $\partial\widetilde{X}$ is trivial, see 2.25(5)), then the Stokes filtration is non-ramified.

(11) We will still denote by $\mathscr{L}_\bullet$ a (possibly ramified) Stokes filtration on $\mathscr{L}$.

**2.d. Extension of scalars.** — Let $(\mathscr{L},\mathscr{L}_\bullet)$ be a Stokes-filtered local system defined over a field $\boldsymbol{k}$ and let $\boldsymbol{k}'$ be an extension of $\boldsymbol{k}$. Then $(\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L},\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_\bullet)$ is a Stokes-filtered local system defined over $\boldsymbol{k}'$. The following properties are satisfied for any local section $\varphi$ of $\mathcal{I}$:

- $(\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L})_{<\varphi}=\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_{<\varphi}$, and $\operatorname{gr}_\varphi(\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L})=\boldsymbol{k}'\otimes\operatorname{gr}_\varphi\mathscr{L}$, so the set of exponential factors of $(\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L},\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_\bullet)$ is equal to that of $(\mathscr{L},\mathscr{L}_\bullet)$;
- $\mathscr{L}_{\leqslant\varphi}=(\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_{\leqslant\varphi})\cap\mathscr{L}$ in $\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}$.

In such a case, we say that the Stoles-filtered local system $(\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L},\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_\bullet)$ is *defined over* $\boldsymbol{k}$.

Conversely, let now $(\mathscr{L},\mathscr{L}_\bullet)$ be a Stokes-filtered local system over $\boldsymbol{k}'$ and let $\widetilde{\Sigma}\subset\mathcal{I}$ be its covering of exponential factors. We wish to find sufficient conditions to ensure that it comes from a Stokes-filtered local system defined over $\boldsymbol{k}$ by extension of scalars.

**Proposition 2.26.** — *Assume that the local system $\mathscr{L}$ is defined over $\boldsymbol{k}$, that is, $\mathscr{L}=\boldsymbol{k}'\otimes\mathscr{L}_{\boldsymbol{k}}$ for some $\boldsymbol{k}$-local system $\mathscr{L}_{\boldsymbol{k}}$ (regarded as a subsheaf of $\mathscr{L}$), and that, for any local section $\varphi$ of $\widetilde{\Sigma}$,*

$$\mathscr{L}_{\leqslant\varphi}=\boldsymbol{k}'\otimes_{\boldsymbol{k}}(\mathscr{L}_{\boldsymbol{k}}\cap\mathscr{L}_{\leqslant\varphi}),$$

*where the intersection is taken in $\mathscr{L}$. Then $(\mathscr{L},\mathscr{L}_\bullet)$ is a Stokes-filtered local system defined over $\boldsymbol{k}$.*

*Proof.* — It is not difficult to reduce to the non-ramified case, so we will assume below that $\mathcal{I}=\mathcal{I}_1$ and replace $\widetilde{\Sigma}$ by $\Phi\subset\mathcal{P}_x$. We set, for any local section $\psi$ of $\mathcal{P}_x$, $\mathscr{L}_{\boldsymbol{k},\leqslant\psi}:=\mathscr{L}_{\boldsymbol{k}}\cap\mathscr{L}_{\leqslant\psi}$, so that the condition reads $\mathscr{L}_{\leqslant\varphi}=\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_{\boldsymbol{k},\leqslant\varphi}$ for $\varphi\in\Phi$. This defines a pre-$\mathcal{I}$-filtration of $\mathscr{L}_{\boldsymbol{k}}$, and we will show that this is indeed a $\mathcal{I}$-filtration.

(1) We start with a general property of Stokes-filtered local systems. Let $(\mathscr{L},\mathscr{L}_\bullet)$ and $\Phi$ be as above, and let $\psi\in\mathcal{P}_x$. Set $\Psi=\Phi\cup\{\psi\}$ and denote by $\operatorname{St}(\Psi,\Psi)$ the (finite) set of Stokes directions of pairs $\varphi,\eta\in\Psi$. The sheaves $\mathscr{L}_{\leqslant\psi}$ and $\mathscr{L}_{<\psi}$ can be described as triples consisting of their restrictions to the open set $S^1\smallsetminus\operatorname{St}(\Psi,\Psi)$, the closed set $\operatorname{St}(\Psi,\Psi)$, and a gluing map from the latter to the restriction to this closed set of the push-forward of the former. We will make this description explicit.

On any connected component $I$ of $S^1\smallsetminus\operatorname{St}(\Psi,\Psi)$, the set $\Psi$ is totally ordered, and there exists $\varphi=\varphi(I,\psi)\in\Phi$ such that $\mathscr{L}_{\leqslant\psi|I}=\mathscr{L}_{\leqslant\varphi|I}$. Similarly, there exists $\eta=\eta(I,\psi)\in\Phi$ such that $\mathscr{L}_{<\psi|I}=\mathscr{L}_{<\eta|I}$.

Let us fix $\theta_o\in\operatorname{St}(\Psi,\Psi)$ and denote by $I_1,I_2$ the two connected components of $S^1\smallsetminus\operatorname{St}(\Psi,\Psi)$ containing $\theta_o$ in their closure, with corresponding inclusions $j_i:I_i\hookrightarrow S^1$, $i=1,2$. We also denote by $i_o:\{\theta_o\}\hookrightarrow S^1$ the closed inclusion and set $\varphi_i:=\varphi(I_i,\psi)$, $i=1,2$ (resp. $\eta_i:=\eta(I_i,\psi)$).

We claim that, in the neighbourhood of $\theta_o$ (and more precisely, on $I_1\cup I_2\cup\{\theta_o\}$), the sheaf $\mathscr{L}_{\leqslant\psi}$ is described by the data $\mathscr{L}_{\leqslant\psi|I_i}=\mathscr{L}_{\leqslant\varphi_i|I_i}$, $i=1,2$, $\mathscr{L}_{\leqslant\psi,\theta_o}=i_o^{-1}j_{1,*}\mathscr{L}_{\leqslant\varphi_1|I_1}\cap i_o^{-1}j_{2,*}\mathscr{L}_{\leqslant\varphi_2|I_2}$, where the intersection is taken in $i_o^{-1}j_{1,*}\mathscr{L}=i_o^{-1}j_{2,*}\mathscr{L}=\mathscr{L}_{\theta_o}$, and the gluing map is the natural inclusion map of the intersection into each of its components. A similar statement holds for $\mathscr{L}_{<\psi}$. This easily follows from the local description $\mathscr{L}_{\leqslant\psi}=\bigoplus_{\varphi\in\Phi}\beta_{\varphi\leqslant\psi}\operatorname{gr}_\varphi\mathscr{L}$, and similarly for $\mathscr{L}_{<\psi}$.

(2) We now claim that $\mathscr{L}_{\boldsymbol{k},\leqslant\psi}:=\mathscr{L}_{\boldsymbol{k}}\cap\mathscr{L}_{\leqslant\psi}$ satisfies $\boldsymbol{k}'\otimes\mathscr{L}_{\boldsymbol{k},\leqslant\psi}=\mathscr{L}_{\leqslant\psi}$. This is by assumption on $S^1\smallsetminus\operatorname{St}(\Psi,\Psi)$, according to the previous description, and it remains to check this at any $\theta_o\in\operatorname{St}(\Psi,\Psi)$. The previous description gives $i_o^{-1}\mathscr{L}_{\boldsymbol{k},\leqslant\psi}=i_o^{-1}j_{1,*}\mathscr{L}_{\boldsymbol{k},\leqslant\varphi_1|I_1}\cap i_o^{-1}j_{2,*}\mathscr{L}_{\boldsymbol{k},\leqslant\varphi_2|I_2}$ and the result follows easily (by considering a suitable basis of $\mathscr{L}_{\boldsymbol{k},\theta_o}$ for instance).

(3) Let us now define $\mathscr{L}_{\boldsymbol{k},<\psi}$ as $\sum_{\psi'\in\mathcal{P}_x}\beta_{\psi'<\psi}\mathscr{L}_{\boldsymbol{k},\psi'}$, as in (2.2). Then the previous description also shows that $\mathscr{L}_{\boldsymbol{k},<\psi}=\mathscr{L}_{\boldsymbol{k}}\cap\mathscr{L}_{<\psi}$ and that $\mathscr{L}_{<\psi}=\boldsymbol{k}'\otimes_{\boldsymbol{k}}\mathscr{L}_{\boldsymbol{k},<\psi}$.

(4) As a consequence, we obtain that $\operatorname{gr}_\psi\mathscr{L}=\boldsymbol{k}'\otimes_{\boldsymbol{k}}\operatorname{gr}_\psi\mathscr{L}_{\boldsymbol{k}}$ for any $\psi$, from which the proposition follows. ☐

***Remark 2.27***. — The condition considered in the proposition is that considered in [**36**] in order to define a $\boldsymbol{k}$-structure on a Stokes-filtered local system defined over $\boldsymbol{k}'$ (e.g. $\boldsymbol{k}=\mathbb{Q}$ and $\boldsymbol{k}'=\mathbb{C}$). This proposition shows that there is no difference with the notion of Stokes-filtered $\boldsymbol{k}$-local system.

**2.e. Stokes-filtered local systems and Stokes data.** — In this section, we make explicit the relationship between Stokes filtrations and the more conventional approach with Stokes data in the simple case of a Stokes-filtered local system of *exponential type.*

*Stokes-filtered local systems of exponential type*

***Definition 2.28* (see [49] and [36]).** — We say that a Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$ is of *exponential type* if it is non-ramified and its exponential factors have a pole of order one at most.

In such a case, we can replace $\mathcal{P}$ with $\mathbb{C}\cdot x^{-1}$, and thus with $\mathbb{C}$, and for each $\theta\in S^1$, the partial order $\leqslant_\theta$ on $\mathbb{C}$ is compatible with addition and satisfies

$$c\leqslant_\theta 0\iff c=0\text{ or }\arg c-\theta\in(\pi/2,3\pi/2)\mod 2\pi.$$

We will use notation of §2.a by replacing $\varphi\in\mathcal{P}$ with $c\in\mathbb{C}$. For each pair $c\neq c'\in\mathbb{C}$, there are exactly two values of $\theta$ mod $2\pi$, say $\theta_{c,c'}$ and $\theta'_{c,c'}$, such that $c$ and $c'$ are not comparable at $\theta$. We have $\theta'_{c,c'}=\theta_{c,c'}+\pi$. These are the Stokes directions of the pair $(c,c')$. For any $\theta$ in one component of $S^1\smallsetminus\{\theta_{c,c'},\theta'_{c,c'}\}$, we have $c<_\theta c'$, and the reverse inequality for any $\theta$ in the other component.

*Stokes data.* — These are linear data which provide a description of Stokes-filtered local system. Given a finite set $C\subset\mathbb{C}$ and given $\theta_o\in S^1$ which is not a Stokes direction of any pair $c\neq c'\in C$, $\theta_o$ defines a total ordering on $C$, that we write $c_1<_{\theta_o}c_2<_{\theta_o}\cdots<_{\theta_o}c_n$.

***Definition 2.29***. — Let $C$ be a finite subset of $\mathbb{C}$ totally ordered by $\theta_o$. The category of Stokes data of type $(C,\theta_o)$ has objects consisting of two families of $\boldsymbol{k}$-vector spaces

$(G_{c,1}, G_{c,2})_{c\in C}$ and a diagram of morphisms

(2.29 $*$)
$$\begin{tikzcd}\displaystyle\bigoplus_{c\in C} G_{c,1} \arrow[r, bend left, "S"] \arrow[r, bend right, "S'"'] & \displaystyle\bigoplus_{c\in C} G_{c,2}\end{tikzcd}$$

such that, for the numbering $C = \{c_1, \dots, c_n\}$ given by $\theta_o$,

(1) $S = (S_{ij})_{i,j=1,\dots,n}$ is block-upper triangular, i.e., $S_{ij} : G_{c_i,1} \to G_{c_j,2}$ is zero unless $i \leqslant j$, and $S_{ii}$ is invertible (so $\dim G_{c_i,1} = \dim G_{c_i,2}$, and $S$ itself is invertible),

(2) $S' = (S'_{ij})_{i,j=1,\dots,n}$ is block-lower triangular, i.e., $S'_{ij} : G_{c_i,1} \to G_{c_j,2}$ is zero unless $i \geqslant j$, and $S'_{ii}$ is invertible (so $S'$ itself is invertible).

A morphism of Stokes data consists of morphisms of $\boldsymbol{k}$-vector spaces $\lambda_{c,\ell} : G_{c,\ell} \to G'_{c,\ell}$, $c \in C$, $\ell = 1, 2$ which are compatible with the diagrams (2.29 $*$).

Fixing bases in the spaces $G_{c,\ell}$, $c \in C$, $\ell = 1, 2$, allows one to present Stokes data by matrices $(\Sigma, \Sigma')$ where $\Sigma = (\Sigma_{ij})_{i,j=1,\dots,n}$ (resp. $\Sigma' = (\Sigma'_{ij})_{i,j=1,\dots,n}$) is block-lower (resp. -upper) triangular and each $\Sigma_{ii}$ (resp. $\Sigma'_{ii}$) is invertible.

The following is a translation of a classical result (see [**46**] and the references given therein, see also [**24**] for applications):

***Proposition 2.30.*** — *There is a natural functor from the category of Stokes-filtered local systems with exponential factors contained in $C$ and the category of Stokes data $(C, \theta_o)$, which is an equivalence of categories.* □

The proof of this proposition mainly uses Theorem 3.5 below, and more precisely Lemma 3.11 to define the functor.

*Duality.* — Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be a Stokes-filtered local system. Recall (see Definition 2.13) the dual local system $\mathscr{L}^\vee$ comes equipped with a Stokes filtration $(\mathscr{L}^\vee)_\bullet$ defined by

$$(\mathscr{L}^\vee)_{\leqslant c} = (\mathscr{L}_{<-c})^\perp,$$

where the orthogonality is relative to duality. In particular, $\mathrm{gr}_c(\mathscr{L}^\vee) = (\mathrm{gr}_{-c}\mathscr{L})^\vee$. Similarly, given Stokes data $((G_{c,1}, G_{c,2})_{c\in C}, S, S')$ of type $(C, \theta_0)$, let us denote by ${}^tS$ the adjoint of $S$ by duality. Define Stokes data $((G_{c,1}, G_{c,2})_{c\in C}, S, S')^\vee$ of type $(-C, \theta_o)$ by the formula $G^\vee_{-c,i} = (G_{c,i})^\vee$ $(i = 1, 2)$ and $S^\vee = {}^tS^{-1}$, $S'^\vee = {}^tS'^{-1}$, so that the diagram (2.29 $*$) becomes

(2.29 $*$)$^\vee$
$$\begin{tikzcd}\displaystyle\bigoplus_{i=1}^{r} (G_{c_i,1})^\vee \arrow[r, bend left, "{}^tS^{-1}"] \arrow[r, bend right, "{}^tS'^{-1}"'] & \displaystyle\bigoplus_{i=1}^{r} (G_{c_i,2})^\vee\end{tikzcd}$$

Then the equivalence of Proposition 2.30 is compatible with duality (see [**24**]).

# LECTURE 3

# ABELIANITY AND STRICTNESS

**Summary.** We prove that the category of $\boldsymbol{k}$-Stokes-filtered local systems on $S^1$ is abelian. The main ingredient, together with vanishing properties of the cohomology, is the introduction of the level structure. Abelianity is also a consequence of the Riemann-Hilbert correspondence considered in Lecture 5, but it is instructive to prove it over the base field $\boldsymbol{k}$.

The purpose of this lecture is to prove the following:

***Theorem 3.1***. — *The category of $\boldsymbol{k}$-Stokes-filtered local systems on $S^1$ is abelian and every morphism is strict. Moreover, it is stable by extension in the category of Stokes-prefiltered sheaves.*

By definition, Stokes-prefiltered sheaves are pre-$\mathcal{I}$-filtered sheaves on $S^1$, with $\mathcal{I}$ as in §2.c.

## 3.a. Strictness and abelianity.

— Let us recall the definition of strictness:

***Definition 3.2* (Strictness).** — A morphism $\lambda : (\mathscr{L}, \mathscr{L}_\bullet) \to (\mathscr{L}', \mathscr{L}'_\bullet)$ of pre-Stokes-filtered local systems (ramified of order $d$) is said to be *strict* if, for any $\varphi' \in \mathcal{P}_{x'}$ and any $\theta \in S^1_x$, $\lambda(\mathscr{L}_{\leqslant \varphi',\theta}) = \mathscr{L}'_{\leqslant \varphi',\theta} \cap \lambda(\mathscr{L}_\theta)$.

If a morphism $\lambda$ is strict, the two naturally defined pre-Stokes filtrations on $\lambda(\mathscr{L})$ coincide. Therefore, the local systems $\operatorname{Ker}\lambda$, $\operatorname{Im}\lambda$ and $\operatorname{Coker}\lambda$ are naturally equipped with pre-Stokes filtrations. Theorem 3.1 asserts that, if $(\mathscr{L}, \mathscr{L}_\bullet)$ and $(\mathscr{L}', \mathscr{L}'_\bullet)$ are Stokes-filtered local systems, then so are $\operatorname{Ker}\lambda$, $\operatorname{Im}\lambda$ and $\operatorname{Coker}\lambda$ (i.e., they also satisfy the local gradedness property).

***Remark 3.3***. — Let $\lambda : (\mathscr{L}, \mathscr{L}_\bullet) \to (\mathscr{L}', \mathscr{L}'_\bullet)$ be a morphism of Stokes-filtered local systems. Assume that, for every $\theta_o \in S^1$, we can find local trivializations (2.5) on some neighbourhood $\mathrm{nb}(\theta_o)$ for $(\mathscr{L}, \mathscr{L}_\bullet)$ and $(\mathscr{L}', \mathscr{L}'_\bullet)$ such that $\lambda : \mathscr{L}_{\theta_o} \to \mathscr{L}'_{\theta_o}$ is block-diagonal. Then $\lambda_{\theta_o}$ is block-diagonal in the neighbourhood of $\theta_o$ and it is strict near $\theta_o$.

***Lemma 3.4.*** — *It is enough to prove the theorem for non-ramified Stokes-filtered local systems.*

*Proof.* — Indeed, the properties considered in the theorem are local properties on $S^1$, and the compatibility property in Remark 2.25(1) implies the result. $\square$

The first part of Theorem 3.1, in the non-ramified case, is a consequence of the following more precise result, proved in §3.c:

***Theorem 3.5.*** — *Given two non-ramified Stokes-filtered local systems* $(\mathscr{L},\mathscr{L}_\bullet)$ *and* $(\mathscr{L}',\mathscr{L}'_\bullet)$*, there exist trivializations of them in the neighbourhood of any point of* $S^1$ *such that any morphism* $\lambda$ *between them is diagonal with respect to these local trivializations. In particular, such a morphism is strict, and the natural pre-Stokes filtrations on the local systems* $\operatorname{Ker}\lambda$, $\operatorname{Im}\lambda$ *and* $\operatorname{Coker}\lambda$ *are Stokes filtrations. Their sets of exponential factors satisfy*

$$\Phi(\operatorname{Ker}\lambda)\subset\Phi,\quad \Phi(\operatorname{Coker}\lambda)\subset\Phi',\quad \Phi(\operatorname{Im}\lambda)\subset\Phi\cap\Phi'.$$

**3.b. Level structure of a Stokes-filtered local system.** — In this paragraph, we work with non-ramified Stokes-filtered local systems without mentioning it explicitly. For every $\ell\in\mathbb{N}$, we define the notion of *Stokes filtration of level* $\geqslant\ell$ *on* $\mathscr{L}$, by replacing the set of indices $\mathcal{P}$ by the set $\mathcal{P}(\ell):=\mathscr{O}(*0)/x^{-\ell}\mathscr{O}$. We denote by $[\cdot]_\ell$ the map $\mathcal{P}\to\mathscr{O}(*0)/x^{-\ell}\mathscr{O}$. The constant sheaf $\mathcal{I}(\ell)$ is ordered as follows: for every connected open set $U$ of $S^1$ and $[\varphi]_\ell,[\psi]_\ell\in\mathcal{P}(\ell)$, we have $[\psi]_\ell\leqslant_U[\varphi]_\ell$ if, for some (or any) representatives $\varphi,\psi$ in $\mathscr{O}(*0)$, $e^{|x|^\ell(\psi-\varphi)}$ has *moderate growth* in a neighbourhood of $U$ in $X$ intersected with $X^*$. In particular, a Stokes structure as defined previously has level $\geqslant 0$.

Let us make this more explicit. Let $\eta\in\mathcal{P}$, $\eta=u(x)x^{-n}$ with either $\eta=0$ or $u(0)\neq 0$. Then, in a way analogous to (1.6 $*$) and (1.6 $**$), we have

$$\begin{aligned}
&[\eta]_\ell<_\theta 0 \iff u(0)\neq 0,\ n>\ell \text{ and } \arg u_n(0)-n\theta\in(\pi/2,3\pi/2)\mod 2\pi,\\
&[\eta]_\ell\leqslant_\theta 0 \iff n\leqslant\ell \text{ or } \arg u_n(0)-n\theta\in(\pi/2,3\pi/2)\mod 2\pi.
\end{aligned}$$

***Lemma 3.6.*** — *The natural morphism* $\mathcal{I}\to\mathcal{I}(\ell)$ *is compatible with the order.*

*Proof.* — Using (1.6 $*$), the previous equivalences show that $\eta\leqslant_\theta 0\Rightarrow[\eta]_\ell\leqslant_\theta 0$. $\square$

We will now introduce a reduction procedure with respect to the level. Given a Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$ (of level $\geqslant 0$), we set, using Notation 2.3,

$$\mathscr{L}_{\leqslant[\varphi]_\ell}=\sum_\psi\beta_{[\psi]_\ell\leqslant[\varphi]_\ell}\mathscr{L}_{\leqslant\psi},$$

where the sum is taken in $\mathscr{L}$. Then

$$\mathscr{L}_{<[\varphi]_\ell}:=\sum_{[\psi]_\ell}\beta_{[\psi]_\ell<[\varphi]_\ell}\mathscr{L}_{\leqslant[\psi]_\ell}=\sum_\psi\beta_{[\psi]_\ell<[\varphi]_\ell}\mathscr{L}_{\leqslant\psi}.$$

Indeed,

$$\mathscr{L}_{<[\varphi]_\ell}=\sum_{[\psi]_\ell}\sum_\eta \beta_{[\psi]_\ell<[\varphi]_\ell}\beta_{[\eta]_\ell\leqslant[\varphi]_\ell}\mathscr{L}_{\leqslant\eta},$$

and for a fixed $\eta$, the set of $\theta\in S^1$ for which there exists $[\psi]_\ell$ satisfying $[\eta]_\ell\leqslant_\theta[\psi]_\ell<_\theta[\varphi]_\ell$ is equal to the set of $\theta$ such that $[\eta]_\ell<_\theta[\varphi]_\ell$. So the right-hand term above is written

$$\sum_\eta\beta_{[\eta]_\ell<[\varphi]_\ell}\mathscr{L}_{\leqslant\eta}.$$

We can also pre-$\mathcal{I}$-filter $\mu^{-1}\operatorname{gr}_{[\varphi]_\ell}\mathscr{L}$ by setting, for $\psi\in\mathcal{P}$,

$$(\operatorname{gr}_{[\varphi]_\ell}\mathscr{L})_{\leqslant\psi}=(\mathscr{L}_{\leqslant\psi}\cap\mathscr{L}_{\leqslant[\varphi]_\ell}+\mathscr{L}_{<[\varphi]_\ell})/\mathscr{L}_{<[\varphi]_\ell}.$$

***Proposition 3.7.*** — *Assume $(\mathscr{L},\mathscr{L}_\bullet)$ is a Stokes-filtered local system (of level $\geqslant 0$) and let $\Phi$ be the finite set of its exponential factors.*

(1) *For each $\ell\in\mathbb{N}$, $\mathscr{L}_{\leqslant[\bullet]_\ell}$ defines a Stokes-filtered local system $(\mathscr{L},\mathscr{L}_{[\bullet]_\ell})$ of level $\geqslant\ell$ on $\mathscr{L}$, $\operatorname{gr}_{[\varphi]_\ell}\mathscr{L}$ is locally isomorphic to $\bigoplus_{\psi,\,[\psi]_\ell=[\varphi]_\ell}\operatorname{gr}_\psi\mathscr{L}$, and the set of exponential factors of $(\mathscr{L},\mathscr{L}_{[\bullet]_\ell})$ is $\Phi(\ell):=\operatorname{image}(\Phi\to\mathcal{P}(\ell))$.*

(2) *For every $[\varphi]_\ell\in\Phi(\ell)$, $(\operatorname{gr}_{[\varphi]_\ell}\mathscr{L},(\operatorname{gr}_{[\varphi]_\ell}\mathscr{L})_\bullet)$ is a Stokes-filtered local system and its set of exponential factors is the pull-back of $[\varphi]_\ell$ by $\Phi\to\Phi(\ell)$.*

(3) *Let us set*

$$\bigl(\operatorname{gr}_\ell\mathscr{L},(\operatorname{gr}_\ell\mathscr{L})_\bullet\bigr):=\bigoplus_{[\psi]_\ell\in\Phi(\ell)}\bigl(\operatorname{gr}_{[\psi]_\ell}\mathscr{L},(\operatorname{gr}_{[\psi]_\ell}\mathscr{L})_\bullet\bigr).$$

*Then $(\operatorname{gr}_\ell\mathscr{L},(\operatorname{gr}_\ell\mathscr{L})_\bullet)$ is a Stokes-filtered local system (of level $\geqslant 0$) which is locally isomorphic to $(\mathscr{L},\mathscr{L}_\bullet)$.*

*Proof.* — All the properties are local, so we can assume that $(\mathscr{L},\mathscr{L}_\bullet)$ is graded. We have then

$$\mathscr{L}_{\leqslant\psi}=\bigoplus_\eta\beta_{\eta\leqslant\psi}\operatorname{gr}_\eta\mathscr{L},$$

hence

$$\mathscr{L}_{\leqslant[\varphi]_\ell}=\bigoplus_\eta\Bigl(\sum_\psi\beta_{[\psi]_\ell\leqslant[\varphi]_\ell}\beta_{\eta\leqslant\psi}\operatorname{gr}_\eta\mathscr{L}\Bigr),\quad \mathscr{L}_{<[\varphi]_\ell}=\bigoplus_\eta\Bigl(\sum_\psi\beta_{[\psi]_\ell<[\varphi]_\ell}\beta_{\eta\leqslant\psi}\operatorname{gr}_\eta\mathscr{L}\Bigr)$$

For a fixed $\eta$, the set of $\theta\in S^1$ such that there exists $\psi$ with $\eta\leqslant_\theta\psi$ and $[\psi]_\ell\leqslant_\theta[\varphi]_\ell$ (resp. $[\psi]_\ell<_\theta[\varphi]_\ell$) is the set of $\theta$ such that $[\eta]_\ell\leqslant_\theta[\varphi]_\ell$ (resp. $[\eta]_\ell<_\theta[\varphi]_\ell$), so

$$\mathscr{L}_{\leqslant[\varphi]_\ell}=\bigoplus_\eta\beta_{[\eta]_\ell\leqslant[\varphi]_\ell}\operatorname{gr}_\eta\mathscr{L},\quad \mathscr{L}_{<[\varphi]_\ell}=\bigoplus_\eta\beta_{[\eta]_\ell<[\varphi]_\ell}\operatorname{gr}_\eta\mathscr{L},$$

which gives 3.7(1).

Let us show 3.7(2). All sheaves entering in the definition of $(\operatorname{gr}_{[\varphi]_\ell}\mathscr{L})_{\leqslant\psi}$, $(\operatorname{gr}_{[\varphi]_\ell}\mathscr{L})_{<\psi}$ and $\operatorname{gr}_\psi\operatorname{gr}_{[\varphi]_\ell}\mathscr{L}$ decompose with respect to $\eta$ as above. Given $\eta\in\Phi$ and $\theta\in S^1$, the component $\operatorname{gr}_\eta\mathscr{L}_\theta$ occurs in $\mathscr{L}_{\leqslant\psi,\theta}\cap\mathscr{L}_{\leqslant[\varphi]_\ell,\theta}$ if and only if $\eta\leqslant_\theta\psi$

and $[\eta]_\ell \leqslant_\theta [\varphi]_\ell$. Arguing similarly, we finally find that $\operatorname{gr}_\eta \mathscr{L}_\theta$ occurs in $\operatorname{gr}_\psi \operatorname{gr}_{[\varphi]_\ell} \mathscr{L}_\theta$ if and only if $\eta = \psi$ and $[\eta]_\ell = [\varphi]_\ell$, that is,

$$\operatorname{gr}_\psi \operatorname{gr}_{[\varphi]_\ell} \mathscr{L}_\theta = \begin{cases} 0 & \text{if } [\psi]_\ell \neq [\varphi]_\ell, \\ \operatorname{gr}_\psi \mathscr{L}_\theta & \text{if } [\psi]_\ell = [\varphi]_\ell. \end{cases}$$

This concludes the proof of 3.7(2). Now, 3.7(3) is clear. ☐

***Remark 3.8***. — Let us explain the meaning and usefulness of this proposition. To a Stokes-filtered local system $(\mathscr{L}, \mathscr{L}_\bullet)$ is associated a partially graded Stokes-filtered local system $(\operatorname{gr}_\ell \mathscr{L}, (\operatorname{gr}_\ell \mathscr{L})_\bullet)$. Going from $(\mathscr{L}, \mathscr{L}_\bullet)$ to $(\operatorname{gr}_\ell \mathscr{L}, (\operatorname{gr}_\ell \mathscr{L})_\bullet)$ consists in

- considering $(\mathscr{L}, \mathscr{L}_\bullet)$ as indexed by $\mathcal{P}(\ell)$ (or $\Phi(\ell)$) and grading it as such,
- remembering the $\mathcal{P}$-filtration on the graded object, making it a Stokes-filtered local system as well.

Conversely, let $(\mathscr{G}_\ell, \mathscr{G}_{\ell,\bullet})$ be a fixed Stokes-filtered local system graded at the level $\ell \geqslant 0$, that is, the associated $\mathcal{P}(\ell)$-filtered local system is graded. As a consequence of the last statement of the proposition, an argument similar to that of Proposition 1.43 implies that the pointed set of isomorphism classes of Stokes-filtered local systems $(\mathscr{L}, \mathscr{L}_\bullet)$ equipped with an isomorphism $f_\ell : (\operatorname{gr}_\ell \mathscr{L}, (\operatorname{gr}_\ell \mathscr{L})_\bullet) \xrightarrow{\sim} (\mathscr{G}_\ell, \mathscr{G}_{\ell,\bullet})$ is in bijection with the pointed set $H^1\big(S^1, \mathscr{A}ut^{<_{[\bullet]_\ell}0}(\mathscr{G}_\ell)\big)$, where $\mathscr{A}ut^{<_{[\bullet]_\ell}0}(\mathscr{G}_\ell)$ is the sheaf of automorphisms $\lambda$ of $(\mathscr{G}_\ell, \mathscr{G}_{\ell,\bullet})$ such that $\operatorname{gr}_{[\varphi]_\ell} \lambda = \mathrm{Id}$ on the local system $\operatorname{gr}_{[\varphi]_\ell} \mathscr{G}_\ell$ for any $\varphi \in \mathcal{P}$ (equivalently, any $[\varphi]_\ell$ in $\mathcal{P}(\ell)$).

In particular, one can reconstruct $(\mathscr{L}, \mathscr{L}_\bullet)$ from $\operatorname{gr} \mathscr{L}$ either in one step, by specifying an element of $H^1(S^1, \mathscr{A}ut^{<0} \operatorname{gr} \mathscr{L})$, or step by step with respect to the level, by specifying at each step $\ell$ an element of $H^1\big(S^1, \mathscr{A}ut^{<_{[\bullet]_\ell}0}(\operatorname{gr}_\ell \mathscr{L})\big)$.

## 3.c. Proof of Theorem 3.5.

— It will be done by induction, using Corollary 3.10 below for the inductive assumption on $\operatorname{gr}_\ell \mathscr{L}$ of Proposition 3.7(3).

Let $\Phi$ be a finite set in $\mathcal{P}$. Assume $\#\Phi \geqslant 2$. We then set $m(\Phi) = \max\{m(\varphi - \psi) \mid \varphi \neq \psi \in \Phi\}$, and we have $m(\Phi) > 0$. If $\#\Phi = 1$, we set $m(\Phi) = 0$. We also set $\ell = m(\Phi) - 1$ if $m(\Phi) > 0$.

***Lemma 3.9***. — *Assume $\#\Phi \geqslant 2$ and fix $\varphi_o \in \Phi$. Then*

$$m(\Phi, \varphi_o) := \max\{m(\varphi - \varphi_o) \mid \varphi \in \Phi\} = m(\Phi).$$

*Proof*. — Let $\eta$ be the sum of common monomials to all $\varphi \in \Phi$. By replacing $\Phi$ with $\Phi - \eta$ we can assume that $\eta = 0$. We will show that, in such a case, $m(\Phi) = \max\{m(\varphi) \mid \varphi \in \Phi\} = \max\{m(\varphi - \varphi_o) \mid \varphi \in \Phi\}$. If all $\varphi \in \Phi$ have the same order $k > 0$, two of them, say $\varphi_1$ and $\varphi_2$, do not have the same dominant monomial. Then $m(\Phi) = k$. A given $\varphi_o \in \Phi$ cannot have the same dominant monomial as $\varphi_1$ and $\varphi_2$, so one of $\varphi_1 - \varphi_o$ and $\varphi_2 - \varphi_o$ has order $k$, and $m(\Phi, \varphi_o) = k$.

Assume now that $\varphi_1$ has maximal order $k$ and $\varphi_2$ has order $< k$. Then $m(\Phi) = k$. Given $\varphi_o \in \Phi$, then either $\varphi_o$ has order $k$, and then $\varphi_2 - \varphi_o$ has order $k$, or $\varphi_o$ has order $< k$, and then $\varphi_1 - \varphi_o$ has order $k$, so in any case we have $m(\Phi, \varphi_o) = k$. $\square$

Let us fix $\varphi_o \in \Phi$. In the following, we will work with $\Phi - \varphi_o$. We will assume that $\#\Phi \geqslant 2$, that is, $m(\Phi) > 0$ (otherwise the theorem is clear).

Let $\varphi \in \Phi - \varphi_o$. If $m(\varphi) = m(\Phi)$, then its image in $(\Phi - \varphi_o)(\ell)$ is nonzero. For every $\varphi \in \Phi - \varphi_o$, the subset $(\Phi - \varphi_o)_{[\varphi]_\ell} := \{\psi \in \Phi - \varphi_o \mid [\psi]_\ell = [\varphi]_\ell\}$ satisfies $m((\Phi - \varphi_o)_{[\varphi]_\ell}) \leqslant \ell < m$. Indeed, if $[\varphi]_\ell = 0$, then any $\psi \in (\Phi - \varphi_o)_{[\varphi]_\ell}$ can be written as $x^{-\ell} u_\psi(x)$ with $u_\psi(x) \in \mathscr{O}$, and the difference of two such elements is written $x^{-\ell} v(x)$ with $v(x) \in \mathscr{O}$. On the other hand, if $[\varphi]_\ell \neq 0$, $\psi$ is written as $\varphi + x^{-\ell} u_\psi(x)$ and the same argument applies.

***Corollary 3.10* (of Prop. 3.7).** — *Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be a Stokes-filtered local system, let $\Phi''$ be a finite subset of $\mathcal{P}$ containing $\Phi(\mathscr{L}, \mathscr{L}_\bullet)$, set $m = m(\Phi'')$, $\ell = m(\Phi'') - 1$, and fix $\varphi_o$ as above.*

*Then, for every $[\varphi]_\ell \in (\Phi'' - \varphi_o)(\ell)$, $(\mathrm{gr}_{[\varphi]_\ell} \mathscr{L}[-\varphi_o], (\mathrm{gr}_{[\varphi]_\ell} \mathscr{L}[-\varphi_o])_\bullet)$ is a Stokes-filtered local system and $m_{\max}(\mathrm{gr}_{[\varphi]_\ell} \mathscr{L}[-\varphi_o], (\mathrm{gr}_{[\varphi]_\ell} \mathscr{L}[-\varphi_o])_\bullet) \leqslant \ell < m$.* $\square$

***Lemma 3.11*.** — *Let $I$ be any open interval of $S^1$ such that, for any $\varphi, \psi \in \Phi$, $\mathrm{card}(I \cap \mathrm{St}(\varphi, \psi)) \leqslant 1$. Then the decompositions* (2.5) *hold on $I$.*

*Proof.* — It is enough to show that $H^1(I, \mathscr{L}_{<\varphi}) = 0$ for any $\varphi \in \Phi$. Indeed, arguing as in the proof of Lemma 2.7, if we restrict to such a $I$, we can lift for any $\varphi$ a basis of global sections of $\mathrm{gr}_\varphi \mathscr{L}$ as sections of $\mathscr{L}_{\leqslant\varphi}$ and get an injective morphism $\mathrm{gr}_\varphi \mathscr{L} \to \mathscr{L}_{\leqslant\varphi}$. We therefore obtain a morphism $\bigoplus_{\varphi\in\Phi} \mathrm{gr}_\varphi \mathscr{L} \to \mathscr{L}$ sending $\mathrm{gr}_\varphi \mathscr{L}$ into $\mathscr{L}_{\leqslant\varphi}$ for each $\varphi \in \Phi$. This is an isomorphism: indeed, as both sheaves are local systems, it is enough to check this at some $\theta \in I$; considering a splitting of $\mathscr{L}$ at $\theta$, the matrix of this morphism is block-triangular with respect to the order at $\theta$ and the diagonal blocks are equal to the identity. It remains to show that this morphism induces the splitting for each $\mathscr{L}_\eta$. For every $\eta$, we have a natural inclusion $\beta_{\varphi\leqslant\eta} \mathscr{L}_{\leqslant\varphi} \hookrightarrow \mathscr{L}_{\leqslant\eta}$, hence the previous isomorphism induces a morphism $\bigoplus_{\varphi\in\Phi} \beta_{\varphi\leqslant\eta} \mathrm{gr}_\varphi \mathscr{L} \to \mathscr{L}_{\leqslant\eta}$, which is seen to be an isomorphism on stalks.

Let us now show that $H^1(I, \mathscr{L}_{<\varphi}) = 0$. It will be easier to argue by duality. According to Lemma 2.17, the sheaf $(\mathscr{L}_{<\varphi})^\vee$ coincides with the Poincaré-Verdier dual of $\mathscr{L}_{<\varphi}$ and has a local decomposition

$$(\mathscr{L}_{<\varphi})^\vee_{|\mathrm{nb}(\theta_o)} \simeq \bigoplus_{\psi\in\Phi} \gamma_{\psi<\varphi} (\mathrm{gr}_\psi \mathscr{L})^\vee_{|\mathrm{nb}(\theta_o)}, \tag{3.12}$$

where $\gamma_{\psi<\varphi}$ denotes the functor of restriction to $S_{\psi<\varphi}$ composed with the usual direct image by the inclusion $S_{\psi<\varphi} \hookrightarrow S^1$, and it is enough to show that $H^0_c(I, (\mathscr{L}_{<\varphi})^\vee) = 0$. Let us set $I = (\theta_0, \theta_{n+1})$ and let us denote by $\theta_1, \dots, \theta_n$ the successive Stokes directions in $I$. We set $I_i = (\theta_{i-1}, \theta_{i+1})$ for $i = 1, \dots, n$. The decomposition (3.12) holds on each $I_i$ and, on $I_i \cap I_{i+1}$, two decompositions (3.12) are related by $\lambda^{(i,i+1)}$ whose component $\lambda^{(i,i+1)}_{\psi,\eta}:\gamma_{\psi<\varphi}(\operatorname{gr}_\psi\mathscr{L})^\vee_{|I_i\cap I_{i+1}}\to\gamma_{\eta<\varphi}(\operatorname{gr}_\eta\mathscr{L})^\vee_{|I_i\cap I_{i+1}}$ is nonzero only if $\psi\leqslant_{I_i\cap I_{i+1}}\eta$, and is equal to Id if $\psi=\eta$.

Assume that $s\in H^0_c(I,(\mathscr{L}_{<\varphi})^\vee)$ is nonzero. For any $i=1,\dots,n$, we denote by $s_i$ its restriction to $I_i$ and $s_{i,\psi}$ its component on $\gamma_{\psi<\varphi}(\operatorname{gr}_\psi\mathscr{L})^\vee_{|I_i}$ for the chosen decomposition on $I_i$. Therefore, $s_{i,\psi}$ and $s_{i+1,\psi}$ may differ on $I_i\cap I_{i+1}$. Let us note, however, that the set $S$ is totally ordered by $\leqslant_{I_i\cap I_{i+1}}$ on $I_i\cap I_{i+1}$, and if we set $\psi_i=\min\{\psi\in S\mid s_{i,\psi}\neq 0 \text{ on } I_i\cap I_{i+1}\}$ (when defined, i.e., if $s_i\not\equiv 0$ on $I_i\cap I_{i+1}$), then we also have $\psi_i=\min\{\psi\in S\mid s_{i+1,\psi}\neq 0\text{ on } I_i\cap I_{i+1}\}$. Indeed, let us denote by $\psi_i'$ the right-and term.

- On the one hand, $s_{i+1,\psi_i}=\sum_{\psi\leqslant\psi_i}\lambda^{(i,i+1)}_{\psi,\psi_i}(s_{i,\psi})=s_{i,\psi_i}$ on $I_i\cap I_{i+1}$ since $\lambda^{(i,i+1)}_{\psi,\psi_i}=0$ if $\psi<\psi_i$ on $I_i\cap I_{i+1}$, and $\lambda^{(i,i+1)}_{\psi_i,\psi_i}=\mathrm{Id}$. Therefore, $\psi_i'\leqslant\psi_i$ on $I_i\cap I_{i+1}$.
- On the other hand, if $\psi_i'<\psi_i$ on $I_i\cap I_{i+1}$, then $s_{i+1,\psi_i'}=\sum_{\psi\leqslant\psi_i'}\lambda^{(i,i+1)}_{\psi,\psi_i'}(s_{i,\psi})=0$, a contradiction.

Since $s$ is compactly supported on $I$, its restriction to $(\theta_0,\theta_1)$ and to $(\theta_n,\theta_{n+1})$ vanishes identically. In the following, we assume that $s_1$ and $s_n$ are not identically zero (otherwise, we just forget these Stokes directions and consider the first and last one for which $s_i$ is not identically zero). Then $\psi_1$ is defined, and we have

- $\psi_1<\varphi$ on $I_1\cap I_2$ since $s_{1,\psi_1}\neq 0$ on $I_1\cap I_2$,
- $\theta_1\in\operatorname{St}(\psi_1,\varphi)$ since $s_{1,\psi_1}=0$ on $(\theta_0,\theta_1)$,
- $\varphi<\psi_1$ on $(\theta_0,\theta_1)$ since $\theta_1\in\operatorname{St}(\psi_1,\varphi)$,
- and $\psi_1<\varphi$ on $(\theta_1,\theta_{n+1})$, as it is so on $(\theta_1,\theta_2)$ and there are no other element of $\operatorname{St}(\psi_1,\varphi)$ in $(\theta_1,\theta_{n+1})$.

It follows that $s_{2,\psi_1}\not\equiv 0$ on $I_1\cap I_2$. Moreover, $\theta_2\notin\operatorname{St}(\psi_1,\varphi)$, so $s_{2,\psi_1}\neq 0$ on $I_2$ (being a section of a local system on this interval). Therefore, $s_2\neq 0$ on $I_2\cap I_3$, $\psi_2$ is defined and (by definition) $\psi_2\leqslant\psi_1$ on $I_2\cap I_3$, and there is a component $s_{2,\psi_2}$ on $I_2$ which is not identically zero. In conclusion, on $I_2\cap I_3$, the following holds:

- $\psi_2\leqslant\psi_1<\varphi$,
- $s_{2,\psi_2}\neq 0$.

We claim that $\psi_2<\varphi$ on $(\theta_2,\theta_{n+1})$ (and in particular $\theta_3\notin\operatorname{St}(\psi_2,\varphi)$):

- this is already proved if $\psi_2=\psi_1$;
- assume therefore that $\psi_2<\psi_1$ on $I_2\cap I_3$; then,
  - if $s_{2,\psi_2}\neq 0$ on $I_1\cap I_2$, then $\psi_1<\psi_2$ on $I_1\cap I_2$ (by definition of $\psi_1$) and so $\theta_2\in\operatorname{St}(\psi_1,\psi_2)$, hence there is no other element of $\operatorname{St}(\psi_1,\psi_2)$ belonging to $(\theta_2,\theta_{n+1})$ and therefore also $\psi_2<\varphi$ since $\psi_1<\varphi$ on $(\theta_1,\theta_{n+1})$;
  - if $s_{2,\psi_2}=0$ on $I_1\cap I_2$, this means that $\theta_2\in\operatorname{St}(\psi_2,\varphi)$, hence there is no other element of $\operatorname{St}(\psi_2,\varphi)$ belonging to $(\theta_2,\theta_{n+1})$, so we keep $\psi_2<\varphi$ on $(\theta_2,\theta_{n+1})$.

Continuing in the same way, we find that $\psi_n$ is defined, which means that $s_n$ is not identically zero on $(\theta_n,\theta_{n+1})$, a contradiction. □

***Remark 3.13.*** — For $I \neq S^1$ as in Lemma 3.11, we also have $H^1(I, \mathscr{L}_{\leqslant\varphi}) = 0$ for any $\varphi$. This follows from the exact sequence

$$0 = H^1(I, \mathscr{L}_{<\varphi}) \longrightarrow H^1(I, \mathscr{L}_{\leqslant\varphi}) \longrightarrow H^1(I, \operatorname{gr}_\varphi \mathscr{L}) = 0,$$

where the last equality holds because $\operatorname{gr}_\varphi \mathscr{L}$ is a local system.

***Corollary 3.14.*** — *In the setting of Corollary* 3.10*, let* $I$ *be any open interval of* $S^1$ *such that, for any* $\varphi, \psi \in \Phi$*,* $\operatorname{card}(I \cap \operatorname{St}(\varphi, \psi)) \leqslant 1$*. Then*

$$(\mathscr{L}, \mathscr{L}_\bullet)_{|I} \simeq (\operatorname{gr}_\ell \mathscr{L}, (\operatorname{gr}_\ell \mathscr{L})_\bullet)_{|I}.$$

*Proof.* — By Proposition 3.7(3), the set of exponential factors of $(\mathscr{L}, \mathscr{L}_\bullet)$ and $(\operatorname{gr}_\ell \mathscr{L}, (\operatorname{gr}_\ell \mathscr{L})_\bullet)$ coincide, hence $I$ satisfies the assumption of Lemma 3.11 for both $(\mathscr{L}, \mathscr{L}_\bullet)$ and $(\operatorname{gr}_\ell \mathscr{L}, (\operatorname{gr}_\ell \mathscr{L})_\bullet)$. Therefore, when restricted to $I$, both are isomorphic to the graded Stokes-filtered local system determined by $\operatorname{gr} \mathscr{L}$ restricted to $I$. ☐

*End of the proof of Theorem* 3.5. — Let $\lambda : \mathscr{L} \to \mathscr{L}'$ be a morphism of Stokes-filtered local systems. The proof that $\lambda$ is strict and that $\operatorname{Ker} \lambda$, $\operatorname{Im} \lambda$ and $\operatorname{Coker} \lambda$ (equipped with the naturally induced pre-Stokes structures) are Stokes-filtered local systems follows from the local decomposition of the morphism, the proof of which will be done by induction on $m = m(\Phi'')$, with $\Phi'' = \Phi \cup \Phi'$. The result is clear if $m = 0$ (so $\Phi'' = \{0\}$), as both Stokes-filtered local systems have only one jump.

Assume now $m \neq 0$. We fix $\varphi_o \in \Phi''$ as in Corollary 3.10. It is enough to prove the assertion for $\lambda : \mathscr{L}[-\varphi_o] \to \mathscr{L}'[-\varphi_o]$. We shall assume that $\varphi_o = 0$ in order to simplify the notation below. Let $I$ be any open interval of $S^1$ of length $\pi/m$ with no Stokes points (of $\Phi''$) as end points. According to the choice of $m$ and to Corollary 3.14, $\lambda_{|I}$ decomposes as $\bigoplus_{[\psi]_\ell, [\psi']_\ell} \lambda_{|I, [\psi]_\ell, [\psi']_\ell}$, with $\lambda_{|I, [\psi]_\ell, [\psi']_\ell} : \operatorname{gr}_{[\psi]_\ell} \mathscr{L}_{|I} \to \operatorname{gr}_{[\psi']_\ell} \mathscr{L}'_{|I}$, where $[\psi]_\ell$ runs in $\Phi(\ell)$ and $[\psi']_\ell$ in $\Phi'(\ell)$ (otherwise, it is zero). Each $\lambda_{|I, [\psi]_\ell, [\psi']_\ell}$ is a morphism of (constant) local systems, hence is constant.

Let us show that $\lambda_{|I, [\psi]_\ell, [\psi']_\ell} = 0$ if $[\psi]_\ell \neq [\psi']_\ell$. We denote by $\mathcal{P}_{[\psi]_\ell}$ the set of $\eta \in \mathcal{P}$ such that $[\eta]_\ell = [\psi]_\ell$. According to the proof of Corollary 3.14, $\lambda_{|I, [\psi]_\ell, [\psi']_\ell}$ itself decomposes as blocks with components $\lambda_{|I, \eta, \eta'}$ where $\eta$ varies in $\mathcal{P}_{[\psi]_\ell}$ and $\eta'$ in $\mathcal{P}_{[\psi']_\ell}$. We will show that each such block is zero.

As in restriction to $I$, $\lambda$ is (isomorphic to) a filtered morphism $\operatorname{gr}_\ell \mathscr{L} \to \operatorname{gr}_\ell \mathscr{L}'$, it sends $(\operatorname{gr}_\ell \mathscr{L})_{\leqslant\varphi}$ to $(\operatorname{gr}_\ell \mathscr{L}')_{\leqslant\varphi}$ for any $\varphi$, that is, after the proof of Corollary 3.14, it sends $\bigoplus_{[\psi]_\ell} \bigoplus_{\eta \in \mathcal{P}_{[\psi]_\ell}} \beta_{\eta \leqslant \varphi} \operatorname{gr}_\eta \mathscr{L}$ in $\bigoplus_{[\psi']_\ell} \bigoplus_{\eta' \in \mathcal{P}_{[\psi']_\ell}} \beta_{\eta' \leqslant \varphi} \operatorname{gr}_{\eta'} \mathscr{L}$, so the block $\lambda_{|I, \eta, \eta'}$ sends $\beta_{\eta \leqslant \varphi} \operatorname{gr}_\eta \mathscr{L}$ in $\beta_{\eta' \leqslant \varphi} \operatorname{gr}_{\eta'} \mathscr{L}$. Choosing $\varphi = \eta$ shows that $\lambda_{|I, \eta, \eta'}$ sends $\operatorname{gr}_\eta \mathscr{L}$ in $\beta_{\eta' \leqslant \eta} \operatorname{gr}_{\eta'} \mathscr{L}$. Now, if $[\psi]_\ell \neq [\psi']_\ell$, we have $\eta - \eta' = x^{-m} u(x)$ with $u(0) \neq 0$ and, $I$ being of length $\pi/m$, contains at least one Stokes point of $\operatorname{St}(\eta, \eta')$. Therefore $\lambda_{|I, \eta, \eta'}$ vanishes on some non-empty open subset of $I$, hence everywhere.

In conclusion, $\lambda_{|I}$ is diagonal with respect to the $[\bullet]_\ell$-decomposition, that is, it coincides with the graded morphism $\operatorname{gr}_{[\bullet]_\ell} \lambda$, graded with respect to the filtration $\mathscr{L}_{[\bullet]}$ of level $\geqslant \ell$.

By Proposition 3.7(2) and Corollary 3.10, the inductive assumption can be applied to $\mathrm{gr}_{[\bullet]_\ell}\lambda$. By induction we conclude that $\lambda$ can be diagonalized (with respect to the decomposition in the third line of (2.5)) in the neighbourhood[(1)] of any point of $I$. As this holds for any such $I$, this ends the proof of the theorem in the non-ramified case. □

***Corollary 3.15* (of Lemma 3.11).** — *Let* $(\mathscr{L},\mathscr{L}_\bullet)$ *be a Stokes-filtered local system. Then, for any* $\varphi\in\mathcal{P}_x$, $\Gamma(S^1,\mathscr{L}_{<\varphi})=0$.

*Proof.* — Let us keep the notation used above, and introduced at the beginning of §3.c. It is easy to reduce to the case where $(\mathscr{L},\mathscr{L}_\bullet)$ is non-ramified, that we assume to hold below.

We will argue by decreasing induction on $m(\Phi)$. If $m(\Phi)=0$, so $\Phi=\{\varphi_o\}$, we have $\mathscr{L}_{<\varphi}=\beta_{\varphi_o<\varphi}\mathscr{L}$ (see Example 2.11(1)), and the assertion is clear.

Assume now $m(\Phi)>0$, that is, $\#\Phi\geqslant 2$. By twisting it is enough to show $\Gamma(S^1,\mathscr{L}_{<0})=0$.

There is a natural inclusion $\mathscr{L}_{\leqslant 0}\hookrightarrow\mathscr{L}_{\leqslant[0]_\ell}$, hence a map $\mathscr{L}_{\leqslant 0}\to\mathrm{gr}_{[0]_\ell}\mathscr{L}$ sending $\mathscr{L}_{\leqslant 0}$ to $(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{\leqslant 0}$ and $\mathscr{L}_{<0}$ to $(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{<0}$. It is enough to show that the induced map $\Gamma(S^1,\mathscr{L}_{<0})\to\Gamma(S^1,(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{<0})$ is injective. Indeed, we can apply the inductive assumption to the Stokes-filtered local system $(\mathrm{gr}_{[0]_\ell}\mathscr{L},(\mathrm{gr}_{[0]_\ell}\mathscr{L})_\bullet)$ to conclude that $\Gamma(S^1,(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{<0})=0$.

Let $I$ be an interval of length $\pi/m(\Phi)$ with no Stokes directions for $\Phi\cup\{0\}$ at its boundary. We will show that $\Gamma(I,\mathscr{L}_{<0})\to\Gamma(I,(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{<0})$ is injective for any such $I$, which is enough. By Lemma 3.11, we have $\mathscr{L}_{<0|I}\simeq\bigoplus_{\varphi\in\Phi}\beta_{\varphi<0}\,\mathrm{gr}_\varphi\mathscr{L}_{|I}$, and the map $\mathscr{L}_{<0|I}\to(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{<0|I}$ is the projection to the sum taken on $\varphi\in[0]_\ell$, that is, the $\varphi\in\Phi$ having a pole of order $<m(\Phi)$. But since $I$ contains a Stokes direction for $(\varphi,0)$ for any $\varphi\notin[0]_\ell$, we have $\Gamma(I,\beta_{\varphi<0}\,\mathrm{gr}_\varphi\mathscr{L}_{|I})=0$ for any such $\varphi$. We conclude that $\Gamma(I,\mathscr{L}_{<0|I})\to\!\!\to\Gamma(I,(\mathrm{gr}_{[0]_\ell}\mathscr{L})_{<0})$ is isomorphic to the identity map from $\Gamma\big(I,\bigoplus_{\varphi\in[0]_\ell}\beta_{\varphi<0}\,\mathrm{gr}_\varphi\mathscr{L}_{|I}\big)$ to itself, by applying the same reasoning to $(\mathrm{gr}_{[0]_\ell}\mathscr{L},(\mathrm{gr}_{[0]_\ell}\mathscr{L})_\bullet)$. □

***Remark 3.16.*** — The dimension of the only non-zero cohomology space $H^1(S^1,\mathscr{L}_{<0})$ is usually called the *irregularity number* of $(\mathscr{L},\mathscr{L}_\bullet)$.

***Corollary 3.17.*** — *Let* $f:(\mathscr{L},\mathscr{L}_\bullet)\to(\mathscr{L}',\mathscr{L}'_\bullet)$ *be a morphism of Stokes-filtered local systems on* $S^1$ *sending* $\mathscr{L}_\leqslant$ *to* $\mathscr{L}'_<$. *Then* $f$ *is equal to zero.*

*Proof.* — Indeed, $f$ is a section on $S^1$ of $\mathscr{H}\!om\big((\mathscr{L},\mathscr{L}_\bullet),(\mathscr{L}',\mathscr{L}'_\bullet)\big)_{<0}$. □

---

(1) In fact, the argument shows that the diagonalization holds on $I$.

**3.d. Proof of the stability by extension.** — We continue to work in the non-ramified case. We consider an exact sequence

$$0 \longrightarrow (\mathscr{L}', \mathscr{L}'_\bullet) \longrightarrow (\mathscr{L}, \mathscr{L}_\bullet) \longrightarrow (\mathscr{L}'', \mathscr{L}''_\bullet) \longrightarrow 0,$$

where $(\mathscr{L}', \mathscr{L}'_\bullet), (\mathscr{L}'', \mathscr{L}''_\bullet)$ are Stokes-filtered local systems, but no assumption is made on $(\mathscr{L}, \mathscr{L}_\bullet)$, except that $(\mathscr{L}, \mathscr{L}_\bullet)$ is a pre-$\mathcal{I}$-filtered sheaf ($\mathcal{I} = \mathcal{I}_1$ here) in the sense of Definition 1.8. Firstly, the extension $\mathscr{L}$ of $\mathscr{L}'$ and $\mathscr{L}''$ is clearly a local system.

Let us now restrict to an interval $I$ as in Lemma 3.11, with $\Phi = \Phi' \cup \Phi''$. Since $H^1(I, \mathscr{L}'_{\leqslant\varphi}) = 0$ for any $\varphi \in \mathcal{P}$ (see Remark 3.13), there is a section of $\mathscr{L}_{\leqslant\varphi} \to \mathscr{L}''_{\leqslant\varphi}$, and therefore an injective morphism $\operatorname{gr}_\varphi \mathscr{L}'' \hookrightarrow \mathscr{L}_{\leqslant\varphi}$ for any $\varphi \in \Phi''$. By the filtration property, it defines for any $\psi \in \mathcal{P}$ an injective morphism $\beta_{\varphi\leqslant\psi} \operatorname{gr}_\varphi \mathscr{L}'' \hookrightarrow \mathscr{L}_{\leqslant\psi}$ and then a section $\mathscr{L}''_{\leqslant\psi} = \bigoplus_{\varphi\in\Phi''} \beta_{\varphi\leqslant\psi} \operatorname{gr}_\varphi \mathscr{L}'' \to \mathscr{L}_{\leqslant\psi}$ of the projection $\mathscr{L}_{\leqslant\psi} \to \mathscr{L}''_{\leqslant\psi}$, giving rise to an isomorphism $(\mathscr{L}, \mathscr{L}_\bullet)_{|I} \simeq (\mathscr{L}', \mathscr{L}'_\bullet)_{|I} \oplus (\mathscr{L}'', \mathscr{L}''_\bullet)_{|I}$. In particular, $(\mathscr{L}, \mathscr{L}_\bullet)$ is a Stokes filtration when restricted to $I$.

As this holds for any such $I$, this proves the assertion. $\square$

**3.e. More on the level structure.** — It will be useful to understand the level structure in an intrinsic way when the Stokes-filtered local system is possibly ramified. We will therefore use the presentation and the notation of Remark 2.23.

Let $\ell$ be any non-negative rational number. We define the subsheaf $(\widetilde{\jmath}_*\mathscr{O}_{X^*})(\ell) \subset \widetilde{\jmath}_*\mathscr{O}_{X^*}$ as the subsheaf of germs $\varphi$ such that $x^\ell\varphi$ is a local section of $(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$, for some (or any) local determination of $x^\ell$. We have, for any $\ell \leqslant \ell'$,

$$(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}} = (\widetilde{\jmath}_*\mathscr{O}_{X^*})(0) \subset (\widetilde{\jmath}_*\mathscr{O}_{X^*})(\ell) \subset (\widetilde{\jmath}_*\mathscr{O}_{X^*})(\ell') \subset \widetilde{\jmath}_*\mathscr{O}_{X^*}.$$

We set $\mathcal{I}(\ell) = \widetilde{\mathcal{I}}/\widetilde{\mathcal{I}} \cap (\widetilde{\jmath}_*\mathscr{O}_{X^*})(\ell)$. We have a natural morphism $\mathcal{I} = \mathcal{I}(0) \to \mathcal{I}(\ell)$ which is compatible with the order (same proof as for Lemma 3.6). We have a commutative diagram

$$\begin{tikzcd}
\mathcal{I}^{\text{ét}} \arrow[r, "q_\ell"] \arrow[dr, "\mu"'] & \mathcal{I}(\ell)^{\text{ét}} \arrow[d, "\mu_\ell"] \\
 & S^1
\end{tikzcd}$$

The map $q_\ell$ is étale and onto. If $\Sigma \subset \mathcal{I}^{\text{ét}}$ is a finite covering of $S^1$ (via $\mu$), then $\Sigma(\ell) := q_\ell(\Sigma)$ is a finite covering of $S^1$ (via $\mu_\ell$), and $q_\ell : \Sigma \to \Sigma(\ell)$ is also a finite covering. Moreover, since $q_\ell : \mathcal{I}^{\text{ét}}_{|\Sigma(\ell)} \to \Sigma(\ell)$ is étale, it corresponds to a sheaf of ordered abelian groups $\mathcal{I}_{\Sigma(\ell)}$ on $\Sigma(\ell)$, and $\Sigma$ can be regarded as a finite covering of $\Sigma(\ell)$ contained in $(\mathcal{I}_{\Sigma(\ell)})^{\text{ét}}$.

***Proposition 3.18*** **(Intrinsic version of Prop. 3.7).** — *Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be a (possibly ramified) Stokes-filtered local system on $S^1$ (in particular we have a subsheaf $\mathscr{L}_\leqslant$ of $\mu^{-1}\mathscr{L}$) and let $\Sigma \subset \mathcal{I}^{\text{ét}}$ be the support of $\operatorname{gr}\mathscr{L}$. For $\ell \in \mathbb{Q}_+$,*

(1) *the subsheaf* $\mathrm{Tr}_{\leqslant q_\ell}(\mathscr{L}_\leqslant, \mu_\ell^{-1}\mathscr{L})$ *of* $\mu_\ell^{-1}\mathscr{L}$ *(see Lemma* 1.32*) defines a pre-*$\mathcal{I}(\ell)$*-filtration of* $\mathscr{L}$ *which is in fact a* $\mathcal{I}(\ell)$*-filtration, that we denote by* $(\mathscr{L}, \mathscr{L}_{[\bullet]_\ell})$*;*

(2) *the sheaf* $\mathrm{gr}_{[\bullet]_\ell}\mathscr{L}$ *is supported on* $\Sigma(\ell) := q_\ell(\Sigma)$ *and is a local system on* $\Sigma(\ell)$*;*

(3) *the sheaf* $\mathscr{L}_\leqslant$ *induces a pre-*$\mathcal{I}_{\Sigma(\ell)}$*-filtration on* $q_\ell^{-1}\mathrm{gr}_{[\bullet]_\ell}\mathscr{L}$*, which is a* $\mathcal{I}_{\Sigma(\ell)}$*-filtration denoted by* $(\mathrm{gr}_{[\bullet]_\ell}\mathscr{L}, (\mathrm{gr}_{[\bullet]_\ell}\mathscr{L})_\bullet)$*, and* $\mathrm{gr}(\mathrm{gr}_{[\bullet]_\ell}\mathscr{L})$ *is a local system on* $\Sigma$*; this local system has locally the same rank as the local system* $\mathrm{gr}\,\mathscr{L}$*.*

*Conversely, let* $\mathscr{L}_\leqslant \subset \mu^{-1}\mathscr{L}$ *be a pre-*$\mathcal{I}$*-filtration of* $\mathscr{L}$*. Let us assume that* $\mathrm{Tr}_{\leqslant q_\ell}(\mathscr{L}_\leqslant, \mu_\ell^{-1}\mathscr{L})$ *is a* $\mathcal{I}(\ell)$*-filtration of* $\mathscr{L}$ *and let us denote by* $\Sigma(\ell)$ *the support of* $\mathrm{gr}_{[\bullet]_\ell}\mathscr{L}$*. Let us also assume that the pre-*$\mathcal{I}_{\Sigma(\ell)}$*-filtration induced by* $\mathscr{L}_\leqslant$ *on the local system* $\mathrm{gr}_{[\bullet]_\ell}\mathscr{L}$ *(on* $\Sigma(\ell)$*) is a* $\mathcal{I}_{\Sigma(\ell)}$*-filtration, and let us denote by* $\Sigma$ *the support of* $\mathrm{gr}(\mathrm{gr}_{[\bullet]_\ell}\mathscr{L})$*. Then* $\mathscr{L}_\leqslant$ *is a* $\mathcal{I}$*-filtration of* $\mathscr{L}$ *and* $\mathrm{gr}\,\mathscr{L}$ *is supported on* $\Sigma$ *and locally isomorphic to* $\mathrm{gr}(\mathrm{gr}_{[\bullet]_\ell}\mathscr{L})$*.* □

**3.f. Comments.** — The notion of level structure is implicitly present since long in the theory of meromorphic differential equations (see [**29, 4**]). It appears as the "dévissage Gevrey" in [**68**]. Later, it was an important tool for the construction of the moduli space of meromorphic connections [**2**], as emphasized in [**17**], and was strongly related to the multisummation property of solutions of differential equations (see e.g. [**55**] and the references given therein). Note also that the basic vanishing property proved during the proof of Lemma 3.11 is already present in [**17**] and explained in the one-slope case in [**46**, §5], see also [**55**, Lemma 4.3]. It is related to the Watson lemma (see loc. cit.). It is related to the existence of asymptotic solutions of a differential equation in "large" Stokes sectors, see e.g. [**4**] and in particular §B for historical references.

This level structure is not a filtration of the filtered local system. It is the structure induced by a suitable filtration of the indexing sheaf $\mathcal{I}$. A visually similar structure appears in knot theory when considering iterated torus knots, for instance the knots obtained from a singularity of an irreducible germ of complex plane curve. The algebraic structure related to the level structure is the "dévissage" by Gevrey exponents or the multi-summability property. Similarly, the algebraic structure related to iterated torus knots of singularities of plane curves are the Puiseux aproximations.

# LECTURE 4

# STOKES-PERVERSE SHEAVES ON CURVES

**Summary.** In this lecture, we introduce the global notion of Stokes-perverse sheaves on a Riemann surface. The Riemann-Hilbert correspondence of the next lecture will be an equivalence between holonomic $\mathscr{D}$-modules on the Riemann surface and Stokes-perverse sheaves on it. If $\boldsymbol{k}$ is a subfield of $\mathbb{C}$, this allows one to speak of a $\boldsymbol{k}$-structure on a holonomic $\mathscr{D}$-module when the corresponding Stokes-perverse is defined over $\boldsymbol{k}$. The constructions in this chapter are taken from [**15, 16, 49**].

**4.a. The setting.** — Let $X$ be a Riemann surface and let $D$ be a discrete set of points of $X$. We set $X^* = X \smallsetminus D$. We denote by $\varpi : \widetilde{X}(D) \to X$ the real oriented blowing up of $X$ at each of the points of $D$. When $D$ is fixed, we set $\widetilde{X} = \widetilde{X}(D)$. Locally near a point of $D$, we are in the situation of Example 1.6. We set $\partial\widetilde{X} = \varpi^{-1}(D)$ and we denote by $j_\partial : X^* \hookrightarrow \widetilde{X}$ and $i_\partial : \partial\widetilde{X} \hookrightarrow \widetilde{X}$ the inclusions. We denote by $\mathcal{I}$ the sheaf which is zero on $X^*$ and is equal to $\bigcup_d \mathcal{I}_{d,x_o}$ on[1] $S^1_{x_o} := \varpi^{-1}(x_o)$ for $x_o \in D$, where $\mathcal{I}_{d,x_o}$ is the sheaf introduced in Remark 2.23. Notice that $\mathcal{I}^{\text{ét}}$ is not Hausdorff, since we consider $\mathcal{I}$ as a sheaf on $\widetilde{X}$, not on $\partial\widetilde{X}$. Since $\mu : \mathcal{I}^{\text{ét}} \to \widetilde{X}$ is a homeomorphism above $X^*$, we shall denote by $\widetilde{\jmath} : X^* = \mathcal{I}^{\text{ét}}_{|X^*} \hookrightarrow \mathcal{I}^{\text{ét}}$ the open inclusion, by $\widetilde{\imath} : \mathcal{I}^{\text{ét}}_{|\partial\widetilde{X}} \hookrightarrow \mathcal{I}^{\text{ét}}$ the closed inclusion, so that $\mu \circ \widetilde{\jmath} = j_\partial$ and we have a commutative diagram

$$\begin{tikzcd}
\mathcal{I}^{\text{ét}}_{|\partial\widetilde{X}} \arrow[r, hook, "\widetilde{\imath}"] \arrow[d, "\mu_\partial"'] & \mathcal{I}^{\text{ét}} \arrow[d, "\mu"] \\
\partial\widetilde{X} \arrow[r, hook, "i_\partial"] & \widetilde{X}
\end{tikzcd} \tag{4.1}$$

***Remark 4.2*** **(Sheaves on $\mathcal{I}^{\text{ét}}$).** — The sheaf $\mathcal{I}$ does not satisfy the Hausdorff property only because it is zero on $X^*$, but it is Hausdorff when restricted to $\partial\widetilde{X}$. It may therefore be convenient to describe a sheaf $\mathscr{F}_{\leqslant}$ on $\mathscr{I}^{\text{ét}}$, up to unique isomorphism inducing the identity on $X^*$ and $\partial\widetilde{X}$, as a triple $(\mathscr{F}^*, \mathscr{L}_{\leqslant}, \nu)$ consisting of a sheaf $\mathscr{F}^*$ on $X^*$, a sheaf $\mathscr{L}_{\leqslant}$ on $\mathscr{I}^{\text{ét}}_{|\partial\widetilde{X}}$, and a morphism $\nu : \mathscr{L}_{\leqslant} \to \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$. Then one can use Remark 1.9 to describe $\mathscr{L}_{\leqslant}$.

---

[1] The notation $S^1_{x_o}$ should not be confused with the notation $S^1_x$ of §2.b.

In a similar way, a pre-$\mathscr{I}$-filtration $\mathscr{F}_{\leqslant}$ on $\mathscr{I}^{\text{ét}}$ corresponds in a one-to-one way to a triple $(\mathscr{F}^*, \mathscr{L}_{\leqslant}, \nu)$, where $\mathscr{L}_{\leqslant}$ is a pre-$\mathscr{I}_{\partial\widetilde{X}}$-filtration and the morphisms $\beta_{\varphi\leqslant\psi}\mathscr{L}_{\leqslant\varphi} \to \mathscr{L}_{\leqslant\psi}$ commute with $\nu$. Indeed, for a pre-$\mathscr{I}$-filtration, the restriction to $X^*$ of the morphisms $\beta_{\varphi\leqslant\psi}\mathscr{F}_{\leqslant\varphi} \to \mathscr{F}_{\leqslant\psi}$ have to be equal to the identity, because $\varphi = \psi = 0$ there.

**4.b. The category of Stokes-ℂ-constructible sheaves on $\widetilde{X}$.** — The category of St-ℂ-constructible sheaves will be a subcategory of the category $\mathrm{Mod}(\boldsymbol{k}_{\mathscr{I}^{\text{ét}},\leqslant})$ of pre-$\mathscr{I}$-filtrations introduced in §1.c, and we use the notation introduced there, with $\mathscr{I}$ as above. Recall that ℂ-constructible means that the stratification giving constructibility is ℂ-analytic in $X$. The coefficient field is $\boldsymbol{k}$.

Let $\mathscr{F}^*$ be a ℂ-constructible sheaf on $X^*$ with singularity set $S$. We assume that $S$ is locally finite on $X$, so that it does not accumulate on $D$. Then $\mathscr{F}^*$ is a locally constant sheaf of finite dimensional $\boldsymbol{k}$-vector spaces on $X^* \smallsetminus S$, and in particular in the punctured neighbourhood of each point of $D$. This implies that $\boldsymbol{R}j_{\partial,*}\mathscr{F}^* = j_{\partial,*}\mathscr{F}^*$ is a local system on $\widetilde{X} \smallsetminus S$, as well as its restriction to each $S^1_{x_o}$, $x_o \in D$. Similarly, $\boldsymbol{R}\widetilde{\jmath}_*\mathscr{F}^* = \widetilde{\jmath}_*\mathscr{F}^*$ is a local system on $\mathscr{I}^{\text{ét}} \smallsetminus S$, since the property is local on $\mathscr{I}^{\text{ét}}$ and $\mu$ is a local homeomorphism. Moreover, the adjunction morphism $\mu^{-1}\mu_* \to \mathrm{Id}$ induces a morphism $\mu^{-1}j_{\partial,*}\mathscr{F}^* \to \widetilde{\jmath}_*\mathscr{F}^*$, which is easily seen to be an isomorphism. We will identify both sheaves.

We will set $\mathscr{L} = i_\partial^{-1}j_{\partial,*}\mathscr{F}^*$, which is a local system on $\partial\widetilde{X}$. We thus have $\mu_\partial^{-1}\mathscr{L} = \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$.

***Definition 4.3*** (St-ℂ-**constructible sheaves**). — Let $\mathscr{F}_{\leqslant}$ be a pre-$\mathscr{I}$-filtration, i.e., an object of $\mathrm{Mod}(\boldsymbol{k}_{\mathscr{I}^{\text{ét}},\leqslant})$. We say that $\mathscr{F}_{\leqslant}$ is St-ℂ-constructible if

(1) $\mathscr{F}^* := \widetilde{\jmath}^{-1}\mathscr{F}_{\leqslant}$ is a ℂ-constructible sheaf on $X^*$ whose singularity set $S \subset X^*$ is locally finite in $X$,

(2) the natural morphism $\nu : \mathscr{L}_{\leqslant} := \widetilde{\imath}^{-1}\mathscr{F}_{\leqslant} \to \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$ is *injective*, i.e., the sheaf $\mathscr{F}_{\leqslant}$ does not have a non-zero subsheaf supported in $\mu^{-1}\partial\widetilde{X}$,

(3) the inclusion $\nu : \mathscr{L}_{\leqslant} \hookrightarrow \mu^{-1}\mathscr{L}$ is a Stokes filtration of $\mathscr{L}$.

We will denote by $\mathrm{Mod}_{\text{St-ℂ-c}}(\boldsymbol{k}_{\mathscr{I}^{\text{ét}},\leqslant})$ the full subcategory of $\mathrm{Mod}(\boldsymbol{k}_{\mathscr{I}^{\text{ét}},\leqslant})$ whose objects are St-ℂ-constructible on $\mathscr{I}^{\text{ét}}$.

We can also regard $\mathscr{F}_{\leqslant}$ as a pre-$\mathscr{I}$-filtration of the sheaf $\mathscr{F} := j_{\partial,*}\mathscr{F}^*$, in the sense of Definition 1.28. For any open set $U \subset \widetilde{X}$ and any $\varphi \in \Gamma(U, \mathscr{I})$, $\mathscr{F}_{\leqslant\varphi} := \varphi^{-1}\mathscr{F}_{\leqslant}$ is a sheaf on $U$.

***Lemma 4.4.*** — *The category of* St-$\mathbb{C}$-*constructible sheaves on* $\mathfrak{I}^{\text{ét}}$ *is abelian and stable by extensions in the category of pre-*$\mathfrak{I}$*-filtrations on* $\mathfrak{I}^{\text{ét}}$.

*Proof.* — When restricted to $X^*$, the result is well-known for the category of $\mathbb{C}$-constructible sheaves in the category of sheaves of $\boldsymbol{k}$-vector spaces. It is therefore enough to prove it over $\partial\widetilde{X}$. Let $\lambda : \mathscr{F}_{\leqslant} \to \mathscr{F}'_{\leqslant}$ be a morphism of St-$\mathbb{C}$-constructible sheaves. Its restriction to $\partial\widetilde{X}$ induces a morphism in the category of Stokes-filtered local systems on $\partial\widetilde{X}$. Note also that

$$j_{\partial,*}j_\partial^{-1}\operatorname{Ker}\lambda = \operatorname{Ker} j_{\partial,*}j_\partial^{-1}\lambda \quad \text{and} \quad j_{\partial,*}j_\partial^{-1}\operatorname{Coker}\lambda = \operatorname{Coker} j_{\partial,*}j_\partial^{-1}\lambda.$$

According to Theorem 3.1 and the previous identification, the kernel and cokernel of $\lambda_{|\partial\widetilde{X}}$ inject into $\mu^{-1}\operatorname{Ker}[\mathscr{L} \to \mathscr{L}']$ and $\mu^{-1}\operatorname{Coker}[\mathscr{L} \to \mathscr{L}']$, and define Stokes-filtered local systems. Therefore, $\operatorname{Ker}\lambda$ and $\operatorname{Coker}\lambda$ (taken in $\operatorname{Mod}(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}},\leqslant})$) belong to $\operatorname{Mod}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}},\leqslant})$.

Similarly, the stability by extension is clear on $X^*$, and an extension of two St-$\mathbb{C}$-constructible sheaves satisfies 4.3(2) above. Then it satisfies 4.3(3) according to Theorem 3.1. ☐

Note that, as a consequence, $\operatorname{Coker}\lambda$ does not have any nonzero subsheaf supported on $\mu^{-1}\partial\widetilde{X}$, so in particular $\lambda$ is strict with respect to the support condition 4.3(2).

***Remark 4.5*** **($\mathscr{H}om$).** — Given two St-$\mathbb{C}$-constructible sheaves $\mathscr{F}_{\leqslant}, \mathscr{F}'_{\leqslant}$, one can define a St-$\mathbb{C}$-constructible sheaf $\mathscr{H}om(\mathscr{F},\mathscr{F}')_{\leqslant}$ whose generic part is the constructible sheaf $\mathscr{H}om(\mathscr{F}^*,\mathscr{F}'^*)$, and whose restriction to $\mathfrak{I}^{\text{ét}}_{\partial\widetilde{X}}$ is $\mathscr{H}om(\mathscr{L},\mathscr{L}')_{\leqslant}$ (see Definition 2.13). Firstly, one remarks that since $\widetilde{\jmath}_*\mathscr{F}^*$ and $\widetilde{\jmath}_*\mathscr{F}'^*$ are local systems, the natural morphism $\mathscr{H}om(\widetilde{\jmath}_*\mathscr{F}^*,\widetilde{\jmath}_*\mathscr{F}'^*) \to \widetilde{\jmath}_*\mathscr{H}om(\mathscr{F}^*,\mathscr{F}'^*)$ is an isomorphism. The natural injection $\mathscr{H}om(\mathscr{L},\mathscr{L}')_{\leqslant} \hookrightarrow \mathscr{H}om(\mathscr{L},\mathscr{L}') = \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{H}om(\mathscr{F}^*,\mathscr{F}'^*)$ defines thus the desired St-$\mathbb{C}$-constructible sheaf.

Let $\mathscr{F}_{\leqslant}$ be a Stokes-$\mathbb{C}$-constructible sheaf on $\widetilde{X}$. Since $\mathfrak{I}$ is Hausdorff on $\partial\widetilde{X}$, the subsheaf $\mathscr{L}_<$ of $\mathscr{L}_{\leqslant}$ is well defined, and we can consider the triple $(\mathscr{F}^*,\mathscr{L}_<,\nu_{|\mathscr{L}_<})$, which defines a subsheaf $\mathscr{F}_{\prec}$ of $\mathscr{F}_{\leqslant}$, according to Remark 4.2 (we avoid the notation $\mathscr{F}_<$, which has a different meaning over $X^*$).

***Definition 4.6*** **(co-St-$\mathbb{C}$-constructible sheaves).** — We will say that a pre-$\mathfrak{I}$-filtration $\mathscr{F}_{\prec}$ is a co-Stokes-$\mathbb{C}$-constructible sheaf if it satisfies the properties 4.3(1), 4.3(2) and

(3′) the inclusion $\nu : \mathscr{L}_< \hookrightarrow \mu^{-1}\mathscr{L}$ is a Stokes co-filtration of $\mathscr{L}$ (see Remark 1.42).

This defines a full subcategory $\operatorname{Mod}_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}},\leqslant})$ of $\operatorname{Mod}(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}},\leqslant})$.

We similarly have:

***Lemma 4.7.*** — *The category of co*-St-$\mathbb{C}$-*constructible sheaves on* $\mathfrak{I}^{\text{ét}}$ *is abelian and stable by extensions in the category of pre-*$\mathfrak{I}$*-filtrations on* $\mathfrak{I}^{\text{ét}}$. ☐

***Exercise 4.8.*** — From a co-Stokes-$\mathbb{C}$-constructible sheaf $\mathscr{F}_{\prec}$, reconstruct the Stokes-$\mathbb{C}$-constructible sheaf $\mathscr{F}_{\leqslant}$ which defines it (this mainly consists in reconstructing $\mathscr{L}_{\leqslant}$ from $\mathscr{L}_{<}$, see Remark 1.42).

Lastly, there is a simpler subcategory of $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ which will also be useful, namely that of graded $\mathbb{C}$-constructible sheaves on $\mathcal{I}^{\text{ét}}$. By definition, such an object takes the form $\widetilde{\imath}_*(\mu_{\partial,!}\mathscr{G})_{\leqslant}$, where $\mathscr{G}$ is a locally constant sheaf of finite dimensional $\boldsymbol{k}$-vector spaces on $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$, on the support of which $\mu_\partial$ is proper. It is easy to check that it is abelian and stable by extensions in $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. The following is then clear:

***Lemma 4.9.*** — *Let $\mathscr{F}_{\leqslant}$ be a Stokes-$\mathbb{C}$-constructible sheaf on $\mathcal{I}^{\text{ét}}$ and let $\mathscr{F}_{\prec}$ be the associated co-Stokes-$\mathbb{C}$-constructible sheaf, together with the natural inclusion $\mathscr{F}_{\prec}\hookrightarrow\mathscr{F}_{\leqslant}$. Then the quotient (in $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$) is a graded $\mathbb{C}$-constructible sheaf on $\mathcal{I}^{\text{ét}}$.* ☐

***Remark 4.10.*** — There is no nonzero morphism (in the category $\operatorname{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$) from a graded $\mathbb{C}$-constructible sheaf $\mathscr{G}$ on $\mathcal{I}^{\text{ét}}$ to a (co-)Stokes-$\mathbb{C}$-constructible sheaf $\mathscr{F}_{\leqslant}$ (or $\mathscr{F}_{\prec}$). Indeed, presenting $\mathscr{F}_{\leqslant}$ as a triple $(\mathscr{F}^*,\mathscr{L}_{\leqslant},\nu)$ and $\mathscr{G}$ by $(0,\widetilde{\imath}^{-1}\mathscr{G},0)$, a morphism $\mathscr{G}\to\mathscr{F}_{\leqslant}$ consists of a morphism $\widetilde{\imath}^{-1}\mathscr{G}\to\mathscr{L}_{\leqslant}$ which is zero when composed with $\nu$. Since $\nu$ is injective, after 4.3(2), the previous morphism is itself zero. The same argument applies to $\mathscr{G}\to\mathscr{F}_{\prec}$. In other words, since $\mathscr{F}_{\leqslant}$ (resp. $\mathscr{F}_{\prec}$) does not have any non-zero subsheaf supported on $\mu^{-1}\partial\widetilde{X}$, and since $\mathscr{G}$ is supported on this subset, any morphism from $\mathscr{G}$ to $\mathscr{F}_{\leqslant}$ (resp. $\mathscr{F}_{\prec}$) is zero.

## 4.c. Derived categories and duality

*Derived categories.* — We denote by $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ the bounded derived category of pre-$\mathcal{I}$-filtrations, and by $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ the full subcategory of $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ consisting of objects having St-$\mathbb{C}$-constructible cohomology. By Lemma 4.4, the category $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ is a full triangulated subcategory of $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. We will now define a $t$-structure on $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ which restricts to the usual(2) perverse one on $X^*$.

The subcategory ${}^p\!D^{b,\leqslant 0}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ consists of objects $\mathscr{F}_{\leqslant}$ satisfying $\mathscr{H}^j(\mathscr{F}_{\leqslant})=0$ for $j>-1$ and $\mathscr{H}^j(i_{x_o}^{-1}\mathscr{F}_{\leqslant})=0$ for any $x_o\in X^*$ and $j>0$.

Similarly, the subcategory ${}^p\!D^{b,\geqslant 0}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ consists of objects $\mathscr{F}_{\leqslant}$ satisfying $\mathscr{H}^j(\mathscr{F}_{\leqslant})=0$ for $j<-1$ and $\mathscr{H}^j(i_{x_o}^!\mathscr{F}_{\leqslant})=0$ for any $x_o\in X^*$ and $j<0$.

The pair $\big({}^p\!D^{b,\leqslant 0}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}),{}^p\!D^{b,\geqslant 0}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})\big)$ is a $t$-structure on $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ (this directly follows from the result on $X^*$).

***Definition 4.11.*** — The category $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ of perverse sheaves on $X^*$ with a Stokes structure at $\partial\widetilde{X}$ (that we also call *Stokes-perverse sheaves* on $\widetilde{X}$) is the heart of this $t$-structure.

---

(2) We refer for instance to [**20**] for basic results on perverse sheaves; recall that the constant sheaf supported at one point is perverse, and a local system shifted by one is perverse, see e.g. [**20**, Ex. 5.2.23].

This is an abelian category (see [**6**], [**35**, §10.1]).

***Remark 4.12.*** — Objects of $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ behave like sheaves, i.e., can be reconstructed, up to isomorphism, from similar objects on each open set of an open covering of $X$, together with compatible gluing isomorphisms. An object of $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ is a sheaf shifted by 1 in the neighbourhood of $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$ and is a perverse sheaf (in the usual sense) on $X^*$.

Notice also that the objects of $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ which have no singularity on $X^*$ are sheaves (shifted by one). We will call them *smooth* St-$\mathbb{C}$-constructible sheaves or smooth Stokes-perverse sheaves (depending on the shifting convention).

However, it will be clear below that this presentation is not suitable for defining a duality functor on it. We will use a presentation which also takes into account $\mathscr{F}_{\prec}$ and $\operatorname{gr}\mathscr{F}$. Anticipating on the Riemann-Hilbert correspondence, $\mathscr{F}_{\leqslant}$ corresponds to horizontal sections of a connection having moderate growth at $\partial\widetilde{X}$, while $\mathscr{F}_{\prec}$ corresponds with horizontal sections having rapid decay there, and duality pairs moderate growth with rapid decay.

Note that similar arguments apply to co-Stokes-$\mathbb{C}$-constructible sheaves, and we get a $t$-structure $\big({}^p\!D^{b,\leqslant 0}_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}), {}^p\!D^{b,\geqslant 0}_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})\big)$ on $D^b_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$.

Lastly, the full subcategory $D^b_{\text{gr-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ of $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ whose objects have graded $\mathbb{C}$-constructible cohomology is endowed with the $t$-structure induced from the shifted natural one $(D^{b,\leqslant -1}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}), D^{b,\geqslant -1}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}))$.

*Duality.* — Recall that the dualizing complex on $\widetilde{X}$ is $j_{\partial,!}\boldsymbol{k}_{X^*}[2]$ (see e.g. [**35**]) and (see Corollary 1.24) we have a functor $\boldsymbol{R}\mathscr{H}om(\iota^{-1}\bullet, \mu^{-1}j_{\partial,!}\boldsymbol{k}_{X^*}[2])$ from the category $D^{b,\text{op}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ to $D^{+}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. We will denote it by $\boldsymbol{D}$. We will prove the following.

***Proposition 4.13.*** — *The duality functor $\boldsymbol{D}$ induces functors $\boldsymbol{D} : D^{b,\text{op}}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \to D^b_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ and $\boldsymbol{D} : D^{b,\text{op}}_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \to D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ which are $t$-exact, and there are isomorphisms of functors $\operatorname{Id} \simeq \boldsymbol{D}\circ\boldsymbol{D}$ in both $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ and $D^b_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$.*

*On the other hand, it induces a functor $\boldsymbol{D} : D^{b,\text{op}}_{\text{gr-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \to D^b_{\text{gr-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ such that $\boldsymbol{D}\circ\boldsymbol{D} \equiv \operatorname{Id}$.*

Let $\mathscr{F}_{\leqslant}$ be an object of $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. For any open set $U\subset\widetilde{X}$ and any $\varphi\in\Gamma(U,\mathcal{I})$, $\mathscr{F}_{\leqslant\varphi} := \varphi^{-1}\mathscr{F}_{\leqslant}$ is an object of $D^b(\boldsymbol{k}_U)$.

***Lemma 4.14.*** — *If $\mathscr{F}_{\leqslant}$ is a Stokes-$\mathbb{C}$-constructible sheaf on $\mathcal{I}^{\text{ét}}$, then for any $\varphi\in\Gamma(U,\mathcal{I})$, $\mathscr{F}_{\leqslant\varphi}$ is $\mathbb{R}$-constructible on $U$. The same result holds for a co-Stokes-$\mathbb{C}$-constructible sheaf $\mathscr{F}_{\prec}$.*

*Proof.* — It suffices to check this on $U\cap\partial\widetilde{X}$, where it follows from Lemma 2.7. ☐

If $\mathscr{F}_{\leqslant}$ is a Stokes-$\mathbb{C}$-constructible sheaf on $\mathcal{I}^{\text{ét}}$, we present it as a triple $(\mathscr{F}^*, \mathscr{L}_{\leqslant}, \nu)$, with $\nu : \mathscr{L}_{\leqslant} \hookrightarrow \mu^{-1}\mathscr{L} := \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$ defining a Stokes filtration of $\mathscr{L} := i_{\partial}^{-1}j_{\partial,*}\mathscr{F}^*$. We have a natural morphism $\nu' : \mu^{-1}\mathscr{L} = \widetilde{\imath}^!\widetilde{\jmath}_!\mathscr{F}^*[1] \to \widetilde{\imath}^!\mathscr{F}_{\leqslant}[1] = \mu^{-1}\mathscr{L}/\mathscr{L}_{\leqslant}$. Then the injectivity of $\nu$ is equivalent to the surjectivity of $\nu'$. Indeed, note that the injectivity condition 4.3(2) means that, for any open set $U$ and any $\varphi \in \Gamma(U, \mathcal{I})$, the distinguished triangle $i_\partial^{-1}\mathscr{F}_{\leqslant\varphi} \to i_\partial^{-1}j_{\partial,*}\mathscr{F}^* \to i_\partial^!\mathscr{F}_{\leqslant\varphi}[1] \xrightarrow{+1}$ reduces to the exact sequence $0 \to \mathscr{L}_{\leqslant\varphi} \to \mathscr{L}_{|U} \to \mathscr{L}_{|U}/\mathscr{L}_{\leqslant\varphi} \to 0$, that is, $i_\partial^!\mathscr{F}_{\leqslant\varphi}[1]$ has cohomology in degree 0 at most, and $\mathscr{H}^0(i_\partial^!\mathscr{F}_{\leqslant\varphi}[1]) = \mathscr{L}_{|U}/\mathscr{L}_{\leqslant\varphi}$. Argue similarly for the surjectivity.

***Lemma 4.15.*** — *If* $\mathscr{F}_{\leqslant}$ *is a Stokes-*$\mathbb{C}$*-constructible sheaf (resp.* $\mathscr{F}_{\prec}$ *is a co-Stokes-*$\mathbb{C}$*-constructible sheaf) on* $\mathcal{I}^{\text{ét}}$*, then on* $\widetilde{X} \smallsetminus \operatorname{Sing}\mathscr{F}^*$*,* $\mathscr{F}_{\prec}^\vee := \boldsymbol{R}\mathscr{H}om(\iota^{-1}\mathscr{F}_{\leqslant}, \widetilde{\jmath}_!\boldsymbol{k}_{X^*})$ *(resp.* $\mathscr{F}_{\leqslant}^\vee := \boldsymbol{R}\mathscr{H}om(\iota^{-1}\mathscr{F}_{\prec}, \widetilde{\jmath}_!\boldsymbol{k}_{X^*})$*) has cohomology in degree* 0 *at most and is equal to the co-Stokes-*$\mathbb{C}$*-constructible sheaf* $(\mathscr{F}^{*,\vee}, (\mathscr{L}^\vee)_<, \nu^\vee)$ *(resp. the Stokes-*$\mathbb{C}$*-constructible sheaf* $(\mathscr{F}^{*,\vee}, (\mathscr{L}^\vee)_{\leqslant}, \nu^\vee)$*), where* $\nu^\vee$ *is the morphism dual to* $\nu'$*. Similarly,* $\nu^{\vee,\prime}$ *is the morphism dual to* $\nu$*.*

*Proof.* — If $U$ is an open set in $\widetilde{X}$ and $\varphi \in \Gamma(U, \mathcal{I}^{\text{ét}})$, we will set

$$\boldsymbol{D}(\mathscr{F}_{\leqslant\varphi}) = \boldsymbol{R}\mathscr{H}om_{\boldsymbol{k}}(\mathscr{F}_{\leqslant\varphi}, j_{\partial,!}\boldsymbol{k}_{U^*}[2]) \quad \text{and} \quad \boldsymbol{D}'(\mathscr{F}_{\leqslant\varphi}) = \boldsymbol{R}\mathscr{H}om_{\boldsymbol{k}}(\mathscr{F}_{\leqslant\varphi}, j_{\partial,!}\boldsymbol{k}_{U^*}).$$

As a consequence of Lemma 4.14 we have (see [**35**, Prop. 3.1.13]),

$$\begin{aligned} i_\partial^!\boldsymbol{D}(\mathscr{F}_{\leqslant\varphi}) &= \boldsymbol{R}\mathscr{H}om(i_\partial^{-1}\mathscr{F}_{\leqslant\varphi}, i_\partial^! j_{\partial,!}\boldsymbol{k}[2]) \\ &= \boldsymbol{R}\mathscr{H}om(i_\partial^{-1}\mathscr{F}_{\leqslant\varphi}, \boldsymbol{k}[1]) =: \boldsymbol{D}(i_\partial^{-1}\mathscr{F}_{\leqslant\varphi}), \end{aligned} \tag{4.16}$$

since $\boldsymbol{k}_{\partial\widetilde{X}}[1]$ is the dualizing complex on $\partial\widetilde{X}$, and by biduality we have $\boldsymbol{D}(i_\partial^!\mathscr{F}_{\leqslant\varphi}) = i_\partial^{-1}\boldsymbol{D}(\mathscr{F}_{\leqslant\varphi})$ (see [**35**, Prop. 3.4.3]). This can also be written as

$$i_\partial^!\boldsymbol{D}'(\mathscr{F}_{\leqslant\varphi})[1] = \boldsymbol{D}'(i_\partial^{-1}\mathscr{F}_{\leqslant\varphi}) \quad \text{and} \quad i_\partial^{-1}\boldsymbol{D}'(\mathscr{F}_{\leqslant\varphi}) = \boldsymbol{D}'(i_\partial^!\mathscr{F}_{\leqslant\varphi}[1]).$$

For the proof, we can assume that $\operatorname{Sing}\mathscr{F}^* = \varnothing$. We will prove the Stokes case, the co-Stokes case being similar.

For the first part of the statement, it suffices to show that $i_\partial^{-1}\boldsymbol{D}'(\mathscr{F}_{\leqslant\varphi})$ is a sheaf. We have $i_\partial^{-1}\boldsymbol{D}'(\mathscr{F}_{\leqslant\varphi}) = \boldsymbol{D}'(i_\partial^!\mathscr{F}_{\leqslant\varphi}[1]) = \boldsymbol{D}'(\mathscr{L}_{|U}/\mathscr{L}_{\leqslant\varphi})$, as remarked above, and thus $i_\partial^{-1}\boldsymbol{D}'(\mathscr{F}_{\leqslant\varphi})$ is a sheaf, according to Lemma 2.17. Therefore, $\mathscr{F}_{\prec}^\vee = \mathscr{H}om(\iota^{-1}\mathscr{F}_{\leqslant}, \widetilde{\jmath}_!\boldsymbol{k}_{X^*})$. It remains to check that the morphism $\beta_{\leqslant}p_1^{-1}\mathscr{F}_{\prec}^\vee \to p_2^{-1}\mathscr{F}_{\prec}^\vee$ given by Corollary 1.24 induces $\nu^\vee$ (dual of $\nu'$) and that $\nu^{\vee,\prime}$ corresponds to the dual of $\nu$. The second point amounts to proving that, for any open set $U$ in $\widetilde{X}$ and any $\varphi, \psi \in \Gamma(U, \mathcal{I})$, the natural morphism $\mathscr{H}om(\mathscr{F}_{\leqslant-\varphi}, j_{\partial,!}\boldsymbol{k}_U)_{U_{\varphi,\psi}} \to \mathscr{H}om(\mathscr{F}_{\leqslant-\psi}, j_{\partial,!}\boldsymbol{k}_U)_{U_{\varphi,\psi}}$ induces the dual of $\nu$ after $i_\partial^!$. This follows from (4.16). The assertion for $\nu^\vee$ follows by biduality. ☐

*Proof of Proposition* 4.13. — The first claim is now a direct consequence of Lemma 4.15. For any $U$ and any $\varphi \in \Gamma(U, \mathcal{I})$ we have the bi-duality isomorphism after applying $\varphi^{-1}$, according to the standard result for $\mathbb{R}$-constructible sheaves and Lemma 4.14. This isomorphism is clearly compatible with restrictions of open sets, hence gives $\mathrm{Id} \simeq \boldsymbol{D} \circ \boldsymbol{D}$.

For the graded case, the situation is simpler, as we work with local systems on $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$ extended by zero on $\mathcal{I}^{\text{ét}}$. □

**4.d. The category of Stokes-perverse sheaves on $\widetilde{X}$.** — In the case of Stokes-filtered local systems on $\partial\widetilde{X}$, the Stokes and co-Stokes aspect (namely $\mathscr{L}_{\leqslant}$ and $\mu^{-1}\mathscr{L}/\mathscr{L}_{<}$) can be deduced one from the other. The same remark applies to (co-)Stokes-$\mathbb{C}$-constructible *sheaves*. However, such a correspondence Stokes ⇔ co-Stokes heavily depends on sheaves operations. In the derived category setting, we will keep them together.

It will be useful to consider the $t$-categories $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ etc. as subcategories of the same one. We can choose the following one. By a $\mathbb{C}$-$\mathbb{R}$-constructible sheaf on $\mathcal{I}^{\text{ét}}$ we will mean a sheaf $\mathscr{G}$ on $\mathcal{I}^{\text{ét}}$ such that for any open set $U \subset \widetilde{X}$ and any $\varphi \in \Gamma(U, \mathcal{I})$, $\varphi^{-1}\mathscr{G}$ is $\mathbb{R}$-constructible on $U$ and $\mathbb{C}$-constructible on $U^* = U \cap X^*$ with locally finite singularity set on $U$. The category $\text{Mod}_{\mathbb{C}\text{-}\mathbb{R}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ is the full subcategory of $\text{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ whose underlying sheaf is $\mathbb{C}$-$\mathbb{R}$-constructible. According to Lemma 4.14, it contains the full subcategories of (co-/gr-)St-constructible sheaves. It is abelian and stable by extensions in $\text{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. The full triangulated subcategory $D^b_{\mathbb{C}\text{-}\mathbb{R}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ of $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ is equipped with a $t$-structure defined as in the beginning of §4.c. This $t$-structure induces the already defined one on the full triangulated subcategories $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ etc.

***Definition 4.17***. — The category $\text{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ is the full subcategory of the category of distinguished triangles of $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ whose objects consist of distinguished triangles $\mathscr{F}_{\prec} \to \mathscr{F}_{\leqslant} \to \operatorname{gr}\mathscr{F} \xrightarrow{+1}$, where $\mathscr{F}_{\prec}$ is an object of $D^b_{\text{coSt-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, $\mathscr{F}_{\leqslant}$ of $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, and $\operatorname{gr}\mathscr{F}$ of $D^b_{\text{gr-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. These triangles are also distinguished triangles in $D^b_{\mathbb{C}\text{-}\mathbb{R}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$.

***Proposition 4.18***. — *The category* $\text{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *is triangulated, and endowed with a $t$-structure* $(\text{St}^{\leqslant 0}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}), \text{St}^{\geqslant 0}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}))$ *defined by the property that* $\mathscr{F}_{\prec}, \mathscr{F}_{\leqslant}, \operatorname{gr}\mathscr{F}$ *belong to the* $\leqslant 0$ *(resp.* $\geqslant 0$*) part of their corresponding categories. There is a duality functor* $\boldsymbol{D} : \text{St}^{\text{op}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant}) \to \text{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *satisfying* $\boldsymbol{D} \circ \boldsymbol{D} \simeq \text{Id}$ *and which is compatible to the $t$-structures.*

*Sketch of proof*. — Any (not necessarily distinguished) triangle $\mathscr{F}_{\prec} \to \mathscr{F}_{\leqslant} \to \operatorname{gr}\mathscr{F} \xrightarrow{+1}$ induces a (not necessarily exact) sequence $\cdots \to {}^p\mathscr{H}^k\mathscr{F}_{\prec} \to {}^p\mathscr{H}^k\mathscr{F}_{\leqslant} \to {}^p\mathscr{H}^k\operatorname{gr}\mathscr{F} \to \cdots$ by taking perverse cohomology of objects of $D^b_{\mathbb{C}\text{-}\mathbb{R}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, and, according to Remark 4.10, each connecting morphism ${}^p\mathscr{H}^k\operatorname{gr}\mathscr{F} \to {}^p\mathscr{H}^{k+1}\mathscr{F}_{\prec}$ is zero. One then shows that such a triangle is distinguished if and only if each short sequence as before is exact. The axioms of a triangulated category are then checked for $\text{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ by using this property.

In order to check the $t$-structure property, the only non trivial point is to insert a given object of $\text{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ in a triangle with first term in $\text{St}^{\leqslant 0}$ and third term in $\text{St}^{\geqslant 1}$.

Denoting by $\tau^{\leqslant 0}$ and $\tau^{\geqslant 1}$ the perverse truncation functors in $D^b_{\mathbb{C}\text{-}\mathbb{R}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, it is enough to check that each of these functors preserves objects of $\operatorname{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$. This follows from the cohomological characterization of the distinguished triangles which are objects of $\operatorname{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$.

The assertion on duality follows from Proposition 4.13. □

The heart of this $t$-structure is an abelian category (see [**6**], [**35**]).

***Lemma 4.19***. — *The "forgetting" functor from* $\operatorname{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *to* $D^b_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$*, sending a triangle* $\mathscr{F}_{\prec} \to \mathscr{F}_{\leqslant} \to \operatorname{gr}\mathscr{F} \xrightarrow{+1}$ *to* $\mathscr{F}_{\leqslant}$ *(which is by definition compatible with the t-structures) induces an equivalence of abelian categories from the heart of the t-structure of* $\operatorname{St}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *to* $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$*.*

*Proof*. — For the essential surjectivity, the point is to recover $\mathscr{F}_{\prec}$ and $\operatorname{gr}\mathscr{F}$ from $\mathscr{F}_{\leqslant}$. As our objects can be obtained by gluing from local data, the question can be reduced to a local one near $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$, where we can apply the arguments for sheaves and define $\mathscr{F}_{\prec}$ and $\operatorname{gr}\mathscr{F}$ as in Lemma 4.9.

Similarly, any morphism $\mathscr{F}_{\leqslant} \to \mathscr{F}'_{\leqslant}$ lifts as a morphism between triangles. It remains to show the uniqueness of this lifting. This is a local problem, that we treat for sheaves. The triangle is then an exact sequence, and the morphism $\mathscr{F}_{\prec} \to \mathscr{F}'^{<}$ coming in a morphism between such exact sequences is simply the one induced by $\mathscr{F}_{\leqslant} \to \mathscr{F}'_{\leqslant}$, so is uniquely determined by it. The same argument holds for $\operatorname{gr}\mathscr{F} \to \operatorname{gr}\mathscr{F}'$. □

***Corollary 4.20* (Duality)**. — *The category* $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ *is stable by duality.* □

**4.e. Direct image to $X$.** — Given a sheaf $\mathscr{F}_{\leqslant}$ on $\mathcal{I}^{\text{ét}}$, any section $\varphi \in \Gamma(U, \mathcal{I})$ produces a sheaf $\mathscr{F}_{\leqslant\varphi} := \varphi^{-1}\mathscr{F}_{\leqslant}$ on $U$. We now use more explicitly that $\mathcal{I}$ has a global section 0, meaning that there is a global section $0 : \widetilde{X} \to \mathcal{I}^{\text{ét}}$ of $\mu$, making $\widetilde{X}$ a closed subset of $\mathcal{I}^{\text{ét}}$. Then $\mathscr{F}_{\leqslant 0}$ is a sheaf on $\widetilde{X}$.

***Proposition 4.21***. — *Let* $\mathscr{F}_{\leqslant}$ *be an object of* $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$*. Then* $\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0}$ *is perverse (in the usual sense) on* $X$*.*

*Proof*. — It is a matter of proving that $i_D^{-1}\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0}$ has cohomology in degrees $-1, 0$ at most and $i^!_D\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0}$ in degrees $0, 1$ at most.

We have $i_D^{-1}\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0} = \boldsymbol{R}\Gamma(\partial\widetilde{X}, i_\partial^{-1}\mathscr{F}_{\leqslant 0})$, because $\varpi$ is proper (see [**35**, Prop. 2.5.11]), so, setting $\mathscr{L}_{\leqslant 0} = i_\partial^{-1}\mathscr{F}_{\leqslant 0}[-1]$, the first assertion reduces to $H^k(\partial\widetilde{X}, \mathscr{L}_{\leqslant 0}) = 0$ for $k \neq 0, 1$, which is clear since $\mathscr{L}_{\leqslant 0}$ is a $\mathbb{R}$-constructible sheaf. Similarly, as we have seen after Lemma 4.14, $i^!_\partial(\mathscr{F}_{\leqslant 0}) = i^!_\partial(\mathscr{F}_{\leqslant 0}[-1])[1]$ is a sheaf, so the second assertion is also satisfied (see [**35**, Prop. 3.1.9(ii)]). □

**4.f. Stokes-perverse sheaves on $\underline{\widetilde{X}}$.** — We will prove in the next lecture that Stokes-perverse sheaves on $\widetilde{X}$ correspond to holonomic $\mathscr{D}$-modules on $X$ which are isomorphic to their localization at $D$, and have at most regular singularities on $X^*$. In order to treat arbitrary holonomic $\mathscr{D}_X$-modules, we need to introduce supplementary data at $D$, which are regarded as filling the space $\widetilde{X}$ by discs in order to obtain a topological space $\underline{\widetilde{X}}$, that we describe now. This construction goes back to [**16**] and is developed in [**49**].

For any $x_o \in D$, let $\widehat{X}_{\widehat{x}_o}$ be an open disc with center denoted by $\widehat{x}_o$, and let $\overline{\widehat{X}}_{\widehat{x}_o}$ be the associated closed disc. Denote by $\widehat{\jmath} : \widehat{X}_{\widehat{x}_o} \hookrightarrow \overline{\widehat{X}}_{\widehat{x}_o}$ the open inclusion and $\widehat{\imath} : S^1_{\widehat{x}_o} \hookrightarrow \overline{\widehat{X}}_{\widehat{x}_o}$ the complementary closed inclusion.

We denote by $\underline{\widetilde{X}}$ the topological space (homeomorphic to $X$) obtained by gluing each closed disc $\overline{\widehat{X}}_{\widehat{x}_o}$ to $\widetilde{X}$ along their common boundary $S^1_{\widehat{x}_o} = S^1_{x_o}$ for $x_o \in D$ (see Figure 1).



FIGURE 1. The space $\underline{\widetilde{X}}$ near $S^1_x$ $(x \in D)$

We first define the category $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ to be the category whose objects are triples $(\mathscr{F}_{\leqslant}, \widehat{\mathscr{F}}, \widehat{\nu})$, where $\mathscr{F}_{\leqslant}$ is a pre-$\mathcal{I}$-filtration, i.e., an object of $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$, $\widehat{\mathscr{F}}$ is a sheaf on $\bigsqcup_{x_o \in D} \widehat{X}_{\widehat{x}_o}$, and $\widehat{\nu}$ is a morphism $i_\partial^{-1}\mathscr{F}_{\leqslant 0} \to \widehat{\imath}^{-1}\widehat{\jmath}_*\widehat{\mathscr{F}}$. The morphisms in this category consist of pairs $(\lambda, \widehat{\lambda})$ of morphisms in the respective categories which are compatible with $\widehat{\nu}$ in an obvious way. We have "forgetting" functors to $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})$ (hence also to $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$) and $\mathrm{Mod}(\boldsymbol{k}_{\widehat{X}})$. Then $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ is an abelian category. A sequence in $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ is exact iff the associated sequences in $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}}})$ and $\mathrm{Mod}(\boldsymbol{k}_{\widehat{X}})$ are exact. The associated bounded derived category is denoted by $D^b(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$.

***Definition 4.22*** **(St-$\mathbb{C}$-constructible sheaves on $\underline{\widetilde{X}}$).** — The objects in the category $\mathrm{Mod}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ consist of triples $(\mathscr{F}_{\leqslant}, \widehat{\mathscr{F}}, \widehat{\nu})$, where

(1) $\mathscr{F}_{\leqslant}$ is a St-$\mathbb{C}$-constructible sheaf on $\widetilde{X}$ (see Definition 4.3), defining a Stokes-filtered local system $(\mathscr{L}, \mathscr{L}_\bullet)$ on each $S^1_{x_o}$, $x_o \in D$,

(2) $\widehat{\mathscr{F}}$ is a $\mathbb{C}$-constructible sheaf on $\bigsqcup_{x_o \in D} \widehat{X}_{\widehat{x}_o}$ with singularity at $\widehat{x}_o$ $(x_o \in D)$ at most,

(3) $\widehat{\nu}$ is an isomorphism $\operatorname{gr}_0\mathscr{L}\xrightarrow{\sim}\widehat{\imath}^{-1}\widehat{\jmath}_*\widehat{\mathscr{F}}$.

Morphisms between such triples consist of pairs $(\lambda,\widehat{\lambda})$ of morphisms in the respective categories which are compatible with $\widehat{\nu}$.

By definition, $\mathrm{Mod}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ is a full subcategory of $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$.

***Lemma 4.23.*** — *The category* $\mathrm{Mod}_{\text{St-}\mathbb{C}\text{-c}}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ *is abelian and stable by extensions in* $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$.

*Proof.* — We apply Lemma 4.4 for the $\mathscr{F}_{\leqslant}$-part, a standard result for the $\widehat{\mathscr{F}}$-part, and the compatibility with $\widehat{\nu}$ follows. □

The definition of the category $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ now proceeds exactly as in §4.d, by only adding the information of the usual $t$-structure on the various $\widehat{X}_{\widehat{x}_o}$, $x_o\in D$. We leave the details to the interested reader. As above, a Stokes-perverse sheaf is a sheaf shifted by one away from the singularities in $X^*$ and of $\widehat{D}$.

***Remark 4.24.*** — The previous presentation makes a difference between singularities $S$ in $X^*$ and singularities in $D$. One can avoid this difference, by considering objects which are local systems on $X^*\smallsetminus S$ and by replacing $D$ with $D\cup S$. Then "regular singularities" are the points of $D$ where the set of exponential factors of $\mathscr{F}_{\leqslant}$ reduces to $\{0\}$. This point of view is equivalent to the previous one through the functor $\mathscr{P}$ at those points (see below, Proposition 4.25).

**4.g. Associated perverse sheaf on $X$.** — We now define a functor $\mathscr{P}$ : $\text{St-Perv}_D(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})\to\mathrm{Perv}(\boldsymbol{k}_X)$ between Stokes-perverse sheaves on $\underline{\widetilde{X}}$ and perverse sheaves (in the usual sense) on $X$. This is an extension of the direct image procedure considered in §4.e. Namely, if $(\mathscr{F}_{\leqslant},\widehat{\mathscr{F}},\widehat{\nu})$ is a Stokes-constructible sheaf on $\underline{\widetilde{X}}$, the triple $(\mathscr{F}_{\leqslant 0},\widehat{\mathscr{F}},\widehat{\nu})$ allows one to define a sheaf $\underline{\mathscr{F}}_{\leqslant 0}$ on $\underline{\widetilde{X}}$: indeed, a sheaf on $\underline{\widetilde{X}}$ can be determined up to isomorphism by a morphism $\mathscr{L}_{\leqslant 0}\to\widehat{\imath}^{-1}\widehat{\jmath}_*\widehat{\mathscr{F}}$; we use the composed morphism $\mathscr{L}_{\leqslant 0}\to\operatorname{gr}_0\mathscr{L}\xrightarrow{\widehat{\nu}}\widehat{\imath}^{-1}\widehat{\jmath}_*\widehat{\mathscr{F}}$.

Extending this construction to Stokes-perverse sheaves and taking the direct image by the continuous projection $\underline{\varpi}:\underline{\widetilde{X}}\to X$ which contracts $\overline{\widehat{X}}$ to $D$ is $\mathscr{P}(\mathscr{F}_{\leqslant},\widehat{\mathscr{F}},\widehat{\nu})$. We will however not explicitly use the previous gluing construction in this perverse setting, but an ersatz of it, in order to avoid a precise justification of this gluing. Let us make this more precise.

We first note that the pull-back by the zero section $0:\widetilde{X}\hookrightarrow\mathcal{I}^{\text{ét}}$ defines an exact functor $0^{-1}:\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant})\to\mathrm{Mod}(\boldsymbol{k}_{\widetilde{X}})$ and therefore, taking the identity on the $\widehat{\mathscr{F}}$-part, an exact functor from $\mathrm{Mod}(\boldsymbol{k}_{\mathcal{I}^{\text{ét}},\leqslant,\widehat{D}})$ to the category $\mathrm{Mod}(\boldsymbol{k}_{\underline{\widetilde{X}},\leqslant 0,\widehat{D}})$ consisting of triples $(\mathscr{F}_{\leqslant 0},\widehat{\mathscr{F}},\widehat{\nu})$.

In the following, we implicitly identify sheaves on $D$ and sheaves on $X$ supported on $D$ via $\boldsymbol{R}i_{D,*}$ and we work in $D^b_{\mathbb{C}\text{-c}}(\boldsymbol{k}_X)$. On the one hand, we have a natural composed morphism

$$\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0}\longrightarrow \boldsymbol{R}\Gamma(\partial\widetilde{X},\widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0})\longrightarrow \boldsymbol{R}\Gamma(\partial\widetilde{X},\mathrm{gr}_0\mathscr{F})\xrightarrow[\sim]{\widehat{\nu}}\boldsymbol{R}\Gamma(\partial\widehat{X},\widetilde{\imath}^{-1}\widehat{\jmath}_*\widehat{\mathscr{F}}).$$

On the other hand, we have a distinguished triangle

$$\boldsymbol{R}\Gamma(\widehat{X},\widehat{\jmath}_*\widehat{\mathscr{F}})\longrightarrow \boldsymbol{R}\Gamma(\partial\widehat{X},\widetilde{\imath}^{-1}\widehat{\jmath}_*\widehat{\mathscr{F}})\longrightarrow \boldsymbol{R}\Gamma(\overline{\widehat{X}},\widehat{\jmath}_!\widehat{\mathscr{F}})[1]\xrightarrow{+1}.$$

We deduce a morphism

$$\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0}\longrightarrow \boldsymbol{R}\Gamma(\overline{\widehat{X}},\widehat{\jmath}_!\widehat{\mathscr{F}})[1]=\boldsymbol{R}\Gamma_{\mathrm{c}}(\widehat{X},\widehat{\mathscr{F}})[1].$$

**Proposition 4.25.** — *If* $(\mathscr{F}_{\leqslant},\widehat{\mathscr{F}},\widehat{\nu})$ *is an object of* $\mathrm{St}\text{-}\mathrm{Perv}_D(\boldsymbol{k}_{\jmath_{\text{ét}},\leqslant,\widehat{D}})$*, then the complex* $\boldsymbol{R}\Gamma_{\mathrm{c}}(\widehat{X},\widehat{\mathscr{F}})[1]$ *has (perverse) cohomology in degrees* $-1$ *and* $0$ *at most and the cone of the previous morphism–that we denote by* $\mathscr{P}(\mathscr{F}_{\leqslant},\widehat{\mathscr{F}},\widehat{\nu})[1]$*–has perverse cohomology in degree* $1$ *at most.*

**Corollary 4.26.** — *The functor* $\mathscr{P}:(\mathscr{F}_{\leqslant},\widehat{\mathscr{F}},\widehat{\nu})\mapsto {}^p\mathscr{H}^0\mathscr{P}(\mathscr{F}_{\leqslant},\widehat{\mathscr{F}},\widehat{\nu})$ *is an exact functor from* $\mathrm{St}\text{-}\mathrm{Perv}_D(\boldsymbol{k}_{\jmath_{\text{ét}},\leqslant,\widehat{D}})$ *to* $\mathrm{Perv}(\boldsymbol{k}_X)$. ☐

*Proof of Proposition* 4.25. — We consider the complementary inclusions

$$\widehat{D}\xhookrightarrow{i_{\widehat{D}}}\widehat{X}\xhookleftarrow{j_{\widehat{D}}}\widehat{X}\smallsetminus\{\widehat{D}\}.$$

We have a distinguished triangle $i^!_{\widehat{D}}\widehat{\mathscr{F}}\to\widehat{\mathscr{F}}\to\boldsymbol{R}j_{\widehat{D},*}j^{-1}_{\widehat{D}}\widehat{\mathscr{F}}\xrightarrow{+1}$, to which we apply $\widehat{\jmath}_!$ (recall that $\widehat{\jmath}$ denotes the inclusion $\widehat{X}_{\widehat{x}_o}\hookrightarrow\overline{\widehat{X}}_{\widehat{x}_o}$). This has no effect to the first term, which remains $i^!_{\widehat{D}}\widehat{\mathscr{F}}$ and has cohomology in degrees 0 and 1 at most by the cosupport condition for $\widehat{\mathscr{F}}$. We then have

$$\boldsymbol{H}^{-1}(\overline{\widehat{X}},\widehat{\jmath}_!\widehat{\mathscr{F}})\lhook\joinrel\longrightarrow\boldsymbol{H}^{-1}_C(\widehat{X}^*,j^{-1}_{\widehat{D}}\widehat{\mathscr{F}}),$$

where $C$ is the family of supports in $\widehat{X}^*$ whose closure in $\widehat{X}$ is compact. We are thus reduced to showing that, if $\widehat{\mathscr{L}}$ is a local system on $\widehat{X}^*$, we have $H^0_C(\widehat{X},\widehat{\mathscr{L}})=0$, which is clear. Therefore, $\boldsymbol{R}\Gamma_{\mathrm{c}}(\widehat{X},\widehat{\mathscr{F}})$ has cohomology in degrees 0 and 1 at most, as wanted.

Let us now show that $\mathscr{P}$ is perverse on $X$. Obviously, we only have to check the support and cosupport conditions at $D$, and we will use Proposition 4.21.

For the support condition, we note that $i_D^{-1}\boldsymbol{R}\varpi_*\mathscr{F}_{\leqslant 0}=\boldsymbol{R}\Gamma(\partial\widetilde{X},\widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0})$, and $i_D^{-1}\mathscr{P}[1]$ is the cone of the diagonal morphism below (where $\widehat{\nu}$ is implicitly used):

(4.27)

$$\begin{tikzcd}
& \boldsymbol{R}\Gamma(\partial\widetilde{X},\widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0}) \arrow[d] \arrow[dr] & & \\
\boldsymbol{R}\Gamma(\widehat{X},\widehat{\mathscr{F}}) \arrow[r] & \boldsymbol{R}\Gamma(\partial\widetilde{X},\mathrm{gr}_0\mathscr{F}) \arrow[r] & \boldsymbol{R}\Gamma_{\mathrm{c}}(\widehat{X},\widehat{\mathscr{F}})[1] \arrow[r, "+1"] & {}
\end{tikzcd}$$

This gives an exact sequence

$$\begin{aligned}(4.28)\quad 0 \longrightarrow \mathscr{H}^{-1}(i_D^{-1}\mathscr{P}) \longrightarrow \boldsymbol{H}^{-1}(\partial\widetilde{X}, \widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0}) \longrightarrow \boldsymbol{H}^0_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) \longrightarrow \mathscr{H}^0(i_D^{-1}\mathscr{P})\\ \longrightarrow \boldsymbol{H}^0(\partial\widetilde{X}, \widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0}) \longrightarrow \boldsymbol{H}^1_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) \longrightarrow \mathscr{H}^1(i_D^{-1}\mathscr{P}) \longrightarrow 0\end{aligned}$$

and our claim is that $\boldsymbol{H}^0(\partial\widetilde{X}, \widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0}) \to \boldsymbol{H}^1_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}})$ is onto, because $\boldsymbol{H}^2_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) = 0$ due to the perversity of $\widehat{\mathscr{F}}$. On the one hand, $\boldsymbol{H}^0(\partial\widetilde{X}, \widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0}) \to \boldsymbol{H}^0(\partial\widetilde{X}, \operatorname{gr}_0\mathscr{F})$ is the morphism $H^1(\partial\widetilde{X}, \mathscr{L}_{\leqslant 0}) \to H^1(\partial\widetilde{X}, \operatorname{gr}_0\mathscr{L})$, which is onto since $\mathscr{L}_{<0}$ is a sheaf on $\partial\widetilde{X}$. On the other hand the distinguished horizontal triangle above gives an exact sequence

$$\boldsymbol{H}^0(\partial\widetilde{X}, \operatorname{gr}_0\mathscr{F}) \longrightarrow \boldsymbol{H}^1_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) \longrightarrow \boldsymbol{H}^1(\widehat{X}, \widehat{\mathscr{F}}),$$

and $\boldsymbol{H}^1(\widehat{X}, \widehat{\mathscr{F}}) = \mathscr{H}^1(i_{\widehat{D}}^{-1}\widehat{\mathscr{F}}) = 0$ by the support condition for $\widehat{\mathscr{F}}$.

Let us now check the cosupport condition. We have $\widetilde{\imath}^!\mathscr{F}_{\leqslant 0} = \mathscr{L}/\mathscr{L}_{\leqslant 0}$, as remarked at the end of the proof of Lemma 4.15. We now argue as above, by replacing $\boldsymbol{R}\Gamma(\partial\widetilde{X}, \widetilde{\imath}^{-1}\mathscr{F}_{\leqslant 0})$ with $\boldsymbol{R}\Gamma(\partial\widetilde{X}, \widetilde{\imath}^!\mathscr{F}_{\leqslant 0})$ in (4.27), so that (4.28) becomes

$$\begin{aligned}0 \longrightarrow \mathscr{H}^{-1}(i_D^!\mathscr{P}) \longrightarrow \boldsymbol{H}^{-1}(\partial\widetilde{X}, \widetilde{\imath}^!\mathscr{F}_{\leqslant 0}) \longrightarrow \boldsymbol{H}^0_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) \longrightarrow \mathscr{H}^0(i_D^!\mathscr{P})\\ \longrightarrow \boldsymbol{H}^0(\partial\widetilde{X}, \widetilde{\imath}^!\mathscr{F}_{\leqslant 0}) \longrightarrow \boldsymbol{H}^1_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) \longrightarrow \mathscr{H}^1(i_D^!\mathscr{P}) \longrightarrow \boldsymbol{H}^1(\partial\widetilde{X}, \widetilde{\imath}^!\mathscr{F}_{\leqslant 0}) \longrightarrow 0,\end{aligned}$$

since $\boldsymbol{H}^2_{\mathrm{c}}(\widehat{X}, \widehat{\mathscr{F}}) = 0$. Our claim is now that $\mathscr{H}^{-1}(i_D^!\mathscr{P}) = 0$, which follows from $\boldsymbol{H}^{-1}(\partial\widetilde{X}, \widetilde{\imath}^!\mathscr{F}_{\leqslant 0}) = H^{-1}(\partial\widetilde{X}, \mathscr{L}/\mathscr{L}_{\leqslant 0}) = 0$. ☐

# LECTURE 5

# THE RIEMANN-HILBERT CORRESPONDENCE FOR HOLONOMIC $\mathscr{D}$-MODULES ON CURVES

**Summary.** In this lecture, we define the Riemann-Hilbert correspondence on a Riemann surface $X$ as a functor from the category of holonomic $\mathscr{D}_X$-modules to that of Stokes-perverse sheaves. It is induced from a functor at the derived category level which is compatible with $t$-structures. Given a discrete set $D$ in $X$, we first define the correspondence for $\mathscr{D}_X(*D)$-modules which are holonomic and have regular singularities away from $D$, on the one hand, and Stokes-perverse sheaves on $\widetilde{X}(D)$ on the other hand. We then extend the correspondence to $\mathscr{D}_X$-modules of the same kind on the one hand, and Stokes-perverse sheaves on $\underline{\widetilde{X}}(D)$ on the other hand. This lecture follows [**15, 16**], [**46**], [**2**] and [**49**, §IV.3].

## 5.a. Some basic sheaves.

— In this lecture, the base field $\boldsymbol{k}$ is $\mathbb{C}$.

*On* $X$. — Let $X$ be a Riemann surface and $D$ be a discrete set of points in $X$ as in §4.a (from which we keep the notation). We denote by $\mathscr{O}_X$ the sheaf of holomorphic functions on $X$, by $\mathscr{D}_X$ the sheaf of holomorphic differential operators, by $\mathscr{O}_X(*D)$ the sheaf of meromorphic functions on $X$ with poles of arbitrary order on $D$, and we set $\mathscr{D}_X(*D) = \mathscr{O}_X(*D) \otimes_{\mathscr{O}_X} \mathscr{D}_X$.

*On* $\widetilde{X} = \widetilde{X}(D)$. — We set

$$\mathscr{A}_{\widetilde{X}} = \operatorname{Ker}\big[\overline{\partial} : \mathscr{C}^\infty_{\widetilde{X}} \longrightarrow \mathscr{C}^\infty_{\widetilde{X}} \otimes_{\varpi^{-1}\mathscr{O}_{\overline{X}}} \varpi^{-1}\Omega^1_{\overline{X}}(\log \overline{D})\big],$$

where $\overline{X}$ denotes the complex conjugate Riemann surface. This is a $\varpi^{-1}\mathscr{D}_X$-module, which coincide with $\mathscr{O}_{X^*}$ on $X^* := X \smallsetminus D = \widetilde{X} \smallsetminus \varpi^{-1}(D)$. We set $\mathscr{D}_{\widetilde{X}} = \mathscr{A}_{\widetilde{X}} \otimes_{\varpi^{-1}\mathscr{O}_X} \varpi^{-1}\mathscr{D}_X$ and we define similarly $\mathscr{A}_{\widetilde{X}}(*D)$ and $\mathscr{D}_{\widetilde{X}}(*D)$ by tensoring with $\varpi^{-1}\mathscr{O}_X(*D)$.

We have already implicitly defined the sheaf $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}}$ (see Example 1.6) of holomorphic functions on $X^*$ having moderate growth along $\partial\widetilde{X}$. Let us recall the definition in the present setting. Given an open set $\widetilde{U}$ of $\widetilde{X}$, a section $f$ of $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}}$ on $\widetilde{U}$ is a holomorphic function on $U^* := \widetilde{U} \cap X^*$ such that, for any compact set $K$ in $\widetilde{U}$, in the neighbourhood of which $D$ is defined by $g_K \in \mathscr{O}_X(K)$, there exists constants $C_K > 0$ and $N_K \geqslant 0$ such that $|f| \leqslant C_K |g_K|^{-N_K}$ on $K$.

Similarly, $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ consists of functions having rapid decay along $\partial\widetilde{X}$: change the definition above by asking that, for any $K$ and any $N \geqslant 0$, there exists a constant $C_{K,N}$ such that $|f| \leqslant C_{K,N}|g_K|^N$ on $K$.

Both sheaves $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$ and $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ are left $\mathscr{D}_{\widetilde{X}}(*D)$-modules and are $\varpi^{-1}\mathscr{O}_X(*D)$-flat. It will be convenient to introduce the quotient sheaf $\mathscr{A}_{\widetilde{X}}^{\mathrm{gr}\,D} := \mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}/\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ with its natural $\mathscr{D}_{\widetilde{X}}(*D)$-module structure. It is supported on $\partial\widetilde{X}$ and is $\varpi^{-1}\mathscr{O}_X$-flat (because it has no nonzero $\varpi^{-1}\mathscr{O}_X$-torsion).

More information on these sheaves can be found for instance in [**46, 42, 49, 73**]. In particular, $\boldsymbol{R}\varpi_*\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D} = \varpi_*\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D} = \mathscr{O}_X(*D)$, which will be reproved in any dimension in Proposition 8.7.

*On* $\mathcal{I}^{\text{ét}}$. — Recall that $\mathcal{I}$ is the extension by 0 of the sheaf $\mathcal{I}_{\partial\widetilde{X}}$. We will use the definition of Remark 2.23 in order to consider $\mathcal{I}_{\partial\widetilde{X}}$ as a subsheaf of $i_\partial^{-1}j_{\partial,*}\mathscr{O}_{X^*}/i_\partial^{-1}(j_{\partial,*}\mathscr{O}_{X^*})^{\mathrm{lb}}$ on $\partial\widetilde{X}$. We will use the following lemma.

**Lemma 5.1.** — *Let $U$ be an open subset of $\partial\widetilde{X}$ and let $\varphi \in \Gamma(U, \mathcal{I}_{\partial\widetilde{X}})$. There exists a unique subsheaf of $(i_\partial^{-1}j_{\partial,*}\mathscr{O}_{X^*})_{|U}$, denoted by $e^\varphi\mathscr{A}_U^{\mathrm{mod}\,D}$ (resp. by $e^\varphi\mathscr{A}_U^{\mathrm{rd}\,D}$) such that, for any $\theta \in U$ and any lifting $\widetilde{\varphi}_\theta \in \widetilde{\mathcal{I}}_\theta$ of the germ $\varphi_\theta \in \mathcal{I}_\theta$, the germ $(e^\varphi\mathscr{A}_U^{\mathrm{mod}\,D})_\theta$ (resp. $(e^\varphi\mathscr{A}_U^{\mathrm{rd}\,D})_\theta$) is equal to $e^{\widetilde{\varphi}_\theta}\mathscr{A}_\theta^{\mathrm{mod}\,D} \subset (j_{\partial,*}\mathscr{O}_{X^*})_\theta$ (resp. $e^{\widetilde{\varphi}_\theta}\mathscr{A}_\theta^{\mathrm{rd}\,D} \subset (j_{\partial,*}\mathscr{O}_{X^*})_\theta$).*

*Proof.* — It is a matter of checking that $e^{\widetilde{\varphi}_\theta}\mathscr{A}_\theta^{\mathrm{mod}\,D}$ (resp. $e^{\widetilde{\varphi}_\theta}\mathscr{A}_\theta^{\mathrm{rd}\,D}$) does not depend on the choice of the lifting $\widetilde{\varphi}_\theta$. This is clear since any two such choices differ by a locally bounded function. $\square$

We will define sheaves $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$ and $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$ on $\mathcal{I}^{\text{ét}}$ by their restrictions to the closed and open subsets above, together with the gluing morphism.

- On $\mathcal{I}^{\text{ét}}_{X^*} = X^*$, we set $\mathscr{A}_{\mathcal{I}^{\text{ét}}_{X^*}}^{\mathrm{mod}\,D} = \mathscr{A}_{\mathcal{I}^{\text{ét}}_{X^*}}^{\mathrm{rd}\,D} = \mathscr{O}_{X^*}$.
- On $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$, we universally twist the sheaves $\mu_\partial^{-1}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D}$ and $\mu_\partial^{-1}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{rd}\,D}$ by $e^\varphi$ as follows. For any open subset $U \subset \partial\widetilde{X}$ and any $\varphi \in \Gamma(U, \mathcal{I})$, we set $\varphi^{-1}\mathscr{A}_{\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}}^{\mathrm{mod}\,D} := e^\varphi\mathscr{A}_U^{\mathrm{mod}\,D} \subset (i_\partial^{-1}j_{\partial,*}\mathscr{O}_{X^*})_{|U}$ and we define $\varphi^{-1}\mathscr{A}_{\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}}^{\mathrm{rd}\,D}$ similarly (see Lemma 5.1 above; note that on the left-hand side, $\varphi$ is considered as a section of $\mu_\partial : \mathcal{I}^{\text{ét}}_{\partial\widetilde{X}} \to \partial\widetilde{X}$, while on the right-hand side, $\varphi$ and $e^\varphi$ are considered as local sections of $i_\partial^{-1}j_{\partial,*}\mathscr{O}_{X^*}$). One can check that these data correspond to sheaves $\mathscr{A}_{\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}}^{\mathrm{mod}\,D}$ and $\mathscr{A}_{\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}}^{\mathrm{rd}\,D}$, with $\mathscr{A}_{\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}}^{\mathrm{rd}\,D} \subset \mathscr{A}_{\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}}^{\mathrm{mod}\,D} \subset \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{O}_{X^*}$.
- The previous inclusion is used for defining the gluing $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$ and $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D}$ as sheaves on $\mathcal{I}^{\text{ét}}$.

We have natural inclusions $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D} \subset \mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D} \subset \widetilde{\jmath}_* \mathscr{O}_{X^*}$ of sheaves on $\mathcal{I}^{\text{ét}}$. These are inclusions of sheaves of $\mu^{-1}\mathscr{D}_{\widetilde{X}}(*D)$-modules. Indeed, this is clear on $X^*$, and on $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$ one remarks that, in a local coordinate $x$ used for defining $\mathcal{I}$, $\partial_x(e^{\varphi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D}) = e^{\varphi}(\partial_x + \partial\varphi/\partial x)\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D} \subset e^{\varphi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D}$, and similarly for $\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{rd}\,D}$.

It will be convenient to introduce the notation $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{gr}\,D}$ for the quotient sheaf $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D}/\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$. This sheaf is supported on $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$, and is also equipped with a natural $\mu^{-1}\mathscr{D}_{\widetilde{X}}(*D)$-module structure.

These sheaves are flat over $\mu^{-1}\varpi^{-1}\mathscr{O}_X$ (i.e., they have no torsion).

**Lemma 5.2.** — *The sheaves* $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D}$, $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$ *are pre-$\mathcal{I}$-filtrations of* $j_{\partial,*}\mathscr{O}_{X^*}$, *the sheaf* $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{gr}\,D}$ *is a graded pre-$\mathcal{I}$-filtration, and the exact sequence*

$$0 \longrightarrow \mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D} \longrightarrow \mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D} \longrightarrow \mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{gr}\,D} \longrightarrow 0$$

*is an exact sequence in* $\mathrm{Mod}(\mathbb{C}_{\mathcal{I}^{\text{ét}},\leqslant})$. *Moreover, the pre-$\mathcal{I}$-filtration morphisms* $\beta_{\leqslant}p_1^{-1} \to p_2^{-1}$ *are compatible with the* $\mu^{-1}\mathscr{D}_{\widetilde{X}}$*-action.*

*Proof.* — In order to prove the abstract pre-$\mathcal{I}$-filtration property, we need to prove that there are natural morphisms $\beta_{\leqslant}p_1^{-1}\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D} \to p_2^{-1}\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{mod}\,D}$, etc. compatible with the exact sequence of the lemma and the $\mu^{-1}\mathscr{D}_{\widetilde{X}}$-action. This amounts to defining natural morphisms $\beta_{\varphi\leqslant\psi}e^{\varphi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D} \to e^{\psi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D}$, etc. for any open set $U \subset \partial\widetilde{X}$ and any pair $\varphi, \psi \in \Gamma(U, \mathcal{I}_{\partial\widetilde{X}})$. On $U_{\varphi\leqslant\psi}$, we have $e^{\varphi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D} = e^{\psi}e^{\varphi-\psi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D} \subset e^{\psi}\mathscr{A}_{\partial\widetilde{X}}^{\mathrm{mod}\,D}$, and this inclusion defines the desired morphism. Notice that it is compatible with the action of $\mu_\partial^{-1}\mathscr{D}_{\widetilde{X}}$ by definition.

Clearly, the natural inclusions of $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$ and $\mathscr{A}_{\mathcal{I}^{\text{ét}}}^{\mathrm{rd}\,D}$ in $\widetilde{\jmath}_*\mathscr{O}_{X^*} = \mu^{-1}j_{\partial,*}\mathscr{O}_{X^*}$ make these sheaves pre-$\mathcal{I}$-filtrations of $j_{\partial,*}\mathscr{O}_{X^*}$. $\square$

### 5.b. The Riemann-Hilbert correspondence for germs

*The Riemann-Hilbert correspondence for germs of meromorphic connections*

We recall here the fundamental results of Deligne [**15, 16**] and Malgrange [**46**], [**49**, §IV.3]. We work here in a local setting, and we denote by $\mathscr{O}$, $\mathscr{D}$, etc. the germs at $0 \in \mathbb{C}$ of the corresponding sheaves. We denote by $S^1$ the fibre over 0 of the real blow-up of $\mathbb{C}$ at 0, and by $\widetilde{\mathscr{O}}$, $\mathscr{A}^{\mathrm{mod}\,0}$, $\mathscr{A}^{\mathrm{rd}\,0}$, $\mathscr{A}^{\mathrm{gr}_0}$ the restriction to $S^1$ of the sheaves introduced above.

By a meromorphic connection $\mathscr{M}$ we mean a free $\mathscr{O}(*0)$-module of finite rank with a connection. We can form various de Rham complexes, depending on the sheaves of coeffcients.

(1) $\mathrm{DR}\,\mathscr{M} = \{\mathscr{M} \xrightarrow{\nabla} \mathscr{M} \otimes \Omega^1\}$ is the ordinary holomorphic de Rham complex.

(2) $\widetilde{\mathrm{DR}}\mathscr{M} = \{\widetilde{\mathscr{O}} \otimes \varpi^{-1}\mathscr{M} \xrightarrow{\nabla} \widetilde{\mathscr{O}} \otimes \varpi^{-1}(\mathscr{M} \otimes \Omega^1)\}$ is the multivalued essential holomorphic de Rham complex.

(3) $\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}=\{\mathscr{A}^{\mathrm{mod}\,0}\otimes\varpi^{-1}\mathscr{M}\xrightarrow{\nabla}\mathscr{A}^{\mathrm{mod}\,0}\otimes\varpi^{-1}(\mathscr{M}\otimes\Omega^1)\}$ is the moderate de Rham complex (the tensor products are taken over $\varpi^{-1}\mathscr{O}$ and the connection is extended in a natural way, by using the $\varpi^{-1}\mathscr{D}$-structure on $\mathscr{A}^{\mathrm{mod}\,0}$).

(4) $\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}=\{\mathscr{A}^{\mathrm{rd}\,0}\otimes\varpi^{-1}\mathscr{M}\xrightarrow{\nabla}\mathscr{A}^{\mathrm{rd}\,0}\otimes\varpi^{-1}(\mathscr{M}\otimes\Omega^1)\}$ is the rapid-decay de Rham complex.

Since $\boldsymbol{R}\varpi_*\mathscr{A}^{\mathrm{mod}\,0}=\varpi_*\mathscr{A}^{\mathrm{mod}\,0}=\mathscr{O}(*0)$ as already recalled, the projection formula for the proper map $\varpi$ gives

$$\mathrm{DR}\,\mathscr{M}\simeq\boldsymbol{R}\varpi_*\,\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}.$$

On the other hand, $\widetilde{\mathrm{DR}}\mathscr{M}$ has cohomology in degree 0 at most, and this cohomology is the locally constant sheaf with the same monodromy as the monodromy of the local system of horizontal sections of $\mathscr{M}$ away from 0. We will denote by $\mathscr{L}$ the corresponding local system on $S^1$.

***Theorem 5.3.*** — *For any germ $\mathscr{M}$ of meromorphic connection, the complexes $\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}$, $\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}$ and $\mathrm{DR}^{\mathrm{gr}_0}\mathscr{M}$ have cohomology in degree 0 at most. The natural morphisms $\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}\to\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}\to\widetilde{\mathrm{DR}}\mathscr{M}$ induce inclusions $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}\hookrightarrow\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}\hookrightarrow\mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M}$, and $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{gr}_0}\mathscr{M}$ is equal to $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}/\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}$.*

*Sketch of the proof.* — We will indicate that $\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}$ has cohomology in degree 0 at most. The proof for $\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}$ and $\mathrm{DR}^{\mathrm{gr}_0}\mathscr{M}$ is similar, and the remaining part is then straightforward.

(1) One first proves the result in the case where $\mathscr{M}=\mathscr{E}^{\varphi}:=(\mathscr{O}(*0),d+d\varphi)$ for $\varphi\in\mathcal{P}_x$ (see e.g. see e.g. [**83**] or [**49**, App. 1]).

(2) Using that, for any complex number $\alpha$, the local branches of the function $x^{\alpha}$ determine invertible local sections of $\mathscr{A}^{\mathrm{mod}\,0}_{\widetilde{X}}$, (1) implies that the result holds for any $\mathscr{M}$ of the form $\mathscr{E}^{\varphi}\otimes\mathscr{R}$, where $\mathscr{R}$ is a free $\mathscr{O}[1/x]$-module of rank one with the connection $d+\alpha dx/x$.

(3) Using the well-known structure of free $\mathscr{O}[1/x]$-modules (of any rank) with a connection having a regular singularity, it follows from (2) that the result holds for any $\mathscr{M}$ of the form $\mathscr{E}^{\varphi}\otimes\mathscr{R}$, where $\mathscr{R}$ has a connection with regular singularity. We call a direct sum of such modules an *elementary model.*

(4) The theorem holds then for any $\mathscr{M}$ such that, locally on $S^1$, $\mathscr{A}^{\mathrm{mod}\,0}_{\widetilde{X}}\otimes_{\varpi^{-1}\mathscr{O}}\varpi^{-1}\mathscr{M}$ is isomorphic to an elementary model.

(5) By the Hukuhara-Turrittin theorem (see e.g. [**83**] or [**49**, App. 1]), there exists a ramification $\rho:S^1_{x'}\to S^1_x$ such that the pull-back $\rho^+\mathscr{M}$ satisfies the previous property (more precisely, $\rho^+\mathscr{M}$ is locally isomorphic to an elementary model after tensoring with the sheaf $\mathscr{A}^{\mathrm{mod}\,0}_{x'}$), hence the result holds for $\rho^+\mathscr{M}$ for any $\mathscr{M}$ and a suitable $\rho$ (that we can assume to be of the form $x'\mapsto x'^d=x$).

(6) We have $\mathscr{A}_{x'}^{\mathrm{mod}\,0} = \rho^{-1}\mathscr{A}_{x}^{\mathrm{mod}\,0}$. Then

$$\mathscr{A}_{x'}^{\mathrm{mod}\,0} \otimes_{\varpi'^{-1}\mathscr{O}} \varpi'^{-1}\mathscr{M} = \rho^{-1}\big(\mathscr{A}_{x}^{\mathrm{mod}\,0} \otimes_{\varpi^{-1}\mathscr{O}} \varpi^{-1}\mathscr{M}\big),$$

and since $x'\partial_{x'}$ acts like as $d{\cdot}x\partial_x$, we conclude that $\mathrm{DR}^{\mathrm{mod}\,0}(\rho^+\mathscr{M}) \simeq \widetilde{\rho}^{-1}\,\mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{M})$. As a consequence, if the theorem holds for $\rho^+\mathscr{M}$, it holds for $\mathscr{M}$. ☐

***Corollary 5.4***. — *Let $U$ be any open set of $S^1$ and let $\varphi \in \Gamma(U, \mathcal{I})$. Set (see Lemma 5.1)*

$$\begin{aligned}
\mathrm{DR}_{\leqslant\varphi}\mathscr{M} &= \mathrm{DR}(e^{\varphi}\mathscr{A}^{\mathrm{mod}\,0} \otimes \mathscr{M})\\
\mathrm{DR}_{<\varphi}\mathscr{M} &= \mathrm{DR}(e^{\varphi}\mathscr{A}^{\mathrm{rd}\,0} \otimes \mathscr{M})\\
\mathrm{DR}_{\mathrm{gr}_\varphi}\mathscr{M} &= \mathrm{DR}([e^{\varphi}\mathscr{A}^{\mathrm{mod}\,0}/e^{\varphi}\mathscr{A}^{\mathrm{rd}\,0}] \otimes \mathscr{M}).
\end{aligned}$$

*Then these complexes have cohomology in degree $0$ at most. The natural morphisms between these complexes induce inclusions $\mathscr{H}^0\,\mathrm{DR}_{<\varphi}\mathscr{M} \hookrightarrow \mathscr{H}^0\,\mathrm{DR}_{\leqslant\varphi}\mathscr{M} \hookrightarrow \mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M}$, and $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{gr}_\varphi}\mathscr{M}$ is equal to $\mathscr{H}^0\,\mathrm{DR}_{\leqslant\varphi}\mathscr{M}/\mathscr{H}^0\,\mathrm{DR}_{<\varphi}\mathscr{M}$.*

*Proof*. — Assume first that $\varphi$ is not ramified, i.e., comes from an element of $\mathcal{P}_x$. Termwise multiplication by $e^{-\varphi}$ induces an isomorphism of complexes

$$\mathrm{DR}_{\leqslant\varphi}\mathscr{M} \xrightarrow{\sim} \mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^{\varphi} \otimes \mathscr{M})$$

and similarly for $\mathrm{DR}_{<\varphi}$ and $\mathrm{DR}_{\mathrm{gr}_\varphi}$. We can apply the previous result to $\mathscr{E}^{\varphi} \otimes \mathscr{M}$ (where we recall that $\mathscr{E}^{\varphi} = (\mathscr{O}(*0), d + d\varphi)$).

If $\varphi$ is ramified, then we first perform a suitable ramification $\rho$ so that $\varphi$ defines a section of $\mathcal{P}_{x'}$, and we argue as in (6) of the proof of Theorem 5.3 to descend to $\mathscr{M}$. ☐

***Proposition 5.5***. — *Let $\rho$ be such that $\mathscr{M}' = \rho^+\mathscr{M}$ is locally isomorphic on $S^1_{x'}$ to an elementary model after tensoring with $\mathscr{A}_{x'}^{\mathrm{mod}\,0}$. Then the subsheaves $\mathscr{L}'_{\leqslant\varphi'} := \mathscr{H}^0\,\mathrm{DR}_{\leqslant\varphi'}\mathscr{M}'$ considered above ($\varphi' \in \mathcal{P}_{x'}$) form a non-ramified Stokes filtration on $\mathscr{L}' := \mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M}$. Moreover, it is a Stokes filtration on $\mathscr{L}$, for which $\mathscr{L}_{\leqslant\varphi} = \mathscr{H}^0\,\mathrm{DR}_{\leqslant\varphi}\mathscr{M}$, $\mathscr{L}_{<\varphi} = \mathscr{H}^0\,\mathrm{DR}_{<\varphi}\mathscr{M}$, and $\mathrm{gr}_\varphi\mathscr{L} = \mathscr{H}^0\,\mathrm{DR}_{\mathrm{gr}_\varphi}\mathscr{M}$, for any local section $\varphi$ of $\mathcal{I}$.*

*Proof*. — For the first point, one has to check the local gradedness property for $(\mathscr{L}', \mathscr{L}'_\bullet)$. This is a direct consequence of the Hukuhara-Turrittin mentioned in (5) of the proof above. Indeed, this theorem reduces the problem to showing that the pre-Stokes filtration associated to a connection with regular singularity is the graded Stokes filtration with zero as only exponential factor. This reduces to checking when the function $e^{\varphi}|x|^{\alpha}(\log|x|)^k$ has moderate growth, and this reduces to checking when the function $e^{\varphi}$ has moderate growth: we recover exactly the graded Stokes filtration.

For the second point, we have to check that $\mathscr{L}'_{\leqslant\sigma^*\varphi'} = \widetilde{\sigma}^{-1}\mathscr{L}'_{\leqslant\varphi'}$ (see Remark 2.25(1)). This follows from

$$\sigma^+(\mathscr{E}^{\varphi'} \otimes \rho^+\mathscr{M}) = \mathscr{E}^{\sigma^*\varphi'} \otimes \rho^+\mathscr{M}.$$

☐

***Definition 5.6* (The Riemann-Hilbert functor).** — The Riemann-Hilbert functor from the category of germs at 0 of meromorphic connections to the category of Stokes-filtered local systems is the functor which sends an object $\mathscr{M}$ to $(\mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M}, \mathscr{H}^0\,\mathrm{DR}_{\leqslant}\,\mathscr{M})$ and to a morphism the associated morphism at the de Rham level.

***Theorem 5.7* (Deligne [15], Malgrange [46]).** — *The Riemann-Hilbert functor* $\mathscr{M}\mapsto(\mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M}, \mathscr{H}^0\,\mathrm{DR}_{\leqslant}\,\mathscr{M})$ *is an equivalence of categories.*

*Proof of the full faithfulness.* — We first define a morphism

$$\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}\,\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')\longrightarrow\mathscr{H}\!om(\mathscr{L},\mathscr{L}')_{\leqslant 0}.$$

The source of the desired morphism consists of local morphisms $\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}\to\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}'$ which are compatible with the connection. These sections also send $e^{\varphi}\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}$ in $e^{\varphi}\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}'$ for any section $\varphi$ of $\mathfrak{I}$, hence, restricting to the horizontal subsheaves, send $\mathscr{L}_{\leqslant\varphi}$ in $\mathscr{L}'_{\leqslant\varphi}$ for any such $\varphi$. In other words, they define sections of $\mathscr{H}\!om(\mathscr{L},\mathscr{L}')_{\leqslant 0}$.

To show that this morphism is an isomorphism, it is enough to argue locally near $\theta_o\in S^1$ and after a ramification, so that one can use the Hukuhara-Turrittin decomposition for $\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')$ coming from that of $\mathscr{M}$ (indexed by $\Phi$) and of $\mathscr{M}'$ (indexed by $\Phi'$). In fixed local bases of $\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}$ and $\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}'$ adapted to this decomposition, the matrix of a local section of $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}\,\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')$ has blocks $e^{\varphi-\varphi'}u_{\varphi,\varphi'}$, where $u_{\varphi,\varphi'}$ is a local horizontal section of $\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{R}_{\varphi},\mathscr{R}_{\varphi'})$, that is, of $\mathscr{H}\!om(\mathrm{gr}_{\varphi}\,\mathscr{L},\mathrm{gr}_{\varphi'}\,\mathscr{L}')$. The condition that it has moderate growth is equivalent to $\varphi\leqslant_{\theta_o}\varphi'$, that is, is a section of $\mathscr{H}\!om(\mathrm{gr}_{\varphi}\,\mathscr{L},\mathrm{gr}_{\varphi'}\,\mathscr{L}')_{\leqslant 0}$, as wanted.

Let us set $\mathscr{N}=\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')$ with its natural connection. Then we have $\mathrm{Hom}_{\mathscr{D}}(\mathscr{M},\mathscr{M}')=\mathscr{H}^0\,\mathrm{DR}\,\mathscr{N}$. On the one hand, by the projection formula, $\mathrm{DR}\,\mathscr{N}=\boldsymbol{R}\varpi_*\,\mathrm{DR}^{\mathrm{mod}\,0}\,\mathscr{N}$ and by applying Theorem 5.3 to $\mathscr{N}$, this is $\boldsymbol{R}\varpi_*\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}\,\mathscr{N}$. Therefore, $\mathrm{Hom}_{\mathscr{D}}(\mathscr{M},\mathscr{M}')=\varpi_*\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}\,\mathscr{N}$.

As a consequence, $\mathrm{Hom}_{\mathscr{D}}(\mathscr{M},\mathscr{M}')=\Gamma\big(S^1,\mathscr{H}\!om(\mathscr{L},\mathscr{L}')_{\leqslant 0}\big)$, and the latter term is $\mathrm{Hom}\,\big((\mathscr{L},\mathscr{L}_{\bullet}),(\mathscr{L}',\mathscr{L}'_{\bullet})\big)$. □

*Proof of the essential surjectivity.* — We will need:

***Lemma 5.8*.** — *Any graded non-ramified Stokes-filtered local system on* $S^1$ *comes from an elementary model by Riemann-Hilbert.*

*Proof.* — Using Exercise 2.6 and the twist, one is reduced to showing the lemma for the Stokes-graded local system with only one exponential factor, this one being equal to zero. We have indicated above that the germ of connection with regular singularity corresponding to the local system $\mathscr{L}$ gives the corresponding Stokes-filtered local system. □

We will first prove the essential surjectivity when $(\mathscr{L},\mathscr{L}_\bullet)$ is non-ramified. The isomorphism class of $(\mathscr{L},\mathscr{L}_\bullet)$ is obtained from the Stokes-graded local system $\operatorname{gr}_\bullet\mathscr{L}$ by giving a class in $H^1(S^1,\mathscr{A}ut^{<0}(\operatorname{gr}_\bullet\mathscr{L}))$.

Let $\mathscr{M}^{\mathrm{el}}$ be the elementary model corresponding to the Stokes-graded local system $\operatorname{gr}_\bullet\mathscr{L}$ as constructed in Lemma 5.8. Applying the last part of Proposition 5.5 to $\mathscr{E}nd(\mathscr{M}^{\mathrm{el}})$ and the full faithfulness, we find

$$\mathscr{E}nd_{\mathscr{D}}^{\mathrm{rd}\,0}(\mathscr{M}^{\mathrm{el}}):=\mathscr{H}^0\,\mathrm{DR}_{<0}\,\mathscr{E}nd_{\mathscr{O}}(\mathscr{M}^{\mathrm{el}})=\mathscr{H}om(\operatorname{gr}\mathscr{L},\operatorname{gr}\mathscr{L})_{<0},$$

and therefore $\mathscr{A}ut^{\mathrm{rd}\,0}(\mathscr{M}^{\mathrm{el}}):=\mathrm{Id}+\mathscr{E}nd_{\mathscr{D}}^{\mathrm{rd}\,0}(\mathscr{M}^{\mathrm{el}})=\mathscr{A}ut^{<0}(\operatorname{gr}_\bullet\mathscr{L})$.

A cocycle of $\mathscr{A}ut^{<0}(\operatorname{gr}_\bullet\mathscr{L})$ determining $(\mathscr{L},\mathscr{L}_\bullet)$ (with respect to some covering $(I_i)$ of $S^1$) determines therefore a cocycle of $\mathscr{A}ut^{\mathrm{rd}\,0}(\mathscr{M}^{\mathrm{el}})$. If we fix a $\mathscr{O}(*0)$-basis of $\mathscr{M}^{\mathrm{el}}$, the *Malgrange-Sibuya theorem* (see [**46, 49**]) allows one to lift it as a zero cochain $(f_i)$ of $\mathrm{GL}_d(\mathscr{A}^{\mathrm{mod}\,0})$, where $d=\operatorname{rk}\mathscr{M}$ (the statement of the Malgrange-Sibuya theorem is in fact more precise). This zero cochain is used to twist the connection $\nabla^{\mathrm{el}}$ of $\mathscr{M}^{\mathrm{el}}$ on $\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}^{\mathrm{el}}$ as follows: on $I_i$ we set $\nabla_i=f_i^{-1}\nabla^{\mathrm{el}}f_i$. Since $f_if_j^{-1}$ is $\nabla^{\mathrm{el}}$-flat on $I_i\cap I_j$, the connections $\nabla_i$ glue together as a connection $\nabla$ on the free $\mathscr{A}^{\mathrm{mod}\,0}$-module $\mathscr{A}^{\mathrm{mod}\,0}\otimes\mathscr{M}^{\mathrm{el}}$. The matrix of $\nabla$ in the fixed basis of $\mathscr{M}^{\mathrm{el}}$ is a global section of $\mathscr{E}nd((\mathscr{A}^{\mathrm{mod}\,0})^d)$, hence a section of $\mathrm{End}(\mathscr{O}(*0)^d)$. In other words, $\nabla$ defines a new meromorphic connection on the $\mathscr{O}(*0)$-module $\mathscr{M}^{\mathrm{el}}$, that we now denote $\mathscr{M}$. By construction, $(\mathscr{L},\mathscr{L}_\bullet)$ and $(\mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M},\mathscr{H}^0\,\mathrm{DR}_{\leqslant}\mathscr{M})$ are isomorphic, since they correspond to the same cocycle.

Let us now prove the theorem in the ramified case. We have seen (see Remark 2.25(1)) that a Stokes-filtered local system on $\mathscr{L}$ is a Stokes-filtered local system on $\widetilde{\rho}^{-1}\mathscr{L}$ for a suitable $\rho:X'\to X$, which is equivariant with respect to the automorphisms $\widetilde{\sigma}$.

Similarly, let $\mathscr{M}$ be a meromorphic connection and let $\mathscr{M}^*=j_*j^{-1}\mathscr{M}$, where $j$ denotes the inclusion $\mathbb{C}\smallsetminus\{0\}\hookrightarrow\mathbb{C}$ and where we implicitly consider germs at 0. Then giving $\mathscr{M}$ as a subspace of $\mathscr{M}^*$ stable by the connection is equivalent to giving $\mathscr{M}'$ as a subspace of $\rho^*\mathscr{M}^*$ compatible with the connection, and such that $\sigma^*\mathscr{M}'=\mathscr{M}'$ in $\rho^*\mathscr{M}^*$, under the natural identification $\sigma^*\rho^*\mathscr{M}^*=\rho^*\mathscr{M}^*$.

The essential surjectivity in the ramified case follows then from the functoriality (applied to $\sigma^*$) of the Riemann-Hilbert functor. $\square$

Let $\widehat{\mathscr{O}}$ be the ring of formal power series in the variable $x$. If $\mathscr{M}$ is a germ of meromorphic connection, we set $\widehat{\mathscr{M}}=\widehat{\mathscr{O}}\otimes_{\mathscr{O}}\mathscr{M}$. Recall (see e.g. [**49**, Th. (1.5) p. 45]) that $\widehat{\mathscr{M}}$ decomposes as $\widehat{\mathscr{M}}_{\mathrm{reg}}\oplus\widehat{\mathscr{M}}_{\mathrm{irr}}$ where the first summand is regular and the second one is purely irregular. Moreover, $\widehat{\mathscr{M}}_{\mathrm{reg}}$ is the formalization of a unique regular meromorphic connection $\mathscr{M}_{\mathrm{reg}}$ (which is in general not a summand of $\mathscr{M}$).

***Proposition 5.9.*** — *The diagram of functors*

$$\begin{tikzcd}
\mathscr{M} \arrow[r, mapsto] \arrow[d, mapsto, "{(\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,0},\,\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0})}"'] & \mathscr{M}_{\mathrm{reg}} \arrow[d, mapsto, "\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,0}"] \\
(\mathscr{L}_{<0},\mathscr{L}_{\leqslant 0}) \arrow[r, mapsto, "\mathrm{gr}_0"] & \mathscr{G}
\end{tikzcd}$$

*commutes.*

We note that the right vertical functor (to the category of local systems on $S^1$) is an equivalence.

*Sketch of proof.* — We will prove the result in the non-ramified case, the ramified case being treated as above. In such a case, we can easily define in a functorial way a lifting $\mathscr{M}_{\mathrm{irr}}$ of $\widehat{\mathscr{M}_{\mathrm{irr}}}$ by lifting each $\widehat{\mathscr{R}}_\varphi$ occurring in the formal decomposition of $\mathscr{M}$. (One can also define such a lifting in the ramified case, but we will not use it.) The functor $\mathscr{M} \mapsto \mathscr{M}_{\mathrm{reg}}$ factorizes thus through $\mathscr{M} \mapsto \mathscr{M}^{\mathrm{el}} = \mathscr{M}_{\mathrm{irr}} \oplus \mathscr{M}_{\mathrm{reg}} \mapsto \mathscr{M}_{\mathrm{reg}}$. By definition, we have $\mathrm{gr}_0(\mathscr{L}_{<0},\mathscr{L}_{\leqslant 0}) = \mathscr{L}_{\leqslant 0}/\mathscr{L}_{<0} = \mathrm{DR}\big((\mathscr{A}^{\mathrm{mod}\,0}/\mathscr{A}^{\mathrm{rd}\,0}) \otimes_{\varpi^{-1}\mathscr{O}} \varpi^{-1}\mathscr{M}\big)$. The local isomorphisms

$$(\mathscr{A}^{\mathrm{mod}\,0}/\mathscr{A}^{\mathrm{rd}\,0}) \otimes_{\varpi^{-1}\mathscr{O}} \varpi^{-1}\mathscr{M} \xrightarrow{\ \sim\ } (\mathscr{A}^{\mathrm{mod}\,0}/\mathscr{A}^{\mathrm{rd}\,0}) \otimes_{\varpi^{-1}\mathscr{O}} \varpi^{-1}\mathscr{M}^{\mathrm{el}}$$

deduced from the Hukuhara-Turrittin theorem glue together as a global isomorphism. It therefore enough to prove the assertion in the case where $\mathscr{M} = \mathscr{M}^{\mathrm{el}}$, so that $\mathscr{M} = \mathscr{M}_{\mathrm{reg}} \oplus \mathscr{M}_{\mathrm{irr}}$.

If $\varphi \neq 0$, then one checks that the natural morphism $\mathrm{DR}^{\mathrm{rd}\,0}(\mathscr{E}^\varphi \otimes \mathscr{R}) \hookrightarrow \mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^\varphi \otimes \mathscr{R})$ is a quasi-isomorphism (it is enough to check that it induces an isomorphism on $\mathscr{H}^0$, since both $\mathscr{H}^1$ are zero: this is Theorem 5.3). If $\varphi = 0$, then $\mathrm{DR}^{\mathrm{rd}\,0}\mathscr{M}_{\mathrm{reg}} = 0$, and $\mathrm{DR}^{\mathrm{mod}\,0}\mathscr{M}_{\mathrm{reg}}$ is the local system defined by $\mathscr{M}_{\mathrm{reg}}$, hence the assertion. □

*The Riemann-Hilbert correspondence for germs of holonomic $\mathscr{D}$-modules*

We remark (see [**49**, p. 59]) that the category of germs of holonomic $\mathscr{D}$-modules is equivalent to the category of triples $(\mathscr{M}_*, \widehat{\mathscr{M}_{\mathrm{reg}}}, \widehat{\mu})$, where

- $\mathscr{M}_*$ is a locally free $\mathscr{O}(*0)$-module with connection,
- $\widehat{\mathscr{M}_{\mathrm{reg}}}$ is a regular holonomic $\widehat{\mathscr{D}}$-module,
- $\widehat{\mu}$ is a $\widehat{\mathscr{D}}$-linear isomorphism $(\widehat{\mathscr{M}_*})_{\mathrm{reg}} \simeq \widehat{\mathscr{M}_{\mathrm{reg}}}(*0)$.

Indeed, to a holonomic $\mathscr{D}$-module $\mathscr{M}$, we associate $\mathscr{M}_* = \mathscr{O}(*0) \otimes_{\mathscr{O}} \mathscr{M} = \mathscr{M}(*0)$, $\widehat{\mathscr{M}} = \widehat{\mathscr{O}} \otimes_{\mathscr{O}} \mathscr{M}$ which has a unique decomposition $\widehat{\mathscr{M}} = \widehat{\mathscr{M}_{\mathrm{reg}}} \oplus \widehat{\mathscr{M}_{\mathrm{irr}}}$ into regular part and irregular part. Lastly, $\widehat{\mu}$ is uniquely determined by the canonical isomorphism $(\widehat{\mathscr{M}})(*0) \simeq \widehat{\mathscr{M}(*0)}$, by restricting to the regular summand of both terms.

On the other hand, one recovers the holonomic $\mathscr{D}$-module $\mathscr{M}$ from $(\mathscr{M}_*, \widehat{\mathscr{M}_{\mathrm{reg}}}, \widehat{\mu})$ as the kernel of the morphism $\mathscr{M}_* \oplus \widehat{\mathscr{M}_{\mathrm{reg}}} \to \widehat{\mathscr{M}}(*0)$, $m_* \oplus m_r \mapsto \widehat{\mu}(\widehat{m_{*r}}) - \widehat{\mathrm{loc}}(\widehat{m}_r)$, where $\widehat{\mathrm{loc}}$ is the natural localization morphism $\widehat{\mathscr{M}_{\mathrm{reg}}} \to \widehat{\mathscr{M}_{\mathrm{reg}}}(*0)$.

The Riemann-Hilbert functor for germs of holonomic $\mathscr{D}$-modules is now defined as follows: to the holonomic $\mathscr{D}$-module $\mathscr{M}=(\mathscr{M}_*,\widehat{\mathscr{M}}_{\mathrm{reg}},\widehat{\mu})$ is associated the triple $((\mathscr{L},\mathscr{L}_\bullet),\widehat{\mathscr{F}},\nu)$ consisting of a Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$, a germ of perverse sheaf $\widehat{\mathscr{F}}$ and an isomorphism $\widehat{\nu}:\mathrm{gr}_0\mathscr{L}\simeq\psi_x\widehat{\mathscr{F}}$, where $\psi_x\widehat{\mathscr{F}}$ is the local system on $S^1$ defined as in §4.f. The previous results make it clear that this functor is an equivalence.

### 5.c. The Riemann-Hilbert correspondence in the global case.

— By a $\mathscr{D}_X$-module (or $\mathscr{D}_X(*D)$-module), we usually mean a left module. We will say that a coherent $\mathscr{D}_X(*D)$-module is holonomic if it is holonomic as a $\mathscr{D}_X$-module. Recall that, in some neighbourhood of $D$ in $X$, holonomic $\mathscr{D}_X(*D)$-modules are locally free $\mathscr{O}_X(*D)$-modules with connection. On the other hand that any holonomic $\mathscr{D}_X$-module $\mathscr{M}$ gives rise to a holonomic $\mathscr{D}_X(*D)$-module $\mathscr{M}(*D):=\mathscr{D}_X(*D)\otimes_{\mathscr{D}_X}\mathscr{M}$. We will set $\mathscr{M}_{\widetilde{X}}:=\mathscr{A}_{\widetilde{X}}\otimes_{\varpi^{-1}\mathscr{O}_X}\varpi^{-1}\mathscr{M}$ and $\mathscr{M}_{\mathscr{I}_{\text{ét}}}=\mu^{-1}\mathscr{M}_{\widetilde{X}}$.

Let $\mathscr{M}^\bullet$ be an object of $D^b(\mathscr{D}_X)$. The de Rham functor ${}^p\mathrm{DR}:D^b(\mathscr{D}_X)\to D^b(\mathbb{C}_X)$ is defined by ${}^p\mathrm{DR}(\mathscr{M}^\bullet)=\omega_X\otimes^{\boldsymbol{L}}_{\mathscr{D}_X}\mathscr{M}^\bullet$, where $\omega_X$ is $\Omega^1_X$ regarded as a right $\mathscr{D}_X$-module. It is usually written, using the de Rham resolution of $\omega_X$ as a right $\mathscr{D}_X$-module, as $\Omega_X^{1+\bullet}\otimes_{\mathscr{O}_X}\mathscr{M}^\bullet$, with a suitable differential (there is a shift by one with respect to the notation DR used in the previous paragraph). We define similarly functors

- ${}^p\mathrm{DR}^{\mathrm{mod}\,D}_{\widetilde{X}}$, ${}^p\mathrm{DR}^{\mathrm{rd}\,D}_{\widetilde{X}}$ and ${}^p\mathrm{DR}^{\mathrm{gr}\,D}_{\widetilde{X}}$ from $D^b(\mathscr{D}_{\widetilde{X}})$ to $D^b(\mathbb{C}_{\widetilde{X}})$,
- ${}^p\mathrm{DR}^{\mathrm{mod}\,D}_{\mathscr{I}_{\text{ét}}}$, ${}^p\mathrm{DR}^{\mathrm{rd}\,D}_{\mathscr{I}_{\text{ét}}}$ and ${}^p\mathrm{DR}^{\mathrm{gr}\,D}_{\mathscr{I}_{\text{ét}}}$ from $D^b(\mu^{-1}\mathscr{D}_{\widetilde{X}})$ to $D^b(\mathbb{C}_{\mathscr{I}_{\text{ét}},\leqslant})$.

These are obtained by replacing $\omega_X$ with $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}}\otimes_{\varpi^{-1}\mathscr{O}_X}\varpi^{-1}\omega_X$, etc. That ${}^p\mathrm{DR}^{\mathrm{mod}\,D}_{\mathscr{I}_{\text{ét}}}$, etc. take value in $D^b(\mathbb{C}_{\mathscr{I}_{\text{ét}},\leqslant})$ and not only in $D^b(\mathbb{C}_{\mathscr{I}_{\text{ét}}})$ follows from Lemma 5.2.

**Theorem 5.10.** — *If $\mathscr{M}^\bullet$ is an object of $D^b_{\mathrm{hol}}(\mathscr{D}_X(*D))$ (i.e., has holonomic cohomology), then*

- ${}^p\mathrm{DR}^{\mathrm{mod}\,D}_{\mathscr{I}_{\text{ét}}}\mathscr{M}^\bullet$ *is an object of* $D^b_{\text{St-}\mathbb{C}\text{-c}}(\mathbb{C}_{\mathscr{I}_{\text{ét}},\leqslant})$,
- ${}^p\mathrm{DR}^{\mathrm{rd}\,D}_{\mathscr{I}_{\text{ét}}}\mathscr{M}^\bullet$ *is an object of* $D^b_{\text{coSt-}\mathbb{C}\text{-c}}(\mathbb{C}_{\mathscr{I}_{\text{ét}},\leqslant})$,
- ${}^p\mathrm{DR}^{\mathrm{gr}\,D}_{\mathscr{I}_{\text{ét}}}\mathscr{M}^\bullet$ *is an object of* $D^b_{\text{gr-}\mathbb{C}\text{-c}}(\mathbb{C}_{\mathscr{I}_{\text{ét}},\leqslant})$,

*and the functor (called the Riemann-Hilbert functor) of triangulated categories*

$$ {}^p\mathrm{DR}_{\mathscr{I}_{\text{ét}}}:=\{{}^p\mathrm{DR}^{\mathrm{rd}\,D}_{\mathscr{I}_{\text{ét}}}\longrightarrow{}^p\mathrm{DR}^{\mathrm{mod}\,D}_{\mathscr{I}_{\text{ét}}}\longrightarrow{}^p\mathrm{DR}^{\mathrm{gr}\,D}_{\mathscr{I}_{\text{ét}}}\xrightarrow{+1}\} $$

*from $D^b_{\mathrm{hol}}(\mathscr{D}_X(*D))$ to $\mathrm{St}(\mathbb{C}_{\mathscr{I}_{\text{ét}},\leqslant})$ is $t$-exact when $D^b_{\mathrm{hol}}(\mathscr{D}_X(*D))$ is equipped with its natural $t$-structure.*

*Proof.* — One first easily reduces to the case when $\mathscr{M}$ is a holonomic $\mathscr{D}_X(*D)$-module. The question is local, and we can assume that $X$ is a disc with coordinate $x$. We will denote by $\mathscr{O}$, $\mathscr{D}$, etc. the germs at the origin 0 (here equal to $D$) of the corresponding sheaves. The result follows then from Corollary 5.4 and Proposition 5.5. ☐

*The Riemann-Hilbert correspondence for meromorphic connections.* — We restrict here the setting to meromorphic connections with poles on $D$ at most.

***Proposition 5.11.*** — *The functor* $\mathscr{H}^{-1}{}^p\mathrm{DR}_{\mathcal{J}\text{ét}}$ *of Theorem* 5.10 *induces an equivalence between the category of meromorphic connections on* $X$ *with poles on* $D$ *at most and the category* $\text{St-Perv}_{D,\text{smooth}}(\mathbb{C}_{\mathcal{J}\text{ét},\leqslant})$ *of Stokes-perverse sheaves which are locally constant on* $X^*$*.*

Recall (see Remark 4.12) that we also consider such objects as St-$\mathbb{C}$-constructible sheaves shifted by one, so we will mainly work with sheaves and the functor DR instead of perverse sheaves and the functor ${}^p\mathrm{DR}$. The proof being completely parallel to that of Theorem 5.7, we only indicate it.

***Lemma 5.12* (Compatibility with $\mathscr{H}om$).** — *If* $\mathscr{M},\mathscr{M}'$ *are meromorphic connections on* $X$ *with poles on* $D$ *at most, then*

$$\mathscr{H}^0\,\mathrm{DR}_{\mathcal{J}\text{ét}}\,\mathscr{H}om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')\simeq\mathscr{H}om(\mathscr{H}^0\,\mathrm{DR}_{\mathcal{J}\text{ét}}\,\mathscr{M},\mathscr{H}^0\,\mathrm{DR}_{\mathcal{J}\text{ét}}\,\mathscr{M}').$$

*Proof.* — It is similar to the first part of the proof of Theorem 5.7. □

*Proof of Proposition* 5.11*: full faithfulness.* — It is analogous to the corresponding proof in Theorem 5.7. □

*Proof of Proposition* 5.11*: essential surjectivity.* — Because both categories consist of objects which can be reconstructed by gluing local pieces, and because the full faithfulness proved above, it is enough to prove the local version of the essential surjectivity. This is obtained by the similar statement in Theorem 5.7. □

*The Riemann-Hilbert correspondence for holonomic* $\mathscr{D}_X$*-modules.* — For the sake of simplicity, we consider here the Riemann-Hilbert correspondence for holonomic $\mathscr{D}_X$-modules, and not the general case of $D^b_{\text{hol}}(\mathscr{D}_X)$ of bounded complexes with holonomic cohomology. We first define the Riemann-Hilbert functor with values in $\text{St-Perv}_D(\mathbb{C}_{\mathcal{J}\text{ét},\leqslant,\widehat{D}})$ (see §4.f). As for germs, we first remark (see [**49**, p. 59]) that the category of holonomic $\mathscr{D}_X$-modules with singularity set contained in $D$ is equivalent to the category of triples $(\mathscr{M}_*,\mathscr{M}_{\text{reg}},\widehat{\mu})$, where

- $\mathscr{M}_*$ is a locally free $\mathscr{O}_X(*D)$-module with connection,
- $\widehat{\mathscr{M}}_{\text{reg}}$ is a regular holonomic $\mathscr{D}_{\widehat{D}}$-module, where $\mathscr{O}_{\widehat{D}}$ is the formal completion of $\mathscr{O}_X$ at $D$ and $\mathscr{D}_{\widehat{D}}=\mathscr{O}_{\widehat{D}}\otimes_{\mathscr{O}_X}\mathscr{D}_X$,
- $\widehat{\mu}$ is a $\mathscr{D}_{\widehat{D}}$-linear isomorphism $(\mathscr{O}_{\widehat{D}}\otimes_{\mathscr{O}_X}\mathscr{M}_*)_{\text{reg}}\simeq\mathscr{O}_{\widehat{D}}\otimes_{\mathscr{O}_X}\mathscr{M}_{\text{reg}}(*D)$.

Indeed, to a holonomic $\mathscr{D}_X$-module $\mathscr{M}$, we associate $\mathscr{M}_*=\mathscr{O}_X(*D)\otimes_{\mathscr{O}_X}\mathscr{M}=\mathscr{M}(*D)$, $\widehat{\mathscr{M}}=\mathscr{O}_{\widehat{D}}\otimes_{\mathscr{O}_X}\mathscr{M}$ which has a unique decomposition $\widehat{\mathscr{M}}=\widehat{\mathscr{M}}_{\text{reg}}\oplus\widehat{\mathscr{M}}_{\text{irr}}$ into regular part and irregular part. Lastly, $\widehat{\mu}$ is uniquely determined by the canonical isomorphism $(\widehat{\mathscr{M}})(*D)\simeq\widehat{\mathscr{M}(*D)}$, by restricting to the regular summand of both terms.

On the other hand, one recovers $\mathscr{M}$ from data $(\mathscr{M}_*, \widehat{\mathscr{M}}_{\mathrm{reg}}, \widehat{\mu})$ as the kernel of the morphism $\mathscr{M}_* \oplus \widehat{\mathscr{M}}_{\mathrm{reg}} \to \widehat{\mathscr{M}}_{\mathrm{reg}}(*D)$, $m_* \oplus m_r \mapsto \widehat{\mu}((\widehat{m}_*)_r) - \widehat{\mathrm{loc}}(\widehat{m}_r)$, where $\widehat{\mathrm{loc}}$ is the natural localization morphism $\widehat{\mathscr{M}}_{\mathrm{reg}} \to \widehat{\mathscr{M}}_{\mathrm{reg}}(*D)$.

A statement similar to that of Proposition 5.9 holds in this global setting, since it is essentially local at $D$.

The Riemann-Hilbert functor for holonomic $\mathscr{D}$-modules is now defined as follows: to the holonomic $\mathscr{D}_X$-module $\mathscr{M} = (\mathscr{M}_*, \widehat{\mathscr{M}}_{\mathrm{reg}}, \widehat{\mu})$ is associated the triple $({}^p\mathrm{DR}_{\mathcal{J}\text{ét}}\mathscr{M}_*, \widehat{\mathscr{F}}, \widehat{\nu})$, where ${}^p\mathrm{DR}_{\mathcal{J}\text{ét}}\mathscr{M}_*$ is the Stokes-perverse sheaf attached to the meromorphic connection $\mathscr{M}_* := \mathscr{O}_X(*D) \otimes_{\mathscr{O}_X} \mathscr{M}$, $\widehat{\mathscr{F}} = {}^p\mathrm{DR}(\widehat{\mathscr{M}}_{\mathrm{reg}})$, and $\widehat{\nu}$ is the isomorphism defined from $\widehat{\mu}$ and Proposition 5.9. At this point, we have proved that this triple is a Stokes-perverse sheaf on $\underline{\widetilde{X}}$ (see §4.f) and have given the tools for the proof of

***Theorem 5.13.*** — *The Riemann-Hilbert functor is an equivalence between the category of holonomic $\mathscr{D}_X$-modules and the category of Stokes-perverse sheaves on $\underline{\widetilde{X}}$. Under this equivalence, the de Rham functor ${}^p\mathrm{DR}$ corresponds to the "associated perverse sheaf" functor $\mathscr{P}$ (see Proposition 4.25).* □

**5.d. Compatibility with duality for meromorphic connections.** — We will check the compatibility of the Riemann-Hilbert functor of Definition 5.6 with duality, of meromorphic connections on the one hand and of Stokes-filtered local systems on the other hand (see Lemma 2.17). For the sake of simplicity, we do not go further in checking the analogous compatibility at the level of holonomic $\mathscr{D}$-modules and Stokes-perverse sheaves. General results in this directions are obtained in [**62**].

Let $\mathscr{M}$ be a germ of meromorphic connection, as in §5.c and let $\mathscr{M}^\vee$ the dual connection. Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be the Stokes-filtered local system associated with $\mathscr{M}$ by the Riemann-Hilbert functor RH of Definition 5.6. On the other hand, let $(\mathscr{L}, \mathscr{L}_\bullet)^\vee$ be the dual Stokes-filtered local system, with underlying local system $\mathscr{L}^\vee$. We have a natural and functorial identification

$$\mathscr{H}^0\widetilde{\mathrm{DR}}(\mathscr{M}^\vee) = \mathscr{L}^\vee.$$

***Proposition 5.14.*** — *Under this identification, the Stokes filtration $\mathscr{H}^0\,\mathrm{DR}_{\leqslant}(\mathscr{M}^\vee)$ is equal to the dual Stokes filtration $(\mathscr{L}_{\leqslant})^\vee$.*

*Proof.* — The question is local, as it amounts to identifying two Stokes filtration of the same local system $\mathscr{L}^\vee$, hence it is enough to consider the case where $\mathscr{M} = \mathscr{E}^\varphi \otimes \mathscr{R}$ (we will neglect to check what happens with ramification). We will work on a neighbourhood $X$ of the origin, and its real blow-up space $\widetilde{X}$. We denote by $\mathscr{F}$ the local system determined by a representative of $\mathscr{M}$ on $X \smallsetminus \{0\}$. We will also restrict to checking the compatibility for $\mathrm{DR}_{\leqslant 0}, \mathrm{DR}_{<0}$, since the compatibility for $\mathrm{DR}_{\leqslant\psi}, \mathrm{DR}_{<\psi}$ (any $\psi$) directly follows, according to Corollary 5.4 and its proof.

The result is clear if $\varphi = 0$. If $\varphi \neq 0$, we denote by $I_\varphi$ the open set of $S^1$ where $e^{-\varphi}$ has moderate growth, and we denote by $\alpha_\varphi : X^* \hookrightarrow (X^* \cup I_\varphi)$ and $\beta_\varphi : (X^* \cup I_\varphi) \hookrightarrow \widetilde{X}$ the open inclusions. On the one hand, $\mathscr{H}^0 \operatorname{DR}^{\operatorname{mod} 0}\big((\mathscr{E}^\varphi \otimes \mathscr{R})^\vee\big)$ is equal to $\mathscr{L}^\vee$ on the open set $I_{-\varphi}$ and is zero on the complementary closed subset $\overline{I_\varphi}$ of $S^1$. On the other hand, $\mathscr{L}_{<0} = \mathscr{L}_{\leqslant 0}$ is equal to $\mathscr{L}$ on $I_\varphi$ and zero on $\overline{I_{-\varphi}}$. Then, $(\mathscr{L}^\vee)_{\leqslant 0}$, as defined in Proposition 2.15 (or Lemma 2.17), satisfies the same property as $\mathscr{H}^0 \operatorname{DR}^{\operatorname{mod} 0}\big((\mathscr{E}^\varphi \otimes \mathscr{R})^\vee\big)$ does, and both subsheaves of $\mathscr{L}^\vee$ are equal. $\square$

# LECTURE 6

# APPLICATIONS OF THE RIEMANN-HILBERT CORRESPONDENCE TO HOLONOMIC DISTRIBUTIONS

**Summary.** To any holonomic $\mathscr{D}$-module on a Riemann surface $X$ is associated its Hermitian dual, according to M. Kashiwara [**33**] (see also [**8**]). We revisit the proof given in [**73**, §II.3] that the Hermitian dual is also holonomic. As an application, we make explicit the local expression of a holonomic distribution, following [**75**]. The conclusion is that working with $C^\infty$ objects hides the Stokes phenomenon.

**6.a. The Riemann-Hilbert correspondence for meromorphic connections of Hukuhara-Turrittin type.** — Let us go back to the local setting of §5.b. Let $\widetilde{\mathscr{M}}$ be a locally free $\mathscr{A}^{\mathrm{mod}\,0}$-module of rank $d < \infty$ on $S^1$ with a connection $\widetilde{\nabla}$. We say that $\widetilde{\mathscr{M}}$ is *regular* if, locally on $S^1$, it admits a $\mathscr{A}^{\mathrm{mod}\,0}$-basis with respect to which the matrix of the connection takes the form $Cdx/x$, where $C$ is constant.

Let us start by noting that, locally on $S^1$ one can fix a choice of the argument of $x$, so the matrix $x^C$ is well defined as an element of $\mathrm{GL}_d(\mathscr{A}^{\mathrm{mod}\,0})$, and therefore a regular $\mathscr{A}^{\mathrm{mod}\,0}$-connection of rank $d$ is locally isomorphic to $(\mathscr{A}^{\mathrm{mod}\,0})^d$ with its standard connection. As already mentioned in the proof of Theorem 5.3 (see [**49**, App. 1]), the derivation $\partial_x : \mathscr{A}^{\mathrm{mod}\,0} \to \mathscr{A}^{\mathrm{mod}\,0}$ is onto, and it is clear that its kernel is the constant sheaf. As a consequence, for a regular $\mathscr{A}^{\mathrm{mod}\,0}$-connection $\widetilde{\mathscr{M}}$, the sheaf $\operatorname{Ker}\widetilde{\nabla}$ is a locally constant sheaf of rank $d$ on $S^1$, and the natural morphism $\operatorname{Ker}\widetilde{\nabla}\otimes_{\mathbb{C}}\mathscr{A}^{\mathrm{mod}\,0} \to \widetilde{\mathscr{M}}$ is an isomorphism of $\mathscr{A}^{\mathrm{mod}\,0}$-connections.

We conclude that any regular $\mathscr{A}^{\mathrm{mod}\,0}$-connection $\widetilde{\mathscr{M}}$ takes the form $\widetilde{\mathscr{M}} = \mathscr{A}^{\mathrm{mod}\,0}\otimes_{\varpi^{-1}\mathscr{O}}\varpi^{-1}\mathscr{M}$ for some regular $\mathscr{O}(*0)$-connection $\mathscr{M}$, which is nothing but the regular meromorphic connection whose associated local system on the punctured disc is the local system isomorphic to $\operatorname{Ker}\widetilde{\nabla}$. According to the projection formula, we also have $\mathscr{M} = \varpi_*\widetilde{\mathscr{M}}$.

We now extend this result to more general $\mathscr{A}^{\mathrm{mod}\,0}$-connections. As before, the sheaf $\mathfrak{I}$ on $S^1$ is that introduced in §2.c.

***Definition 6.1.*** — Let $\widetilde{\mathscr{M}}$ be a locally free $\mathscr{A}^{\mathrm{mod}\,0}$-module of finite rank on $S^1$. We say that $\widetilde{\mathscr{M}}$ is of *Hukuhara-Turrittin type* if, for any $\theta \in S^1$, there exists a finite set $\Phi_\theta \subset \mathcal{I}_\theta$ such that, in some neighbourhood of $\theta$, $\widetilde{\mathscr{M}}$ is isomorphic to the direct sum indexed by $\varphi \in \Phi_\theta$ of $\mathscr{A}^{\mathrm{mod}\,0}$-connections $(\mathscr{A}^{\mathrm{mod}\,0}, d + d\varphi)^{d_\varphi}$, for some $d_\varphi \in \mathbb{N}^*$.

As indicated above, the regular part of each summand can be reduced to the trivial connection $d$. On the other hand, to any $\mathscr{A}^{\mathrm{mod}\,0}$-connection one can associate de Rham complexes $\mathrm{DR}_{\leqslant\psi}\widetilde{\mathscr{M}}$ and $\mathrm{DR}_{<\psi}\widetilde{\mathscr{M}}$ for any local section $\psi$ of $\mathcal{I}$, by the same formulas as in Corollary 5.4. In particular, $\mathscr{H}^0\,\mathrm{DR}_{\leqslant\psi}\widetilde{\mathscr{M}}$ is a subsheaf of the locally constant sheaf $\mathscr{H}^0\widetilde{\mathrm{DR}}\widetilde{\mathscr{M}}$ (horizontal sections with arbitrary growth along $S^1$) and defines a pre-$\mathcal{I}$-filtration of this sheaf.

***Proposition 6.2.*** — *Assume that $\widetilde{\mathscr{M}}$ is of Hukuhara-Turrittin type. Then $\mathscr{L}_\leqslant := \mathscr{H}^0\,\mathrm{DR}_\leqslant\widetilde{\mathscr{M}}$ is a $\mathcal{I}$-filtration of $\mathscr{L} := \mathscr{H}^0\widetilde{\mathrm{DR}}\widetilde{\mathscr{M}}$, for which $\mathscr{L}_<$ coincides with $\mathscr{H}^0\,\mathrm{DR}_<\widetilde{\mathscr{M}}$. Moreover, the correspondence $\widetilde{\mathscr{M}} \mapsto (\mathscr{L}, \mathscr{L}_\bullet)$ is an equivalence between the category of $\mathscr{A}^{\mathrm{mod}\,0}$-connections of Hukuhara-Turrittin type and the category of Stokes-filtered local systems.*

*Proof.* — We first note that, near $\theta$, we have at most one non-zero morphism (up to a scalar constant) from $\mathscr{A}^{\mathrm{mod}\,0} \otimes \mathscr{E}^\varphi = (\mathscr{A}^{\mathrm{mod}\,0}, d + d\varphi)$ to $\mathscr{A}^{\mathrm{mod}\,0} \otimes \mathscr{E}^\psi = (\mathscr{A}^{\mathrm{mod}\,0}, d + d\psi)$ as $\mathscr{A}^{\mathrm{mod}\,0}$-connections, which is obtained by sending 1 to $e^{\varphi-\psi}$, and this morphism exists if and only if $\varphi \leqslant_\theta \psi$. In particular, both $\mathscr{A}^{\mathrm{mod}\,0}$-connections are isomorphic near $\theta$ if and only if $\varphi = \psi$ near $\theta$, hence everywhere. As a consequence, the set $\Phi_\theta$ does not depend on $\theta$, and we may repeat the proof of Corollary 5.4 for $\widetilde{\mathscr{M}}$. Similarly, the proof of the full faithfulness in Theorem 5.7 can be repeated for $\widetilde{\mathscr{M}}$, since the main argument is local.

The essential surjectivity follows from the similar statement in Theorem 5.7. ☐

***Corollary 6.3.*** — *Let $\widetilde{\mathscr{M}}$ be of Hukuhara-Turrittin type. Then $\mathscr{M} := \varpi_*\widetilde{\mathscr{M}}$ is a $\mathscr{O}(*0)$-connection and $\widetilde{\mathscr{M}} = \mathscr{A}^{\mathrm{mod}\,0} \otimes_{\varpi^{-1}\mathscr{O}} \varpi^{-1}\mathscr{M}$.*

*Proof.* — Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be the Stokes-filtered local system associated with $\widetilde{\mathscr{M}}$ by the previous proposition and let $\mathscr{N}$ be a $\mathscr{O}(*0)$-connection having $(\mathscr{L}, \mathscr{L}_\bullet)$ as associated Stokes-filtered local system, by Theorem 5.7. Let us set $\widetilde{\mathscr{N}} = \mathscr{A}^{\mathrm{mod}\,0} \otimes_{\varpi^{-1}\mathscr{O}} \varpi^{-1}\mathscr{N}$. Since $\mathscr{N}$ is $\mathscr{O}(*0)$-free, we have $\varpi_*\widetilde{\mathscr{N}} = (\varpi_*\mathscr{A}^{\mathrm{mod}\,0}) \otimes_{\mathscr{O}} \mathscr{N} = \mathscr{N}$. The identity morphism $(\mathscr{L}, \mathscr{L}_\bullet) \to (\mathscr{L}, \mathscr{L}_\bullet)$ lifts in a unique way as a morphism $\widetilde{\mathscr{M}} \to \widetilde{\mathscr{N}}$, by the full faithfulness in Proposition 6.2, and the same argument shows that it is an isomorphism. Therefore, $\varpi_*\widetilde{\mathscr{M}} \simeq \mathscr{N}$. ☐

**6.b. The Hermitian dual of a holonomic $\mathscr{D}_X$-module.** — We now assume that $X$ is a Riemann surface, and we denote by $\overline{X}$ the complex conjugate surface (equipped with the structure sheaf $\mathscr{O}_{\overline{X}} := \overline{\mathscr{O}_X}$). The sheaf of distributions $\mathfrak{Db}_X$ on the underlying $C^\infty$ manifold is at the same time a left $\mathscr{D}_X$ and $\mathscr{D}_{\overline{X}}$-module, and both actions commute. If $\mathscr{M}$ is a left $\mathscr{D}_X$-module, then $C_X\mathscr{M} := \mathscr{H}om_{\mathscr{D}_X}(\mathscr{M}, \mathfrak{Db}_X)$ is equipped with a natural structure of left $\mathscr{D}_{\overline{X}}$-module, induced by that on $\mathfrak{Db}_X$. This object, called the *Hermitian dual* of $\mathscr{M}$, has been introduced by M. Kashiwara in [**33**].

***Theorem 6.4*** **([33], [73]).** — *Assume that* $\mathscr{M}$ *is a holonomic* $\mathscr{D}_X$*-module. Then* $\mathscr{E}xt^k_{\mathscr{D}_X}(\mathscr{M}, \mathfrak{Db}_X) = 0$ *for* $k > 0$ *and* $C_X\mathscr{M}$ *is* $\mathscr{D}_{\overline{X}}$*-holonomic.*

We refer to [**33**], [**8**] and [**73**] for various applications of this result. For instance, the vanishing of $\mathscr{E}xt^1$ implies the solvability in $\mathfrak{Db}(\Omega)$ for any linear differential operator with meromorphic coefficients in an open set $\Omega \subset \mathbb{C}$.

*Proof.* — Let us recall various reductions used in [**33**] and [**73**]. If $D$ is the singular divisor of $\mathscr{M}$, it is enough to prove the result for $\mathscr{M}(*D)$, and one can replace $\mathfrak{Db}_X$ with the sheaf $\mathfrak{Db}_X^{\mathrm{mod}\,D}$ of distributions having moderate growth at $D$ (whose sections on an open set consist of continuous linear forms on the space of $C^\infty$ functions with compact support on this subset and having rapid decay at the points of $D$ in this open set). In such a case, $\mathscr{M}(*D)$ is a meromorphic bundle with connection, and the statement is that $C_X^{\mathrm{mod}\,D}(\mathscr{M}(*D)) := \mathscr{H}om_{\mathscr{D}_X(*D)}(\mathscr{M}(*D), \mathfrak{Db}_X^{\mathrm{mod}\,D})$ is an anti-meromorphic bundle with connection, and the corresponding $\mathscr{E}xt^k$ vanish for $k > 0$.

One can then work locally near each point of $D$, and prove a similar result on the real blow-ups space $\widetilde{X}(D)$, by replacing $\mathscr{M}(*D)$ with $\widetilde{\mathscr{M}} = \mathscr{A}^{\mathrm{mod}\,D} \otimes \mathscr{M}(*D)$ and $\mathfrak{Db}_X^{\mathrm{mod}\,D}$ with the similar sheaf $\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\,D}$ on $\widetilde{X}$. One can then reduce to the case where $\widetilde{\mathscr{M}}(*D)$ has no ramification. It is then proved in [**73**, Prop. II.3.2.6] that $\mathscr{E}xt^k_{\mathscr{D}_{\widetilde{X}}}(\widetilde{\mathscr{M}}, \mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\,D}) = 0$ for $k > 0$ and that $C_{\widetilde{X}}^{\mathrm{mod}\,D}(\widetilde{\mathscr{M}})$ is of Hukuhara-Turrittin type (see [**73**, p. 69]). One concludes in loc. cit. by analyzing the Stokes matrices, but this can be avoided by using Corollary 6.3 above: we directly conclude that Prop. II.3.2.6(2) of loc. cit. is fulfilled. $\square$

**6.c. Asymptotic expansions of holonomic distributions.** — We will apply Theorem 6.4 to give the general form of a germ of holonomic distribution of one complex variable. We follow [**75**].

Since the results will be of a local nature, we will denote by $X$ a disc centered at 0 in $\mathbb{C}$, with coordinate $x$. We denote by $\mathfrak{Db}_X^{\mathrm{mod}\,0}$ the sheaf on $X$ of distributions on $X \smallsetminus \{0\}$ with moderate growth at the origin (see above) and by $\mathfrak{Db}^{\mathrm{mod}\,0}$ its germ at the origin. In particular $\mathfrak{Db}^{\mathrm{mod}\,0}$ is a left $\mathscr{D}$ and $\overline{\mathscr{D}}$-module. Let $u \in \mathfrak{Db}^{\mathrm{mod}\,0}$ be holonomic, that is, the $\mathscr{D}$-module $\mathscr{D} \cdot u$ generated by $u$ in $\mathfrak{Db}^{\mathrm{mod}\,0}$ is holonomic. In other words, $u$ is a solution of a non-zero linear differential operator with holomorphic coefficients. Notice that Theorem 6.4 implies that $\overline{\mathscr{D}} \cdot u$ is also holonomic as a $\overline{\mathscr{D}}$-module.

Let $\rho : X' \to X$, $x' \mapsto x'^d = x$, be a ramified covering of degree $d \in \mathbb{N}^*$. Then the pull-back by $\rho$ of a moderate distribution at 0 is well defined as a moderate distribution at 0 on $X'$. If $u$ is holonomic, so is $\rho^*u$ (if $u$ is annihilated by $P \in \mathscr{D}$, then $\rho^*P$ is well defined in $\mathscr{D}_{X',0}(*0)$ and annihilates $\rho^*u$).

***Theorem 6.5.*** — *Let $u$ be a germ at* 0 *of a moderate holonomic distribution on $X$. Then there exists:*

- *an integer $d$, giving rise to a ramified covering $\rho : X' \to X$,*
- *a finite set $\Phi \subset x'^{-1}\mathbb{C}[x'^{-1}]$,*
- *for all $\varphi \in \Phi$, a finite set $B_\varphi \in \mathbb{C}$ and an integer $L_\varphi \in \mathbb{N}$,*
- *for all $\varphi \in \Phi$, $\beta \in B_\varphi$ and $\ell = 0, \dots, L_\varphi$, a function $f_{\varphi,\beta,\ell} \in C^\infty(X')$*

*such that, in $\mathfrak{Db}^{\bmod 0}(U')$ and in particular in $C^\infty(U'^*)$ (if $U'$ is a sufficiently small neighbourhood of* 0 *in $X'$),*

$$\rho^*u = \sum_{\varphi\in\Phi}\sum_{\beta\in B_\varphi}\sum_{\ell=0}^{L_\varphi} f_{\varphi,\beta,\ell}(y)e^{\varphi-\overline{\varphi}}|x'|^{2\beta}\mathrm{L}(x')^{\ell}, \quad \textit{with } \mathrm{L}(x') := \bigl|\log|x'|^2\bigr|. \tag{6.5 $*$}$$

Notice that, for $\varphi \in x'^{-1}\mathbb{C}[x'^{-1}]$, the function $e^{\varphi-\overline{\varphi}}$ is a multiplier in $\mathfrak{Db}^{\bmod 0}$ (since it is $C^\infty$ away from 0, with moderate growth, as well as all its derivatives, at the origin). So are the functions $|x'|^{2\beta}$ and $\mathrm{L}(x')^\ell$.

We will first assume the existence of an expansion like (6.5 $*$) and we will make precise in Corollary 6.11 below the $\varphi, \beta$ such that $f_{\varphi,\beta,\ell} \neq 0$ for some $\ell$. We will allow to restrict the neighbourhood $U$ or $U'$ as needed.

Let $\mathscr{M}$ denote the $\mathscr{D}[x^{-1}]$-module generated by $u$ in $\mathfrak{Db}^{\bmod 0}$. Then $\mathscr{M}$ is a free $\mathscr{O}[x^{-1}]$-module of finite rank equipped with a connection, induced by the action $\partial_x$. Let $\rho : X' \to X$ be a ramification such that $\mathscr{M}' := \rho^+\mathscr{M}$ is formally isomorphic to $\mathscr{M}'^{\mathrm{el}} = \oplus_{\varphi\in\Phi}(\mathscr{E}^\varphi \otimes \mathscr{R}_\varphi)$. The germ $\rho^+\mathscr{M}$ is identified to the $\mathscr{D}'[x'^{-1}]$-submodule of $\mathfrak{Db}^{\bmod 0}_{X',0}$ generated the moderate distribution $v = \rho^*u$. In particular, $v$ is holonomic if $u$ is so.

***Definition 6.6.*** — We say that the holonomic moderate distribution $u$ has *no ramification* if one can choose $\rho = \mathrm{Id}$.

In the following, we will assume that $u$ has no ramification, since the statement of the theorem is given after some ramification. We will however denote by $v$ such a distribution, in order to avoid any confusion. For $\psi \in x^{-1}\mathbb{C}[x^{-1}]$, let $\mathscr{M}_\psi$ be the $\mathscr{D}[x^{-1}]$-module generated by $e^{\overline{\psi}-\psi}v$ in $\mathfrak{Db}^{\bmod 0}$. Note that $e^{\overline{\psi}-\psi}v$ is also holonomic and that this module is $\mathscr{O}[x^{-1}]$-locally free of finite rank with a connection.

We can write a differential equation of minimal degree satisfied by $e^{\overline{\psi}-\psi}v$ in the following way:

$$\Biggl[\prod_{k=0}^{k(1)}\prod_{\beta\in B'_\psi(v)}\bigl[-(x\partial_x-\beta-k)\bigr]^{L'_{\psi,\beta}} - xP_{\psi,1}\Biggr]\cdot e^{\overline{\psi}-\psi}v = 0, \tag{6.7}$$

for some minimal finite set $B'_\psi(v)\subset\mathbb{C}$, $P_{\psi,1}\in\mathbb{C}\{x\}\langle x\partial_x\rangle$, and for each $\beta\in B'_\psi(v)$, $L'_{\psi,\beta}(v)\in\mathbb{N}$ minimal so as to satisfy this equation. Iterating this relation gives a relation of the same kind, for any $j\in\mathbb{N}^*$:

$$\Big[\prod_{k=0}^{k(j)}\prod_{\beta\in B'_\psi(v)}\big[-(x\partial_x-\beta-k)\big]^{L'_{\psi,\beta}}-x^jP_{\psi,j}\Big]\cdot e^{\overline{\psi}-\psi}v=0. \tag{6.8}$$

***Remark 6.9.*** — For almost all $\psi$, we have $B'_\psi(v)=\varnothing$. We denote by $\Phi(v)$ the set of $\varphi$ for which the component $\mathscr{E}^\varphi\otimes\mathscr{R}'_\varphi$ of the formal module associated with $\mathscr{D}[x^{-1}]\cdot v$ is non-zero. It is also the set of $\varphi$ for which the component $\overline{\mathscr{E}^{-\varphi}\otimes\mathscr{R}''_\varphi}$ of the formal module associated with $\overline{\mathscr{D}}[1/\overline{x}]\cdot v$ is non-zero, as a consequence of the proof of Theorem 6.4.

One defines in a conjugate way the objects $L''_{\varphi,\beta}$ and $B''_\varphi(v)$. One then sets

$$B_\varphi(v)=\Big[(B'_\varphi(v)-\mathbb{N})\cap B''_\varphi(v)\Big]\cup\Big[B'_\varphi(v)\cap(B''_\varphi(v)-\mathbb{N})\Big]. \tag{6.10}$$

In other words, $\beta\in B_\varphi(v)$ if and only if $\beta\in B'_\varphi(v)\cup B''_\varphi(v)$ and both $(\beta+\mathbb{N})\cap B'_\varphi(v)$ and $(\beta+\mathbb{N})\cap B''_\varphi(v)$ are non-empty. For all $\beta\in\mathbb{C}$, let us set $L_{\varphi,\beta}(v)=\min\{L'_{\varphi,\beta}(v),L''_{\varphi,\beta}(v)\}$ and $L_{\varphi,\beta+k}(v)=L_{\varphi,\beta}(v)$ for any $k\in\mathbb{Z}$.

Lastly, for $f\in C^\infty(X)$, the Taylor expansion of $f$ at 0 expressed with $x,\overline{x}$ allows us to define a minimal set $E(f)\subset\mathbb{N}^2$ such that $f=\sum_{(\nu',\nu'')\in E(f)}x^{\nu'}\overline{x}^{\nu''}g_{(\nu',\nu'')}$ with $g_{(\nu',\nu'')}\in C^\infty(X)$ and $g_{(\nu',\nu'')}(0)\neq 0$. We will set $E(f)=\varnothing$ if $f$ has rapid decay at 0.

**Corollary 6.11.** — *Let $v$ be a holonomic moderate distribution with no ramification. Then $v$ has an expansion* (6.5 ∗) *in $\mathfrak{Db}^{\mathrm{mod}\,0}$, with $\Phi=\Phi(v)$ and $B_\varphi=B_\varphi(v)$. Moreover, if $f_{\varphi,\beta,\ell}\neq 0$ and if the point $(k',k'')\in\mathbb{N}^2$ is in $E(f_{\varphi,\beta,\ell})$, then $\beta+k'\in B'_\varphi(v)+\mathbb{N}$ and $\beta+k''\in B''_\varphi(v)+\mathbb{N}$.*

*Proof.* — We assume that the theorem is proved. We will use the Mellin transformation to argue on each term of (6.5 ∗). Let $\chi$ be a $C^\infty$ function with compact support contained in an open set where $v$ is defined, and identically equal to 1 near 0. We will also denote by $\chi$ the differential form $\chi\frac{i}{2\pi}\,dx\wedge d\overline{x}$. On the other hand, let us choose a distribution $\widetilde{v}$ inducing $v$ on $X\smallsetminus\{0\}$ and let $p$ be its order on the support of $\chi$. Let us first consider the terms in (6.5 ∗) for which $\varphi=0$.

For all $k',k''\in\mathbb{N}$, the function $s\mapsto\langle\widetilde{v},|x|^{2s}x^{-k'}\overline{x}^{-k''}\chi\rangle$ is defined and holomorphic on the half plane $\{s\mid 2\operatorname{Re}s>p+k'+k''\}$. For all $j\geqslant 1$, let us denote by $Q_j$ the operator such that (6.8) (for $\psi=0$) reads $Q_j\cdot v=0$. Then $Q_j\cdot\widetilde{v}$ is supported at the origin. It will be convenient below to use the notation $\alpha$ for $-\beta-1$ and to set $A'_\varphi(v)=\{\alpha\mid\beta=-\alpha-1\in B'_\varphi(v)\}$. We deduce then that, on some half plane $\operatorname{Re}s\gg 0$, the function

$$\Big[\prod_{k=0}^{k(j)}\prod_{\alpha\in A'_0(v)}(s-\alpha-k'+k)^{L'_{0,\alpha}}\Big]\langle\widetilde{v},|x|^{2s}x^{-k'}\overline{x}^{-k''}\chi\rangle$$

coincides with a holomorphic function defined on $\{s \mid 2\operatorname{Re} s > p + k' + k'' - j\}$. Applying the same argument in a anti-holomorphic way, we obtain that, for each $k', k'' \in \mathbb{N}$, the function $s \mapsto \langle \widetilde{v}, |x|^{2s} x^{-k'} \overline{x}^{-k''} \chi\rangle$ can be extended as a meromorphic function on $\mathbb{C}$ with poles contained in $(A'_0(v) + k' - \mathbb{N}) \cap (A''_0(v) + k'' - \mathbb{N})$, the order of the pole at $\alpha + \mathbb{Z}$ being bounded by $L_{0,\alpha}(v)$. Moreover, this function does not depend on the choice of the lifting $\widetilde{v}$ of $v$.

Let us now compute the Mellin transform of the expansion (6.5 $*$) for $v$.

***Lemma 6.12.*** — *If $\varphi \neq 0$, then for any function $g \in C^\infty(X)$, the Mellin transform of $g(x) e^{\varphi - \overline{\varphi}} |x|^{2\beta} \mathrm{L}(x)^\ell$ is an entire function of the variable $s$.*

*Proof.* — We will show that this Mellin transform is holomorphic on any half plane $\{s \mid \operatorname{Re} s > -q\}$ ($q \in \mathbb{N}$). In order to do so, for $q$ fixed, we decompose $g$ as the sum of a polynomial in $x, \overline{x}$ and a remaining term which vanishes with high order at the origin so that the corresponding part of the Mellin transform is holomorphic for $\operatorname{Re} s > -p$. One is thus reduced to showing the lemma when $g$ is a monomial in $x, \overline{x}$. One can then find differential equations for $g(x) e^{\varphi - \overline{\varphi}} |x|^{2\beta} \mathrm{L}(x)^\ell$ of the kind (6.8) ($j \in \mathbb{N}^*$) with exponents $L'$ equal to zero, and anti-holomorphic analogues. Denoting by $p$ the order of a distribution lifting this moderate function, we find as above that the Mellin transform is holomorphic on a half plane $2\operatorname{Re} s > p + k' + k'' - j$. As this holds for any $j$, the Mellin transform, when $g$ is a monomial, is entire. The Mellin transform for general $g$ is thus holomorphic on any half plane $\{s \mid \operatorname{Re} s > -q\}$, thus is also entire. □

Let us then consider the terms of the expansion (6.5 $*$) for $v$ for which $\varphi = 0$. It is not a restriction to assume (and we will do so) that any two distinct elements of the index set $B_0$ occurring in (6.5 $*$) do not differ by an integer, and that each $\beta \in B_0$ is maximal, meaning that the set $\bigcup_\ell E(f_{0,\beta,\ell})$ is contained in $\mathbb{N}^2$ and in no subset $(m, m) + \mathbb{N}^2$ with $m \in \mathbb{N}^*$. Let $\beta \in B_0$. We will use the property that, for all $(\nu', \nu'') \in \mathbb{Z}^2$ *not both negative* and any function $g \in C^\infty(X)$ such that $g(0) \neq 0$, the poles of the meromorphic function $s \mapsto \langle g(x) |x|^{2\beta} \mathrm{L}(x)^\ell, |x|^{2s} x^{\nu'} \overline{x}^{\nu''} \chi\rangle$ are contained in $\alpha - \mathbb{N}$ (with $\alpha = -\beta - 1$), and there is a pole at $\alpha$ if and only if $\nu' = 0$ and $\nu'' = 0$, this pole having order $\ell + 1$ exactly.

For $\beta \in B_0$, let $E_\beta \subset \mathbb{N}^2$ be minimal such that $E_\beta + \mathbb{N}^2 = \bigcup_\ell (E(f_{0,\beta,\ell}) + \mathbb{N}^2)$. It follows from the previous remark and from the expansion (6.5 $*$) that, for each $(k', k'') \in E_\beta$, the function $s \mapsto \langle \widetilde{v}, |x|^{2s} x^{-k'} \overline{x}^{-k''} \chi\rangle$ has a non trivial pole at $\alpha$; from the first part of the proof we conclude that $\alpha - k' \in A'_0(v) - \mathbb{N}$ and $\alpha - k'' \in A''_0(v) - \mathbb{N}$, that is, $\beta + k' \in B'_0(v) + \mathbb{N}$ and $\beta + k'' \in B''_0(v) + \mathbb{N}$. By assumption on $\beta \in B_0$, there exists $(k', k'') \in E_\beta$ with $k' = 0$ or $k'' = 0$. As a consequence, $\beta \in B_\varphi(v)$ (defined by (6.10)), and the property of the corollary is fulfilled by the elements $(k', k'')$ of $E_\beta$. It is then trivially fulfilled by the elements $(k', k'')$ of all the subsets $E(f_{0,\beta,\ell})$.

The same result holds for the coefficients $f_{\varphi,\beta,\ell}$ by applying the previous argument to the moderate distribution $e^{\overline{\varphi}-\varphi}v$. $\square$

*Proof of Theorem* 6.5. — We go back to the setting of the theorem. Recall that we set $\mathscr{M} = \mathscr{D}(*0)\cdot u \subset \mathfrak{Db}^{\mathrm{mod}\,0}$ and $C^{\mathrm{mod}\,0}\mathscr{M} = \mathscr{H}om_{\mathscr{D}(*0)}(\mathscr{M},\mathfrak{Db}^{\mathrm{mod}\,0})$. There is thus a canonical $\mathscr{D}\otimes_{\mathbb{C}}\overline{\mathscr{D}}$-linear pairing

$$k : \mathscr{M}\otimes_{\mathbb{C}} C^{\mathrm{mod}\,0}\mathscr{M} \longrightarrow \mathfrak{Db}^{\mathrm{mod}\,0}, \qquad (m,\mu)\longmapsto \mu(m).$$

Since $\mathscr{M}$ is generated by $u$ as a $\mathscr{D}(*0)$-module, an element $\mu\in C^{\mathrm{mod}\,0}\mathscr{M}$ is determined by its value $\mu(u)\in\mathfrak{Db}^{\mathrm{mod}\,0}$. Therefore, there exists a section $\mathbf{1}_u$ of $C^{\mathrm{mod}\,0}\mathscr{M}$ such that $\mathbf{1}_u(u)=u$.

We are thus reduced to proving the theorem in the case where

$$k : \mathscr{M}'\otimes_{\mathbb{C}}\overline{\mathscr{M}''}\longrightarrow\mathfrak{Db}^{\mathrm{mod}\,0}$$

is a sesquilinear pairing between two free $\mathscr{O}(*0)$-modules of finite rank with connection, $m',m''$ are local sections of $\mathscr{M}',\mathscr{M}''$, and $u=k(m',\overline{m''})$.

As we allow cyclic coverings, we can also reduce to the case where both $\mathscr{M}'$ and $\mathscr{M}''$ have a formal decomposition with model $\mathscr{M}'^{\mathrm{el}}$ and $\mathscr{M}''^{\mathrm{el}}$ (when $\mathscr{M}''=\overline{C^{\mathrm{mod}\,0}\mathscr{M}'}$, if this assumption is fulfilled for $\mathscr{M}'$, it is also fulfilled for $\mathscr{M}''$, as follows from the proof of Theorem 6.4). We will still denote by $x$ the variable after ramification, and by $X$ the corresponding disc.

We then introduce the real blow-up space $e:\widetilde{X}\to X$ at the origin, together with the sheaves $\mathscr{A}_{\widetilde{X}}$ (see §5.a) and $\mathfrak{Db}^{\mathrm{mod}\,0}_{\widetilde{X}}$ (sheaf on $\widetilde{X}$ of moderate distributions along $e^{-1}(0)=S^1$). Lastly, we set $\widetilde{\mathscr{M}}=\mathscr{A}_{\widetilde{X}}\otimes_{e^{-1}\mathscr{O}}e^{-1}\mathscr{M}$ (for $\mathscr{M}=\mathscr{M}',\mathscr{M}''$). This is a left $\mathscr{D}_{\widetilde{X}}$-module which is $\mathscr{A}_{\widetilde{X}}(*0)$-free.

The pairing $k$ can be uniquely extended as a $\mathscr{D}_{\widetilde{X}}\otimes_{\mathbb{C}}\mathscr{D}_{\overline{\widetilde{X}}}$-linear pairing

$$\widetilde{k} : \widetilde{\mathscr{M}'}\otimes_{\mathbb{C}}\overline{\widetilde{\mathscr{M}''}}\longrightarrow\mathfrak{Db}^{\mathrm{mod}\,0}_{\widetilde{X}}$$

(because $\mathscr{M}',\mathscr{M}''$ are $\mathscr{O}(*0)$-free). One can then work locally on $\widetilde{X}$ with $\widetilde{k}$ and, according to the Hukuhara-Turrittin theorem, we can replace $\widetilde{\mathscr{M}'}$ et $\widetilde{\mathscr{M}''}$ by their respective elementary models $\bigoplus_{\varphi}(\mathscr{E}^{\varphi}\otimes\mathscr{R}'_{\varphi})$ and $\bigoplus_{\varphi}(\mathscr{E}^{\varphi}\otimes\mathscr{R}''_{\varphi})$.

**Lemma 6.13.** — *If $\varphi,\psi\in x^{-1}\mathbb{C}[x^{-1}]$ are distinct, any sesquilinear pairing $\widetilde{k}_{\varphi,\psi}$ : $(\mathscr{E}^{\varphi}\otimes\mathscr{R}'_{\varphi})\otimes_{\mathbb{C}}(\mathscr{E}^{-\overline{\psi}}\otimes\overline{\mathscr{R}''_{-\psi}})\to\mathfrak{Db}^{\mathrm{mod}\,0}_{\widetilde{X}}$ takes values in the subsheaf of $C^{\infty}$ functions with rapid decay.*

*Proof.* — Since $e^{\psi-\overline{\psi}}$ is a multiplier on $\mathfrak{Db}^{\mathrm{mod}\,0}_{\widetilde{X}}$, we can assume that $\psi=0$ for example. By induction on the rank of $\mathscr{R}'_{\varphi}$ and $\mathscr{R}''_0$, we can reduce to the rank-one case, and since the functions $x^{\alpha}$ and $\overline{x}^{\beta}$ are also multipliers, we can reduce to the case where $\mathscr{R}'_{\varphi}$ and $\mathscr{R}''_0$ are both equal to $\mathscr{O}(*0)$. Now, denoting by "$e^{\varphi}$" the generator 1 of $\mathscr{E}^{\varphi}$, the germ of moderate distribution $\widetilde{u}=\widetilde{k}(\text{“}e^{\varphi}\text{”},1)$ on $\widetilde{X}$ satisfies $\overline{\partial}_x\widetilde{u}=0$ and

$\partial_x\widetilde{u}=\varphi'(x)\widetilde{u}$. It follows that $\widetilde{u}_{|X^*}=e^{\varphi}$ with $0\neq\varphi\in x^{-1}\mathbb{C}[x^{-1}]$. Therefore, $\widetilde{u}$ has moderate growth at $\theta\in S^1\Leftrightarrow\widetilde{u}$ has rapid decay at $\theta$. $\square$

In a similar way (using the Jordan normal form for the matrix of the connection on $\mathscr{R}'_0,\mathscr{R}''_0$), one checks that the diagonal terms $\widetilde{k}_{\varphi,\varphi}(\widetilde{m}',\overline{\widetilde{m}''})$ decompose as a sum, with coefficients in $\mathscr{C}^\infty_{\widetilde{X}}$, of terms $e^{\varphi-\overline{\varphi}}x^{\beta'}\overline{x}^{\beta''}(\log x)^j(\log\overline{x})^k$ ($\beta',\beta''\in\mathbb{C}$, $j,k\in\mathbb{N}$). One rewrites each term as a sum, with coefficients in $\mathscr{C}^\infty_{\widetilde{X}}$, of terms $|x|^{2\beta}\mathrm{L}(x)^\ell$ ($\beta\in\mathbb{C}$, $\ell\in\mathbb{N}$).

If $m',m''$ are local sections of $\mathscr{M}',\mathscr{M}''$, one uses a partition of the unity on $\widetilde{X}$ to obtain for $e^*k(m',\overline{m''})$ an expansion like in (6.5 $*$), with coefficients $\widetilde{f}_{\varphi,\beta,\ell}$ in $e_*\mathscr{C}^\infty_{\widetilde{X}}$, up to adding a $C^\infty$ function with rapid decay along $e^{-1}(0)$; such a function can be incorporated in one of the coefficients $\widetilde{f}_{\varphi,\beta,\ell}$. We denote then by $\widetilde{B}_\varphi$ the set of indices $\beta$ corresponding to $\varphi$. Since $|x|$ is $C^\infty$ on $\widetilde{X}$, one can assume that two distinct elements of $\widetilde{B}_\varphi$ do not differ by half an integer.

It remains to check that this expansion can be written with coefficients $f_{\varphi,\beta,\ell}$ in $C^\infty(X)$. We will use the Mellin transformation, as in Corollary 6.11, from which we only take the notation.

We will use polar coordinates $x=re^{i\theta}$. A function $\widetilde{f}\in e_*\mathscr{C}^\infty_{\widetilde{X}}$ has a Taylor expansion $\sum_{m\geqslant0}\widetilde{f}_m(\theta)r^m$, where $\widetilde{f}_m(\theta)$ is $C^\infty$ on $S^1$ and expands as a Fourier series $\sum_n\widetilde{f}_{mn}e^{in\theta}$. Such a function can be written as $\sum_{k=-2k_0}^0 g_k(x)|x|^k$ with $k_0\in\mathbb{N}$ and $g_k\in C^\infty(X)$ if and only if

$$\widetilde{f}_{m,n}\neq0\Longrightarrow\frac{m\pm n}{2}\geqslant-k_0. \tag{6.14}$$

Indeed, if (6.14) is fulfilled, let us set $k=\min(m-n,m+n)$. We have $k\in\mathbb{Z}$ and $k\geqslant-2k_0$. One writes $\widetilde{f}_{m,n}e^{in\theta}r^m=\widetilde{f}_{m,n}|x|^kx^{\ell'}\overline{x}^{\ell''}$ with $\ell',\ell''\in\mathbb{N}$. The part of the Fourier expansion of $\widetilde{f}$ corresponding to $k$ fixed gives, up to multiplying by $|x|^k$, according to Borel's lemma, a function $g_k(x)\in C^\infty(X)$. The difference $\widetilde{f}-\sum_{k=-2k_0}^0 g_k(x)|x|^k$ is $C^\infty$ on $\widetilde{X}$, with rapid decay along $S^1$. It is thus $C^\infty$ on $X$, with rapid decay at the origin. One can add it to $g_0$ in order to obtain the desired decomposition of $\widetilde{f}$. The converse is clear.

The condition (6.14) can be expressed in terms of Mellin transform. Indeed, one notices that, for all $k',k''\in\frac12\mathbb{N}$ such that $k'+k''\in\mathbb{N}$, the Mellin transform $s\mapsto\langle\widetilde{f},|x|^{2s}x^{-k'}\overline{x}^{-k''}\chi\rangle$, which is holomorphic for $\mathrm{Re}(s)\gg0$, extends as a meromorphic function on $\mathbb{C}$ with simple poles at most, and these poles are contained in $\frac12\mathbb{Z}$. The condition (6.14) is equivalent to the existence of $k_0\in\mathbb{N}$ such that, for all $k',k''\in\frac12\mathbb{N}$ with $k'+k''\in\mathbb{N}$, the poles of $s\mapsto\langle\widetilde{f},|x|^{2s}x^{-k'}\overline{x}^{-k''}\chi\rangle$ are contained in the intersection of the sets $k_0-1+k'-\frac12\mathbb{N}^*$ and $k_0-1+k''-\frac12\mathbb{N}^*$.

Arguing by decreasing induction on $\ell$, that is also on the maximal order of the poles, we conclude that a function $\widetilde{f}=\sum_{\ell=0}^L\widetilde{f}_\ell\mathrm{L}(x)^\ell$ with coefficients in $C^\infty(\widetilde{X})$ can be rewritten as $\sum_{-2k_0\leqslant k\leqslant0}\sum_{\ell=0}^L g_{k,\ell}(x)|x|^k\mathrm{L}(x)^\ell$ with $g_{k,\ell}\in C^\infty(X)$ if and only if the same property is fulfilled by $s \mapsto \langle \widetilde{f}, |x|^{2s} x^{-k'} \overline{x}^{-k''} \chi \rangle$ (and the poles have order $\leqslant L+1$).

Now, if $\widetilde{B}_0 \subset \mathbb{C}$ is a finite set such that two distinct elements do not differ by half an integer, a function $\widetilde{f} = \sum_{\beta \in \widetilde{B}_0} \sum_{\ell=0}^{L} \widetilde{f}_{\beta,\ell} |x|^{2\beta} \mathrm{L}(x)^{\ell}$ with coefficients in $C^{\infty}(\widetilde{X})$ can be rewritten $\sum_{\beta \in B_0} \sum_{\ell=0}^{L} f_{\beta,\ell} |x|^{2\beta} \mathrm{L}(x)^{\ell}$ for some subset $B_0$, with $f_{\beta,\ell} \in C^{\infty}(X)$, if and only if there exists a finite subset $A_0 \subset \mathbb{C}$ such that, for all $k', k'' \in \frac{1}{2}\mathbb{N}$ with $k'+k'' \in \mathbb{N}$, the poles of $s \mapsto \langle \widetilde{f}, |x|^{2s} x^{-k'} \overline{x}^{-k''} \chi \rangle$ are contained in $(A_0 + k' - \mathbb{N}) \cap (A_0 + k'' - \mathbb{N})$.

Lastly, if $\widetilde{f}$ has an expansion of the kind (6.5 $*$) with coefficients in $C^{\infty}(\widetilde{X})$, the condition above applied to $\widetilde{f}$ is equivalent to the fact that $\widetilde{f}$ can be rewriten with coefficients $f_{0,\beta,\ell} \in C^{\infty}(X)$, according to an obvious analogue of Lemma 6.12.

We apply this to $k(m', \overline{m''})$: that the condition on the Mellin transform is fulfilled is seen by using equations like (6.7) for $m'$ and $m''$, in the same way as in Corollary 6.11 and this gives the result for the coefficients corresponding to $\varphi = 0$. If $\varphi \neq 0$, one applies the same reasoning to $\mathscr{E}^{-\varphi} \otimes \mathscr{M}'$ et $\mathscr{E}^{\varphi} \otimes \mathscr{M}''$. ☐

**6.d. Comments.** — The notion of Hermitian dual of a $\mathscr{D}$-module was introduced by M. Kashiwara in [**33**], where many applications of the property that the Hermitian dual of a holonomic $\mathscr{D}$-module remains holonomic have been given (see also [**8**]). Kashiwara only treated the case of regular holonomic $\mathscr{D}$-modules (in arbitrary dimension), and examples of such regular holonomic distributions also appear in [**5**].

The vanishing of $\mathscr{E}xt^1$ in Theorem 6.4 is already apparent in [**45**, Th. 10.2], restricting to the real domain however. The analysis of the $\mathscr{H}om$ has been done in [**82**], in particular Th. 3.1 which is a real version of Theorem 6.5, still in the real domain (I thank J.-E. Björk for pointing this reference out to me).

# LECTURE 7

# RIEMANN-HILBERT AND LAPLACE ON THE AFFINE LINE (THE REGULAR CASE)

**Summary.** The Laplace transform of a holonomic $\mathscr{D}$-module $M$ on the affine line $\mathbb{A}^1$ is a holonomic $\mathscr{D}$-module ${}^{\mathrm{F}}M$ on the affine line $\widehat{\mathbb{A}}^1$. If $M$ has only regular singularities (included at infinity), ${}^{\mathrm{F}}M$ provides the simplest example of an irregular singularity (at infinity). We will describe the Stokes-filtered local system attached to ${}^{\mathrm{F}}M$ at infinity in terms of data of $M$. More precisely, we define the topological Laplace transform of the perverse sheaf $\mathrm{DR}^{\mathrm{an}}\,M$ as a perverse sheaf on $\widehat{\mathbb{A}}^1$ equipped with a Stokes structure at infinity. We make explicit this topological Laplace transform. As a consequence, if $\boldsymbol{k}$ is a subfield of $\mathbb{C}$ and if we have a $\boldsymbol{k}$-structure on $\mathrm{DR}^{\mathrm{an}}\,M$, we find a natural $\boldsymbol{k}$-structure on $\mathrm{DR}^{\mathrm{an}}\,{}^{\mathrm{F}}M$ which extends to the Stokes filtration at infinity. In other words, the Stokes matrices can be defined over $\boldsymbol{k}$. We end this lecture by analyzing the behaviour of duality by Laplace and topological Laplace transformation, as well as the relations between them.

**7.a. Introduction.** — Let $M$ be a holonomic $\mathbb{C}[t]\langle\partial_t\rangle$-module, that we regard as an algebraic holonomic $\mathscr{D}_{\mathbb{A}^1}$-module. Its *Laplace transform* ${}^{\mathrm{F}}M$ is a holonomic $\mathbb{C}[\tau]\langle\partial_\tau\rangle$-module, where $\tau$ is a new variable. Recall that ${}^{\mathrm{F}}M$ can be defined in various equivalent ways. Below, we consider the Laplace transform with kernel $e^{t\tau}$, and a similar description can be made for the inverse Laplace transform, which has kernel $e^{-t\tau}$.

(1) The simplest way to define ${}^{\mathrm{F}}M$ is to set ${}^{\mathrm{F}}M = M$ as $\mathbb{C}$-vector spaces and to define the action of $\mathbb{C}[\tau]\langle\partial_\tau\rangle$ in such a way that $\tau$ acts as $-\partial_t$ and $\partial_\tau$ as $t$ (this is modelled on the behaviour of the action of differential operators under Fourier transform of temperate distributions).

(2) One can mimic the Laplace integral formula, replacing the integral by the direct image of $\mathscr{D}$-modules. We consider the diagram

$$\begin{tikzcd} & \mathbb{A}^1 \times \widehat{\mathbb{A}}^1 \arrow[dl, "p"'] \arrow[dr, "\widehat{p}"] & \\ \mathbb{A}^1 & & \widehat{\mathbb{A}}^1 \end{tikzcd}$$

where $t$ is the coordinate on $\mathbb{A}^1$ and $\tau$ that on $\widehat{\mathbb{A}}^1$. Then ${}^{\mathrm{F}}M = \widehat{p}_+(p^+M \otimes E^{t\tau})$, where $E^{t\tau}$ is $\mathbb{C}[t,\tau]$ equipped with the connection $d+d(t\tau)$, and $p^+M$ is $\mathbb{C}[\tau]\otimes_{\mathbb{C}} M$ equipped with its natural connection. Recall also that $\widehat{p}_+$ is the direct image of $\mathscr{D}$-modules, which is defined here in a simple way: $\widehat{p}_+(p^+M\otimes E^{t\tau})$ is the complex

$$0\longrightarrow (p^+M\otimes E^{t\tau})\xrightarrow{\ \partial_t\ }(p^+M\otimes E^{t\tau})\longrightarrow 0$$

where the source of $\partial_t$ is in degree $-1$ and the target in degree $0$.

(3) It will be useful to work with the analytic topology (not the Zariski topology, as above). In order to do so, one has to consider the projective completions $\mathbb{P}^1$ of $\mathbb{A}^1$ (resp. $\widehat{\mathbb{P}}^1$ of $\widehat{\mathbb{A}}^1$) obtained by adding the point $\infty$ to $\mathbb{A}^1$ (resp. the point $\widehat{\infty}$ to $\widehat{\mathbb{A}}^1$). In the following, we shall denote by $t'$ (resp. $\tau'$) the coordinate centered at $\infty$ (resp. $\widehat{\infty}$), so that $t'=1/t$ (resp. $\tau'=1/\tau$) on $\mathbb{A}^1\smallsetminus\{0\}$ (resp. on $\widehat{\mathbb{A}}^1\smallsetminus\{\widehat{0}\}$). We consider the diagram

(7.1)
$$\begin{tikzcd} & \mathbb{P}^1\times\widehat{\mathbb{P}}^1 \arrow[dl, "p"'] \arrow[dr, "\widehat{p}"] & \\ \mathbb{P}^1 & & \widehat{\mathbb{P}}^1 \end{tikzcd}$$

Let $\mathcal{M}$ (resp. ${}^{\mathrm{F}}\mathcal{M}$) be the algebraic $\mathscr{D}_{\mathbb{P}^1}$-module (resp. $\mathscr{D}_{\widehat{\mathbb{P}}^1}$-module) determined by $M$ (resp. ${}^{\mathrm{F}}M$), which is still holonomic, and let $\mathscr{M}$ (resp. ${}^{\mathrm{F}}\mathscr{M}$) be its analytization. We now write $\mathscr{D}_{\mathbb{P}^1}$ (resp. $\mathscr{D}_{\widehat{\mathbb{P}}^1}$) instead of $\mathscr{D}^{\mathrm{an}}_{\mathbb{P}^1}$ (resp. $\mathscr{D}^{\mathrm{an}}_{\widehat{\mathbb{P}}^1}$). Notice that, by definition, $\mathscr{M}=\mathscr{O}_{\mathbb{P}^1}(*\infty)\otimes_{\mathscr{O}_{\mathbb{P}^1}}\mathscr{M}$ and ${}^{\mathrm{F}}\mathscr{M}=\mathscr{O}_{\widehat{\mathbb{P}}^1}(*\widehat{\infty})\otimes_{\mathscr{O}_{\widehat{\mathbb{P}}^1}}{}^{\mathrm{F}}\mathscr{M}$. Applying a similar construction to $E^{t\tau}$ we get $\mathscr{E}^{t\tau}$ on $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$, which is a free $\mathscr{O}_{\mathbb{P}^1\times\widehat{\mathbb{P}}^1}(*(D_\infty\cup D_{\widehat{\infty}}))$-module of rank one, with $D_\infty\cup D_{\widehat{\infty}}=\mathbb{P}^1\times\widehat{\mathbb{P}}^1\smallsetminus(\mathbb{A}^1\times\widehat{\mathbb{A}}^1)$. Then we have

$${}^{\mathrm{F}}\mathscr{M}=\mathscr{H}^0\widehat{p}_+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}).$$

If $M$ is a regular holonomic $\mathbb{C}[t]\langle\partial_t\rangle$-module (i.e., regular at finite distance *and* at infinity), the following is well known (see e.g. [**49**] for the results and the definition of the vanishing cycle functor):

(a) The Laplace transform ${}^{\mathrm{F}}M$ is holonomic, has a regular singularity at the origin $\tau=0$, no other singularity at finite distance, and possibly irregular at infinity.

(b) The formal structure of ${}^{\mathrm{F}}M$ at infinity can be described exactly from the vanishing cycles of $M$ (or of $\mathrm{DR}^{\mathrm{an}}\,M$) at its critical points at finite distance. More precisely, denoting by $\widehat{{}^{\mathrm{F}}\mathscr{M}}$ the formalized connection at $\widehat{\infty}$, we have a decomposition $\widehat{{}^{\mathrm{F}}\mathscr{M}}\simeq\bigoplus_c(\mathscr{R}_c\otimes\mathscr{E}^{c/\tau'})$, where the sum is taken over the singular points $c\in\mathbb{A}^1$ of $M$, and $\mathscr{R}_c$ is a regular formal meromorphic connection corresponding in a one-to-one way to the data of the vanishing cycles of the perverse sheaf ${}^p\mathrm{DR}^{\mathrm{an}}M$ at $c$. As a consequence, the set of exponential factors of the Stokes filtration of ${}^{\mathrm{F}}M$ at $\widehat{\infty}$ consists of these $c/\tau'$, and the Stokes filtration is non-ramified.

The purpose of this lecture is to give an explicit formula for the Stokes filtration of ${}^{\mathrm{F}}M$ at infinity, in terms of topological data obtained from $M$. More precisely, let $\mathscr{F}={}^p\mathrm{DR}^{\mathrm{an}}M$ be the analytic de Rham complex of $M$ (shifted according to the usual perverse convention). The question we address is a formula for the Stokes filtration of ${}^{\mathrm{F}}M$ at $\tau = \infty$ in terms of $\mathscr{F}$ only. In other words, we will define a topological Laplace transform of $\mathscr{F}$ as being a perverse sheaf on $\widehat{\mathbb{A}}^1$ with a Stokes filtration at infinity, in such a way that the topological Laplace transform of $\mathrm{DR}^{\mathrm{an}} M$ is $\mathrm{DR}^{\mathrm{an}}\, {}^{\mathrm{F}}M$ together with its Stokes filtration at infinity (i.e., $\mathrm{DR}^{\mathrm{an}}\, {}^{\mathrm{F}}M$ as a Stokes-perverse sheaf on $\widehat{\mathbb{P}}^1$, see Definition 4.11).

A consequence of this result is that, if the perverse sheaf $\mathscr{F}$ is defined over $\mathbb{Q}$ (say), then the Stokes filtration of ${}^{\mathrm{F}}M$ at infinity is also defined over $\mathbb{Q}$.

### 7.b. Direct image of the moderate de Rham complex.

— Our goal is to obtain the Stokes filtration of ${}^{\mathrm{F}}M$ at infinity as the direct image by $\widehat{p}$ of a sheaf defined over a completion of $\mathbb{A}^1 \times \widehat{\mathbb{A}}^1$ (integral formula).

Let $X := \widetilde{\mathbb{P}}^1$ be the real blow-up of $\mathbb{P}^1$ at $\infty$ (in order to simplify the notation, we do not use the same notation as in Lecture 4). This space is homeomorphic to a closed disc with boundary $S^1_\infty := S^1 \times \{\infty\}$. A similar construction can be done starting from $\widehat{\mathbb{A}}^1$ and its projective completion $\widehat{\mathbb{P}}^1$. We get a space $\widehat{X} := \widetilde{\widehat{\mathbb{P}}^1}$ with boundary $S^1_{\widehat{\infty}}$. We set $X^* = X \smallsetminus \{0\}$ and $\widehat{X}^* := \widehat{X} \smallsetminus \{\widehat{0}\}$.

We thus have a diagram

(7.2)
$$\begin{tikzcd}
 & \mathbb{P}^1 \times \widehat{X} \arrow[d, "\mathrm{Id} \times \widehat{\varpi}"'] \arrow[ddrr, bend left, "\widehat{q}"] & & \\
 & \mathbb{P}^1 \times \widehat{\mathbb{P}}^1 \arrow[dl, "p"'] \arrow[dr, "\widehat{p}"] & & \\
\mathbb{P}^1 & & \widehat{\mathbb{P}}^1 & \widehat{X} \arrow[l, "\widehat{\varpi}"']
\end{tikzcd}$$

We denote by $\mathscr{A}_{\widehat{X}}^{\mathrm{mod}\,\widehat{\infty}}$ the sheaf on $\widehat{X}$ of holomorphic functions on $\widehat{\mathbb{A}}^1$ which have moderate growth along $S^1_{\widehat{\infty}}$.

Similarly we denote by $\mathscr{A}_{\mathbb{P}^1 \times \widehat{X}}^{\mathrm{mod}\,\widehat{\infty}}$ the sheaf on $\mathbb{P}^1 \times \widehat{X}$ of holomorphic functions on $\mathbb{P}^1 \times \widehat{\mathbb{A}}^1$ which have moderate growth along $\mathbb{P}^1 \times S^1_{\widehat{\infty}}$. We have a natural inclusion

$$\widehat{q}^{-1} \mathscr{A}_{\widehat{X}}^{\mathrm{mod}\,\widehat{\infty}} \hookrightarrow \mathscr{A}_{\mathbb{P}^1 \times \widehat{X}}^{\mathrm{mod}\,\widehat{\infty}}. \tag{7.3}$$

We will denote by $\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}$ the de Rham complex of a $\mathscr{D}$-module with coefficients in such rings.

***Lemma 7.4.*** — *There is a functorial morphism*

$$\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}({}^{\mathrm{F}}\mathscr{M}) \longrightarrow \boldsymbol{R}\widehat{q}_* \, \mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+ \mathscr{M} \otimes \mathscr{E}^{t\tau})[1] \tag{$7.4_{\leqslant 0}$}$$

*and this morphism is injective on the zero-th cohomology sheaves.*

*Proof.* — Let us first consider the relative de Rham complex $\mathrm{DR}_{\widehat{q}}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ defined as

$$\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\mathbb{P}^1\times\widehat{X}}\otimes(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\xrightarrow{\ \nabla'\ }\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\mathbb{P}^1\times\widehat{X}}\otimes\big[\Omega^1_{\mathbb{P}^1\times\widehat{X}/\widehat{X}}\otimes(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\big],$$

where $\nabla'$ is the relative part of the connection on $(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$, i.e., which differentiates only in the $\mathbb{P}^1$ direction. Then $\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ is the single complex associated to the double complex

$$\mathrm{DR}_{\widehat{q}}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\xrightarrow{\ \nabla''\ }\Omega^1_{\mathbb{P}^1\times\widehat{\mathbb{P}}^1/\mathbb{P}^1}\otimes\mathrm{DR}_{\widehat{q}}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}),$$

where $\nabla''$ is the connection in the $\widehat{\mathbb{P}}^1$ direction. Hence, $\boldsymbol{R}q_*\,\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ is the single complex associated to the double complex

$$\boldsymbol{R}\widehat{q}_*\,\mathrm{DR}_{\widehat{q}}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\xrightarrow{\ \nabla''\ }\Omega^1_{\widehat{\mathbb{P}}^1}\otimes\boldsymbol{R}\widehat{q}_*\,\mathrm{DR}_{\widehat{q}}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}).$$

The point in using the relative de Rham complex is that it is a complex of $(\widehat{\varpi}\circ\widehat{q})^{-1}\mathscr{O}_{\widehat{\mathbb{P}}^1}$-modules and the differential is linear with respect to this sheaf of rings. On the other hand, one knows that $\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\widehat{X}}$ is flat over $\widehat{\varpi}^{-1}\mathscr{O}_{\widehat{\mathbb{P}}^1}$, because it has no $\widehat{\varpi}^{-1}\mathscr{O}_{\widehat{\mathbb{P}}^1}$-torsion. So we can apply the projection formula to get (in a functorial way)

$$\begin{aligned}\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\widehat{X}}\underset{\widehat{\varpi}^{-1}\mathscr{O}_{\widehat{\mathbb{P}}^1}}{\otimes}&\boldsymbol{R}\widehat{q}_*(\mathrm{Id}\times\widehat{\varpi})^{-1}\,\mathrm{DR}_{\widehat{p}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\\&\simeq\boldsymbol{R}\widehat{q}_*\Big(\widehat{q}^{-1}\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\widehat{X}}\underset{\widehat{q}^{-1}\widehat{\varpi}^{-1}\mathscr{O}_{\widehat{\mathbb{P}}^1}}{\otimes}(\mathrm{Id}\times\widehat{\varpi})^{-1}\,\mathrm{DR}_{\widehat{p}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\Big).\end{aligned}$$

Using the natural morphism (7.3), we find a morphism

$$\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\widehat{X}}\otimes\boldsymbol{R}\widehat{q}_*(\mathrm{Id}\times\widehat{\varpi})^{-1}\,\mathrm{DR}_{\widehat{p}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\longrightarrow\boldsymbol{R}\widehat{q}_*\,\mathrm{DR}_{\widehat{q}}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}),$$

and therefore, taking the double complex with differential $\nabla''$ and the associated single complex, we find

$$\mathscr{A}^{\mathrm{mod}\,\widehat{\infty}}_{\widehat{X}}\otimes\boldsymbol{R}\widehat{q}_*(\mathrm{Id}\times\widehat{\varpi})^{-1}\,\mathrm{DR}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\longrightarrow\boldsymbol{R}\widehat{q}_*\,\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}).\tag{7.5}$$

On the other hand, we know (see e.g. [**50**]) that the (holomorphic) de Rham functor has a good behaviour with respect to the direct image of $\mathscr{D}$-modules, that is, we have a functorial isomorphism

$$\boldsymbol{R}\widehat{p}_*\,\mathrm{DR}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})[1]\simeq\mathrm{DR}\,\big[\widehat{p}_+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\big]$$

where the shift by one comes from the definition of $p_+$ for $\mathscr{D}$-modules. Because $\widehat{p}$ is proper and the commutative diagram in (7.2) is cartesian, the base change $\widehat{\varpi}^{-1}\boldsymbol{R}\widehat{p}_*\simeq(\mathrm{Id}\times\widehat{\varpi})^{-1}\boldsymbol{R}\widehat{q}_*$ shows that the left-hand side of (7.5) is $\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}({}^{\mathrm{F}}\mathscr{M})[-1]$. Shifting (7.5) by one gives the functorial morphism $(7.4_{\leqslant 0})$.

Notice that, over $\widehat{\mathbb{A}}^1$, this morphism is an isomorphism, as it amounts to the compatibility of DR and direct images. On the other hand, when restricted to $S^1_{\widehat{\infty}}$, the left-hand term has cohomology in degree 0 at most (see Theorem 5.3).

Let us show that $\mathscr{H}^0(7.4_{\leqslant 0})$ is injective. It is enough to check this on $S^1_{\widehat\infty}$. If $\widetilde\jmath : \widehat{\mathbb{A}}^1 \hookrightarrow \widehat X$ denotes the inclusion, we have a commutative diagram

$$\begin{tikzcd}
\mathscr{H}^0 \mathrm{DR}^{\mathrm{mod}\,\widehat\infty}({}^{\mathrm{F}}\mathscr{M}) \arrow[r, "\mathscr{H}^0(7.4_{\leqslant 0})"] \arrow[d, hook] & \mathscr{H}^0 \boldsymbol{R}\widehat q_* \mathrm{DR}^{\mathrm{mod}\,\widehat\infty}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})[1] \arrow[d] \\
\widetilde\jmath_*\widetilde\jmath^{-1}\mathscr{H}^0 \mathrm{DR}^{\mathrm{mod}\,\widehat\infty}({}^{\mathrm{F}}\mathscr{M}) \arrow[r, equal] & \widetilde\jmath_*\widetilde\jmath^{-1}\mathscr{H}^0 \boldsymbol{R}\widehat q_* \mathrm{DR}^{\mathrm{mod}\,\widehat\infty}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})[1]
\end{tikzcd}$$

and the injectivity of $\mathscr{H}^0(7.4_{\leqslant 0})$ follows. $\square$

***Theorem 7.6.*** — *The morphism* $(7.4_{\leqslant 0})$ *is an isomorphism.*

This theorem reduces our problem of expressing the Stokes filtration of ${}^{\mathrm{F}}M$ at $\widehat\infty$ in terms of $\mathrm{DR}^{\mathrm{an}}\,M$ to the question of expressing $\mathrm{DR}^{\mathrm{mod}\,\widehat\infty}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ in terms of $\mathrm{DR}\,\mathscr{M}$. Indeed, applying $\boldsymbol{R}\widehat p_*$ would then give us the answer for the $\leqslant 0$ part of the Stokes filtration of ${}^{\mathrm{F}}M$, as recalled in (b) of §7.a. The $\leqslant c/\tau'$ part is obtained by replacing $t$ with $t-c$ and applying the same argument.

By Lemma 7.4, it is enough to prove that the right-hand term of $(7.4_{\leqslant 0})$ has cohomology in degree zero at most, and that $\mathscr{H}^0(7.4_{\leqslant 0})$ is onto on $S^1_\infty$, or equivalently that the germs of both $\mathscr{H}^0$ at any $\theta\in S^1$ have the same dimension.

The proof of the theorem will be done by identifying $\mathrm{DR}^{\mathrm{mod}\,\widehat\infty}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ with a complex constructed from $\mathrm{DR}^{\mathrm{an}}\,M$ and by computing explicitly its direct image. This computation will be topological. This complex obtained from $\mathrm{DR}^{\mathrm{an}}\,M$ will be instrumental for defining the topological Laplace transform. Moreover, the identification will be more easily done on a space obtained by blowing-up $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$, in order to use asymptotic analysis (see also Remark 7.16 below). So, before proving the theorem, we first do the topological computation and define the topological Laplace transform.

***Remark 7.7.*** — If we replace the sheaf $\mathscr{A}_{\widehat X}^{\mathrm{mod}\,\widehat\infty}$ by the sheaf $\mathscr{A}_{\widehat X}^{\mathrm{rd}\,\widehat\infty}$ of functions having rapid decay at infinity, and similarly for $\mathscr{A}_{\mathbb{P}^1\times\widehat X}^{\mathrm{rd}\,\widehat\infty}$, the same proof as for Lemma 7.4 gives a morphism for the corresponding rapid decay de Rham complexes, that we denote by $(7.4)_{<0}$. Then Lemma 7.4 and Theorem 7.6 are valid for the rapid-decay complexes, and the proofs are similar.

**7.c. Topological spaces.** — The behaviour of the function $e^{t\tau}$ near the divisor $D_\infty\cup D_{\widehat\infty}=\mathbb{P}^1\times\widehat{\mathbb{P}}^1\smallsetminus(\mathbb{A}^1\times\widehat{\mathbb{A}}^1)$, where we set $D_\infty:=\{\infty\}\times\widehat{\mathbb{P}}^1$ and $D_{\widehat\infty}:=\mathbb{P}^1\times\{\widehat\infty\}$, is not clearly understood near the two points $(0,\widehat\infty)$ and $(\infty,\widehat 0)$, where it is written in local coordinates as $e^{t/\tau'}$ and $e^{\tau/t'}$ respectively. This leads us to blow up these points.

Let $e: Z\to\mathbb{P}^1\times\widehat{\mathbb{P}}^1$ be the complex blowing-up of $(0,\widehat\infty)$ and $(\infty,\widehat 0)$ in $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$. Above the chart with coordinates $(t,\tau')$ in $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$, we have two charts of $Z$ with respective coordinates denoted by $(t_1,u)$ and $(v',\tau'_1)$, so that $e$ is given respectively by $t\circ e=t_1$, $\tau'\circ e=t_1u$ and $t\circ e=v'\tau'_1$, $\tau'\circ e=\tau'_1$. The exceptional divisor $E$ of $e$ above $(0,\widehat{\infty})$ is defined respectively by $t_1=0$ and $\tau'_1=0$.

Similarly, above the chart with coordinates $(t',\tau)$ in $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$, we have two charts of $Z$ with respective coordinates denoted by $(t'_1,u')$ and $(v,\tau_1)$, so that $e$ is given respectively by $t'\circ e=t'_1$, $\tau\circ e=t'_1u'$ and $t'\circ e=v\tau_1$, $\tau\circ e=\tau_1$. The exceptional divisor $F$ of $e$ above $(\infty,\widehat{0})$ is defined respectively by $t'_1=0$ and $\tau_1=0$.

Away from $E\cup F$ (in $Z$) and $\{(0,\widehat{\infty}),(\infty,\widehat{0})\}$ (in $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$), $e$ is an isomorphism. In particular, we regard the two sets $\mathbb{A}^1\times\widehat{\mathbb{A}}^1$ and $(\mathbb{P}^1\smallsetminus\{0\})\times(\widehat{\mathbb{P}}^1\smallsetminus\{\widehat{0}\})$ as two open subsets of $Z$ whose union is $Z\smallsetminus(E\cup F)=\mathbb{P}^1\times\widehat{\mathbb{P}}^1\smallsetminus\{(0,\widehat{\infty}),(\infty,\widehat{0})\}$.

Moreover, the strict transform of $D_\infty$ in $Z$ is a divisor in $Z$ which does not meet $E$, and meets transversally $F$ and the strict transform of $D_{\widehat{\infty}}$. Similarly, the strict transform of $D_{\widehat{\infty}}$ in $Z$ is a divisor which meets transversally both $E$ and the strict transform of $D_\infty$, and does not meet $F$. We still denote by $D_\infty, D_{\widehat{\infty}}$ these strict transform, and we denote by $D$ the normal crossing divisor $D=e^{-1}(D_\infty\cup D_{\widehat{\infty}})=D_\infty\cup D_{\widehat{\infty}}\cup E\cup F$. This is represented on Figure 1.



FIGURE 1. The natural divisors on $Z$

Let $\widetilde{Z}$ be the real blowing-up of $Z$ along the four components $D_\infty, D_{\widehat{\infty}}, E, F$ of the normal crossing divisor $D$. Then the map $e$ lifts as $\widetilde{e}:\widetilde{Z}\to X\times\widehat{X}$. We have the following commutative diagram of maps:

$$\begin{tikzcd}
\widetilde{Z} \arrow[r, "\widetilde{e}"] \arrow[dr, "\widetilde{\varepsilon}"'] \arrow[dd, "\varpi_Z"'] & X\times\widehat{X} \arrow[d, "\varpi\times\mathrm{Id}"] \\
& \mathbb{P}^1\times\widehat{X} \arrow[d, "\mathrm{Id}\times\widehat{\varpi}"] \\
Z \arrow[r, "e"] & \mathbb{P}^1\times\widehat{\mathbb{P}}^1
\end{tikzcd} \tag{7.8}$$

The set $\widetilde{E}:=\varpi_Z^{-1}(E)$ can be described as follows: over the chart $\mathbb{A}^1_{v'}$ of $E$ with coordinate $v'$, it is equal to $\mathbb{A}^1_{v'}\times S^1$, and over $u=0$ it is equal to $S^1\times S^1$, so it can be identified with $\widetilde{\mathbb{P}}^1_{v'}\times S^1$, but the gluing above $u=0$ with the rest of $\widetilde{Z}$ is done by the diffeomorphism $(\arg v', \arg\tau_1') \mapsto (\arg u = -\arg v', \arg t_1 = \arg v' + \arg\tau_1')$. Similarly, the set $\widetilde{F} := \varpi_Z^{-1}(F)$ can be described as follows: over the chart $\mathbb{A}^1_{u'}$ of $F$ with coordinate $u'$, it is equal to $\mathbb{A}^1_{u'} \times S^1$, and over $v = 0$ it is equal to $S^1 \times S^1$, so it can be identified with $\widetilde{\mathbb{P}}^1_{u'} \times S^1$, but the gluing above $v = 0$ with the rest of $\widetilde{Z}$ is done by the diffeomorphism $(\arg u', \arg t_1') \mapsto (\arg v = -\arg u', \arg\tau_1 = \arg u' + \arg t_1')$. The complement $\widetilde{Z} \smallsetminus (\widetilde{E} \cup \widetilde{F})$ is identified with $X \times \widehat{X} \smallsetminus [(\{0\} \times S^1_{\widehat{\infty}}) \cup (S^1_\infty \times \{\widehat{0}\})]$.

We define two nested closed subsets $L''_{\leqslant 0} \subset L''_{<0}$ of $\widetilde{Z}$ as the subset of points in the neighbourhood of which $e^{-t\tau}$ does not have moderate growth (resp. is not exponentially decreasing). More specifically:

(1) $L''_{\leqslant 0} \subset \varpi_Z^{-1}(D_\infty \cup D_{\widehat{\infty}})$,
(2) over the chart $(t_1, u)$, $L''_{\leqslant 0} = \{\arg u \in [\pi/2, 3\pi/2] \bmod 2\pi\} \cap \{u = 0\}$,
(3) over the chart $(t', \tau')$, $L''_{\leqslant 0} = \{\arg t' + \arg\tau' \in [\pi/2, 3\pi/2] \bmod 2\pi\}$,
(4) over the chart $(v, \tau_1)$, $L''_{\leqslant 0} = \{\arg v \in [\pi/2, 3\pi/2] \bmod 2\pi\} \cap \{v = 0\}$.
(5) We have $L''_{<0} = L''_{\leqslant 0} \cup \widetilde{E} \cup \widetilde{F}$.

We denote by $L'_{\leqslant 0}$ (resp. $L'_{<0}$) the complementary open set of $L''_{\leqslant 0}$ (resp. $L''_{<0}$) in $\widetilde{Z}$. So $L'_{\leqslant 0} \supset L'_{<0} \supset \mathbb{A}^1 \times \widehat{\mathbb{A}}^1$ and, away from $u = 0$ (resp. away from $v = 0$), $L'_{\leqslant 0} \cap \widetilde{E}$ (resp. $L'_{\leqslant 0} \cap \widetilde{F}$) coincides with $\widetilde{E}$ (resp. $\widetilde{F}$). Moreover, $L'_{<0} \cap \widetilde{E} = \varnothing$ and $L'_{<0} \cap \widetilde{F} = \varnothing$.

We have a diagram

$$\begin{tikzcd}
\mathbb{A}^1 \times \widehat{\mathbb{A}}^1 \arrow[r, hook, "\alpha"] \arrow[d, "p"'] & L'_{<0} \arrow[r, hook, "\gamma"] & L'_{\leqslant 0} \arrow[r, hook, "\beta"] & \widetilde{Z} \arrow[d, "\widetilde{\widehat{q}}"] \\
\mathbb{A}^1 & & & \widehat{X}
\end{tikzcd} \tag{7.9}$$

**7.d. Topological Laplace transform.** — Let $\mathscr{F}$ be a perverse sheaf on $\mathbb{A}^1$. We will usually denote by $\mathscr{G}$ the perverse sheaf $p^{-1}\mathscr{F}[1]$ on $\mathbb{A}^1 \times \widehat{\mathbb{A}}^1$.

***Definition 7.10.*** — We set:

- for any perverse sheaf $\mathscr{G}$ on $\mathbb{A}^1 \times \widehat{\mathbb{A}}^1$, $({}^{\mathcal{F}}\mathscr{G})_{\leqslant 0} = \beta_! \boldsymbol{R}(\gamma \circ \alpha)_* \mathscr{G}$ and $({}^{\mathcal{F}}\mathscr{G})^{\ll 0} = (\beta \circ \gamma)_! \boldsymbol{R}\alpha_* \mathscr{G}$,
- by using $\mathscr{G} = p^{-1}\mathscr{F}[1]$, $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0} = \boldsymbol{R}\widetilde{\widehat{q}}_* ({}^{\mathcal{F}}\mathscr{G})_{\leqslant 0}$ and $({}^{\mathrm{F}}\mathscr{F})^{<0} = \boldsymbol{R}\widetilde{\widehat{q}}_* ({}^{\mathcal{F}}\mathscr{G})^{\ll 0}$.

***Proposition 7.11.*** — *When restricted to $\widehat{\mathbb{A}}^1$, $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0|\widehat{\mathbb{A}}^1} = ({}^{\mathrm{F}}\mathscr{F})^{<0}_{|\widehat{\mathbb{A}}^1}$ is a perverse sheaf with singularity at $\widehat{0}$ at most, and $i_{\widehat{0}}^{-1}({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0} = \boldsymbol{R}\Gamma_{\mathrm{c}}(\mathbb{A}^1, \mathscr{F})[1]$. On the other hand, when restricted to $\widehat{X}^*$, $({}^{\mathrm{F}}\mathscr{F})^{<0}[-1]$ and $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}[-1]$ are two nested sheaves, and $\mathrm{gr}_0\, {}^{\mathrm{F}}\mathscr{F}[-1] := ({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}[-1] / ({}^{\mathrm{F}}\mathscr{F})^{<0}[-1]$ is a local system on $S^1_{\widehat{\infty}}$ isomorphic to $({}^p\phi_t\mathscr{F}, T)$ when considered as a vector space with monodromy.*

As usual, $\Gamma_{\mathrm{c}}$ denotes the sections with compact support and $\boldsymbol{R}\Gamma_{\mathrm{c}}$ denotes its derived functor, whose associated cohomology is the cohomology with compact support.

Given a perverse sheaf $\mathscr{F}$ on $\mathbb{A}^1$, the nearby cycle complex $({}^p\psi_t\mathscr{F}, T)$ and the vanishing cycle complex $({}^p\phi_t\mathscr{F}, T)$ (equipped with their monodromy) have cohomology in degree zero at most, and there is a canonical morphism *can* : $({}^p\psi_t\mathscr{F}, T) \to ({}^p\phi_t\mathscr{F}, T)$ whose cone represents the inverse image $i_0^{-1}\mathscr{F}[-1]$ (see e.g. [**49**]).

*Proof.* — Let $C \subset \mathbb{A}^1$ denote the set of singular points of $\mathscr{F}$, and let us still denote by $C \times \widehat{\mathbb{P}}^1 \subset Z$ the strict transform by $e$ of $C \times \widehat{\mathbb{P}}^1 \subset \mathbb{P}^1 \times \widehat{\mathbb{P}}^1$. Since $\mathscr{F}[-1]$ is a local system away from $C$, one checks that $\boldsymbol{R}\alpha_* p^{-1}\mathscr{F}[-1]$ (resp. $\boldsymbol{R}(\gamma \circ \alpha)_* p^{-1}\mathscr{F}[-1]$) is a local system on $L'_{<0} \smallsetminus \varpi_Z^{-1}(C \times \widehat{\mathbb{P}}^1)$ (resp. on $L'_{\leqslant 0} \smallsetminus \varpi_Z^{-1}(C \times \widehat{\mathbb{P}}^1)$). This will justify various restrictions to fibres that we will perform below.

For the first assertion, let us work over $\widetilde{V} := X \times \widehat{\mathbb{A}}^1$, and still denote by $\widetilde{e} : \widetilde{W} \to \widetilde{V}$ the restriction of $\widetilde{e}$ over this open set. Over $\widehat{\mathbb{A}}^{1*}$, $\widetilde{W}$ coincides with $X \times \widehat{\mathbb{A}}^{1*}$. Then it is known (see e.g. [**49**] or [**78**, §1b]) that $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0|\widehat{\mathbb{A}}^{1*}}$ is a smooth perverse sheaf (i.e., a local system shifted by one) with germ at $\tau_o \neq 0$ equal to $\boldsymbol{H}^0_{\Phi_{\tau_o}}(\mathbb{A}^1, \mathscr{F})[1]$, where $\Phi_{\tau_o}$ is the family of closed sets in $\mathbb{A}^1$ whose closure in $\widetilde{\mathbb{P}}^1$ does not cut $L''_{\leqslant 0} \cap (S^1_\infty \times \{\tau_o\})$. Moreover, $L''_{\leqslant 0}$ and $L''_{<0}$ coincide above $\widehat{\mathbb{A}}^{1*}$, so $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0|\widehat{\mathbb{A}}^{1*}} = ({}^{\mathrm{F}}\mathscr{F})_{<0|\widehat{\mathbb{A}}^{1*}}$.

On the other hand, we claim that

$$\boldsymbol{R}\widetilde{e}_*({}^{\mathfrak{F}}\mathscr{G})_{\leqslant 0|S^1_\infty \times \{\widehat{0}\}} = \boldsymbol{R}\widetilde{e}_*({}^{\mathfrak{F}}\mathscr{G})^{\ll 0}{}_{|S^1_\infty \times \{\widehat{0}\}} = 0.$$

This will show that

$$\boldsymbol{R}\widetilde{e}_*({}^{\mathfrak{F}}\mathscr{G})_{\leqslant 0|X \times \{\widehat{0}\}} = \boldsymbol{R}\widetilde{e}_*({}^{\mathfrak{F}}\mathscr{G})^{\ll 0}{}_{|X \times \{\widehat{0}\}} \simeq \widetilde{\jmath}_!\mathscr{F}[1],$$

where $\widetilde{\jmath} : \mathbb{A}^1 \hookrightarrow X$ denotes the inclusion, concluding the proof of the first assertion by taking $\boldsymbol{R}\Gamma$. The canonical map $i_{\widehat{0}}^{-1}({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}[-1] \to {}^p\psi_\tau({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}$ is induced by the natural map $\boldsymbol{R}\Gamma_{\mathrm{c}}(\mathbb{A}^1, \mathscr{F}) \to \boldsymbol{R}\Gamma_{\Phi_{\tau_o}}(\mathbb{A}^1, \mathscr{F}) = \boldsymbol{H}^0_{\Phi_{\tau_o}}(\mathbb{A}^1, \mathscr{F})$. The cone of this map is the complex $\boldsymbol{R}\Gamma_{\mathrm{c}}\big(L'_{\leqslant 0} \cap (S^1_\infty \times \{\tau_o\}), \mathscr{L}[1]\big)$, where $\mathscr{L}$ is the local system defined by $\mathscr{F}$ on $S^1_\infty$. In particular, it has cohomology in degree 0 at most. As a consequence, ${}^p\phi_\tau({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}$ is a vector space in degree 0, and it follows that $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}$ is perverse in the neighbourhood of $\widehat{0}$.

***Remark 7.12.*** — Developing the proof more carefully at this point, one would obtain an identification of ${}^p\phi_\tau({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}$ with ${}^p\psi_{t'}\mathscr{F}$, compatible with monodromies when suitably oriented. The second part of the lemma, that we consider now, is an analogous statement, where the roles of $\mathbb{A}^1$ and $\widehat{\mathbb{A}}^1$ are exchanged.

Let us now prove the claim. On the one hand, $({}^{\mathfrak{F}}\mathscr{G})^{\ll 0}$ is zero on $L''_{<0}$ by definition, hence on $\widetilde{F}$, so its restriction to $\widetilde{e}^{-1}(S^1_\infty \times \{\widehat{0}\})$ is zero.

On the other hand, $\widetilde{e}^{-1}(\theta', \widehat{0})$ is homeomorphic to the real blow-up of $F$ at $v = 0$ (topologically a closed disc), and $L'_{\leqslant 0} \cap \widetilde{e}^{-1}(\theta', \widehat{0})$ is homeomorphic to a closed disc with a closed interval deleted on its boundary. On this set, $\boldsymbol{R}(\gamma \circ \alpha)_*\mathscr{G}[-2]$ is a local system (which is thus constant). Since the cohomology with compact support (due to $\beta_!$) of such a closed disc with a closed interval deleted on its boundary is identically zero (a particular case of Lemme 7.13 below), we get the vanishing of $\boldsymbol{R}\widetilde{e}_*({}^{\mathscr{F}}\mathscr{G})_{\leqslant 0,(\theta',\widehat{0})}$ for any $\theta' \in S^1_\infty$, as claimed.

We now prove the second part of the proposition. Let us compute the fibre of $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}$ and $({}^{\mathrm{F}}\mathscr{F})^{<0}$ at $(|\tau'_o| = 0, \arg\tau'_o = \widehat{\theta}'_o)$. Notice that, above this point that we also denote by $\widehat{\theta}'_o$, we have

$$L'_{<0,\widehat{\theta}'_o} = X^* \smallsetminus \{\arg t \in [\widehat{\theta}'_o + \pi/2, \widehat{\theta}'_o + 3\pi/2] \bmod 2\pi\},$$

and this set is equal to $\widetilde{\mathbb{P}}^1$ minus the closure of the half-plane $\mathrm{Re}(te^{-i\widehat{\theta}'_o}) \leqslant 0$. Then $({}^{\mathrm{F}}\mathscr{F})^{<0}_{\widehat{\theta}'_o} = \boldsymbol{R}\Gamma_{\mathrm{c}}(L'_{<0,\widehat{\theta}'_o}, \boldsymbol{R}\alpha_*\mathscr{F})[1]$.

***Lemma 7.13.*** — *Let $\mathscr{P}$ be a perverse sheaf with finite singularity set on an open disc $\Delta \subset \mathbb{C}$ and let $\Delta'$ be the open subset of the closed disc $\overline{\Delta}$ obtained by deleting from $\partial\overline{\Delta}$ a nonempty closed interval. Let $\alpha : \Delta \hookrightarrow \Delta'$ denote the open inclusion. Then $\boldsymbol{H}^k_{\mathrm{c}}(\Delta', \boldsymbol{R}\alpha_*\mathscr{P}) = 0$ if $k \neq 0$ and $\dim \boldsymbol{H}^0_{\mathrm{c}}(\Delta', \boldsymbol{R}\alpha_*\mathscr{P})$ is the sum of dimensions of the vanishing cycles of $\mathscr{P}$ at points of $\Delta$ (i.e., at the singular points of $\mathscr{P}$ in $\Delta$).*

*Proof.* — This can be proved as follows: one reduces to the case of a sheaf supported on some point (trivial), and to the case of $\mathscr{P} = j_*\mathscr{L}[1]$, where $j$ is the inclusion $\Delta \smallsetminus \mathrm{Sing}(\mathscr{P}) \hookrightarrow \Delta$ and $\mathscr{L}$ is a local system on $\Delta \smallsetminus \mathrm{Sing}(\mathscr{P})$; clearly, there is no $H^0_{\mathrm{c}}(\Delta', \alpha_*j_*\mathscr{L})$ and, arguing by duality, there is no $H^2_{\mathrm{c}}(\Delta', \alpha_*j_*\mathscr{L})$. The computation of the dimension of $H^1_{\mathrm{c}}(\Delta', \alpha_*j_*\mathscr{L})$ is then an exercise. ☐

As topologically, $L'_{<0,\widehat{\theta}'_o}$ is homeomorphic to such a $\Delta'$, we find that $({}^{\mathrm{F}}\mathscr{F})^{<0}_{\widehat{\theta}'_o}$ has cohomology in degree $-1$ only.

Taking into account the description of $\widetilde{E}$ given above, the difference $L'_{\leqslant 0} \smallsetminus L'_{<0} = L'_{\leqslant 0} \cap \widetilde{E}$ is identified to the set $\widetilde{\mathbb{P}}^1_{v'} \times S^1$ with the subset

$$\{|v'| = \infty, \arg v' \in [\pi/2, 3\pi/2] \bmod 2\pi\}$$

deleted. Then $L'_{\leqslant 0,\widehat{\theta}'_o}$ is homeomorphic to the space obtained by gluing $S^1 \times [0,\infty] \smallsetminus \{\arg t_1 \in [\widehat{\theta}'_o + \pi/2, \widehat{\theta}'_o + 3\pi/2] \bmod 2\pi\}$ (with coordinates $(\arg t_1, |t_1|)$) with $\widetilde{\mathbb{P}}^1_{v'} \smallsetminus \{|v'| = \infty, \arg v' \in [\pi/2, 3\pi/2] \bmod 2\pi\}$ along $|t_1| = 0$, $|v'| = \infty$, by identifying $\arg t_1$ with $\arg v' + \widehat{\theta}'_o$. So $L'_{\leqslant 0,\widehat{\theta}'_o}$ is also homeomorphic to $\Delta'$ like in the lemma.

Note that we have an isomorphism $\mathbb{A}^1 \times \widehat{\mathbb{A}}^{1*} \simeq \mathbb{A}^1_{v'} \times \widehat{\mathbb{A}}^{1*}_{\tau'_1}$ given by $(t,\tau) \mapsto (v' = t\tau, \tau'_1 = 1/\tau)$. It follows that the restriction $\mathscr{P}$ of $\boldsymbol{R}(\gamma \circ \alpha)_*\mathscr{G}$ to $\mathbb{A}^1_{v'} \times S^1 \subset \widetilde{E}$ is equal to the pull-back of $\mathscr{F}$ by the map $(v', \arg\tau'_1) \mapsto t = v'e^{i\arg\tau'_1}$. Its restriction to $L'_{\leqslant 0,\widehat{\theta}'_o}$ is therefore isomorphic to the pull-back of $\mathscr{F}$ by the map $\varpi_{\widehat{\theta}'_o} : v' \mapsto v'e^{i\widehat{\theta}'_o}$. In other words, we can regard the picture of $L'_{\leqslant 0,\widehat{\theta}'_o}$ (Figure 2) as the corresponding subset of $\widetilde{\mathbb{P}}^1$ and $\mathscr{P}$ as the restriction of $\widetilde{\jmath}_*\mathscr{F}$ to this subset. We can apply Lemma 7.13 to get that $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0,\widehat{\theta}'_o}$ has cohomology in degree $-1$ only.



Figure 2

Let us now compute $({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0,\widehat{\theta}'_o}/({}^{\mathrm{F}}\mathscr{F})^{<0}_{\widehat{\theta}'_o}$. By the previous computation, this is

$$\boldsymbol{R}\Gamma_{\mathrm{c}}\Big(\widetilde{\mathbb{P}}^1_{v'}\smallsetminus\{|v'|=\infty,\ \arg v'\in[\pi/2,3\pi/2]\bmod 2\pi\},\widetilde{\varpi}^{-1}_{\widehat{\theta}'_o}\mathscr{F}[1]\Big).$$

In the coordinate $t$, let $\Delta$ be a small open disc centered at 0 and let $I_{\widehat{\theta}'_o}$ be the closed interval of $\partial\overline{\Delta}$ defined by $\arg t\in[\widehat{\theta}'_o+\pi/2,\widehat{\theta}'_o+3\pi/2]$. Then the previous complex is the relative cohomology complex $\boldsymbol{R}\Gamma(\overline{\Delta},I_{\widehat{\theta}'_o};\mathscr{F}_{|\overline{\Delta}}[1])$. This complex has cohomology in degree zero at most and $\boldsymbol{H}^0(\overline{\Delta},I_{\widehat{\theta}'_o};\mathscr{F}_{|\overline{\Delta}}[1])\simeq{}^p\phi_t\mathscr{F}$. When $\widehat{\theta}'_o$ runs counterclockwise once around $S^1_{\widehat{\infty}}$, then $I_{\widehat{\theta}'_o}$ also moves counterclockwise once around $\partial\overline{\Delta}$. The monodromy of $\mathrm{gr}_0({}^{\mathrm{F}}\mathscr{F})$ is thus identified with the monodromy on ${}^p\phi_t\mathscr{F}$. $\square$

We now extend this construction in order to define $({}^{\mathrm{F}}\mathscr{F})_{\leqslant c/\tau'}$ for any $c\in\mathbb{C}$. Let $Z(c)\to\mathbb{P}^1\times\widehat{\mathbb{P}}^1$ be the complex blowing-up at the points $(c,\infty)$ and $(\infty,\widehat{0})$, and let $\widetilde{Z}(c)$ be the corresponding real blowing-up space (so that $Z(0)$ and $\widetilde{Z}(0)$ are respectively equal to $Z$ and $\widetilde{Z}$ introduced above). We also define $L''_*(c)$ and $L'_*(c)$, where $*$ is for $<0$ or $\leqslant 0$, as we did for $L''_*$ and $L'_*$, and we denote by $\alpha_c$, etc. the corresponding maps.

***Definition 7.14.*** — For any $c\in\mathbb{C}$, we set

- $({}^{\mathcal{F}}\mathscr{G})_{\leqslant c/\tau'}=\beta_!\boldsymbol{R}(\gamma\circ\alpha)_*\mathscr{G}$ and $({}^{\mathcal{F}}\mathscr{G})^{\ll c/\tau'}=(\beta\circ\gamma)_!\boldsymbol{R}\alpha_*\mathscr{G}$,
- $({}^{\mathrm{F}}\mathscr{F})_{\leqslant c/\tau'}=\boldsymbol{R}\widetilde{\widehat{q}}_*({}^{\mathcal{F}}\mathscr{G})_{\leqslant c/\tau'}$ and $({}^{\mathrm{F}}\mathscr{F})^{<c/\tau'}=\boldsymbol{R}\widetilde{\widehat{q}}_*({}^{\mathcal{F}}\mathscr{G})^{\ll c/\tau'}$.

Clearly, Lemma 7.11 also applies similarly to $({}^{\mathrm{F}}\mathscr{F})_{\leqslant c/\tau'}$ and $({}^{\mathrm{F}}\mathscr{F})^{<c/\tau'}$. It is important to notice that the spaces $\widetilde{Z}(c)$ $(c\in\mathbb{C})$ all coincide when restricted over $\mathbb{P}^1\times\widehat{\mathbb{A}}^1$. As a consequence, the restrictions of $({}^{\mathrm{F}}\mathscr{F})_{\leqslant c/\tau'}$ and $({}^{\mathrm{F}}\mathscr{F})_{<c/\tau'}$ to $\widehat{\mathbb{A}}^1$ are the same perverse sheaf, that we denote by ${}^{\mathrm{F}}\mathscr{F}$. The previous construction defines thus a $\mathcal{I}$-filtration on ${}^{\mathrm{F}}\mathscr{F}$, according to Proposition 7.11 applied with any $c\in\mathbb{C}$. Here, we will denote by $\mathcal{I}$ the sheaf equal to 0 on $\widehat{\mathbb{A}}^1$ and to the constant sheaf with fibre $\mathbb{C}\cdot(1/\tau')$ on $S^1_{\widehat{\infty}}$.

**Proposition 7.15.** — *The triangle* $[({}^{\mathrm{F}}\mathscr{F})^{<c/\tau'} \to ({}^{\mathrm{F}}\mathscr{F})_{\leqslant c/\tau'} \to \mathrm{gr}_{c/\tau}\mathscr{F} \xrightarrow{+1}]_{c\in\mathbb{C}}$ *defines an object of* $\mathrm{St}_{\widehat{\infty}}(\mathbb{C}_{\mathcal{J}\text{ét}}, \leqslant)$ *(see Definition* 4.17*).*

*Proof.* — It is enough to argue on Stokes filtrations and we will use the definition given in Lemma 2.7. As indicated above, it will be enough to take the space $\{c/\tau' \mid c \in \mathbb{C}\}$ as index set. Let us check the filtration property. Let us fix $\widehat{\theta}'_o \in S^1_{\widehat{\infty}}$. From Figure 2, it is clear that if $c$ belongs to $L'_{<0}$, then $({}^{\mathrm{F}}\mathscr{F})_{\leqslant c/\tau', \widehat{\theta}'_o} \subset ({}^{\mathrm{F}}\mathscr{F})^{<0}_{\widehat{\theta}'_o}$. This condition on $c$ reads $\mathrm{Re}(ce^{-i\widehat{\theta}'_o}) < 0$, or equivalently $c/\tau' <_{\widehat{\theta}'_o} 0$. We therefore get a pre-Stokes filtration, with jumps at $c/\tau'$ where $c$ is a singular point of $\mathscr{F}$. The same argument shows that $({}^{\mathrm{F}}\mathscr{F})^{<c/\tau'}$ is the subsheaf defined from $({}^{\mathrm{F}}\mathscr{F})_{\leqslant}$ by Formula (2.2).

Lastly, the dimension property if obtained by the last part of Lemma 7.13. ☐

**Remark 7.16.** — One can ask whether the sheaves ${}^{\mathrm{F}}\mathscr{F}_{\leqslant c/\tau'}$ could be defined without using the blowing-up map $e$ or not. Recall that this blowing-up was used in order to determine in a clear way whether $e^{t/\tau'}$ (resp. $e^{\tau/t'}$) has moderate growth or not near $t = 0$, $\tau' = 0$ (resp. $t' = 0$, $\tau = 0$). As we already indicated in the proof of Proposition 7.11, the blowing up of $(\infty, \widehat{0})$ is not needed. We introduced it by symmetry, and (mainly) in order to treat duality later. On the other hand, working over $\mathbb{A}^1 \times \widehat{X}$, if the sheaves ${}^{\mathcal{F}}\mathscr{G}_{\leqslant c/\tau'}$ were to be defined without blowing-up, they should be equal to $\boldsymbol{R}\widetilde{e}_* {}^{\mathcal{F}}\mathscr{G}_{\leqslant c/\tau'}$, in order that the definition remains consistent. But one can check that $\boldsymbol{R}\widetilde{e}_* {}^{\mathcal{F}}\mathscr{G}_{\leqslant c/\tau'}$ are not sheaves, but complexes, hence do not enter in the frame of Stokes filtrations of a local system in two variables.

### 7.e. Proof of Theorem 7.6 and compatibility with Riemann-Hilbert

Let us first indicate the steps of the proof. We anticipate on the notation and results explained in Lecture 8, which we refer to.

(1) The first step is to compute $\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau})$ (a complex living on $\mathbb{P}^1 \times \widehat{X}$) from a complex defined on $\widetilde{Z}$. We will use the notation of the commutative diagram (7.8). We consider the sheaf $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{Z}}$ on $\widetilde{Z}$ of holomorphic functions on $\widetilde{Z} \smallsetminus \partial\widetilde{Z} = Z \smallsetminus D$ having moderate growth along $\partial\widetilde{Z}$. In particular, $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{Z}}{}_{|\widetilde{Z} \smallsetminus \widetilde{D}} = \mathscr{O}_{Z \smallsetminus D}$. Applying Proposition 8.9 to $e : Z \to \mathbb{P}^1 \times \widehat{\mathbb{P}}^1$ and then its variant to $\varpi \times \mathrm{Id}$ gives

$$\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau}) \simeq \boldsymbol{R}\widetilde{\varepsilon}_* \, \mathrm{DR}^{\mathrm{mod}\,D}\big(e^+(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau})\big). \tag{7.17}$$

**Remark 7.18.** — In fact, (7.17) is a statement similar to Theorem 7.6, but is much easier, in particular because of Proposition 8.9, which would not apply in the setting of Theorem 7.6: indeed, $e$ is a proper modification while $\widehat{p}$ is not.

As a consequence we get

$$\boldsymbol{R}\widehat{q}_* \, \mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau}) \simeq \boldsymbol{R}\widetilde{q}_* \, \mathrm{DR}^{\mathrm{mod}\,D}\big(e^+(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau})\big). \tag{7.19}$$

(2) The second step consists in comparing $\mathrm{DR}^{\mathrm{mod}\,D}\big(e^+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\big)$ with $\beta_!\boldsymbol{R}(\gamma\circ\alpha)_*p^{-1}(\mathscr{F})[-1]$, where $\mathscr{F}=\mathrm{DR}^{\mathrm{an}}\,M[1]$. Both complexes coincide on $\widetilde{Z}\smallsetminus\partial\widetilde{Z}$. We therefore have a natural morphism

$$\mathrm{DR}^{\mathrm{mod}\,D}\big(e^+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\big)\longrightarrow\boldsymbol{R}\beta_*\boldsymbol{R}(\gamma\circ\alpha)_*p^{-1}(\mathscr{F})[-1]. \tag{7.20}$$

That it factorizes through $\beta_!$... would follow from $\delta^{-1}\,\mathrm{DR}^{\mathrm{mod}\,D}\big(e^+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\big)=0$ (which will be proved below), where $\delta$ is the closed inclusion $L''_{\leqslant 0}\hookrightarrow\widetilde{Z}$ complementary to $\beta$ (see Diagram (7.9)). Therefore, proving

(2a) that a morphism $\mathrm{DR}^{\mathrm{mod}\,D}\big(e^+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\big)\to\beta_!\boldsymbol{R}(\gamma\circ\alpha)_*p^{-1}(\mathscr{F})[-1]$ exists

(2b) and that it is an isomorphism

are both local statements on $\partial\widetilde{Z}$. We will also anticipate on results proved in Lecture 8.

Let $C\subset\mathbb{A}^1$ be the union of the singular set of $M$ and $\{0\}$. We also set $C'=C\smallsetminus\{0\}$. We can reduce both local statements to the case where $M$ is supported on $C$ and the case where $M$ is localized along $C$. The first case will be left as an exercise, and we will only consider the second one.

We note that, when expressed in local coordinates adapted to $D$, the pull-back $e^+\mathscr{E}^{t\tau}$ satisfies the assumption in Proposition 8.15 (this is one reason for using the complex blowing-up $e$). Therefore, by a simple inductive argument on the rank, Proposition 8.15 applies to $e^+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ away from $(p\circ e)^{-1}(C')$ (where the polar divisor of $e^+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})$ contains other components than those of $D$). Both local statements are then clear on such a set, by using that $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}\,\mathscr{E}^{t\tau}$ vanishes where $e^{-t\tau}$ does not have moderate growth. We are thus reduced to considering the situation above a neighbourhood of a point $c\in C'$.



We denote by $t_c$ a local coordinate on $\mathbb{A}^1$ centered at $c$ and we set $\tau'=1/\tau$ as above. We will work near the point $(c,\infty)\in Z$ with coordinates $(t_c,\tau')$ (recall that the complex blow-up $e$ is an isomorphism there, so we identify $Z$ and $\mathbb{P}^1\times\widehat{\mathbb{P}}^1$, and we still denote by $Z$ a sufficiently small neighbourhood of $(c,\infty)$). The divisor $D$ is locally defined by $\tau'=0$, and $p^+\mathscr{M}=\mathscr{E}^{(c+t_c)/\tau'}\otimes\mathscr{R}$, where $\mathscr{R}$ is a free $\mathscr{O}_Z(*(D\cup D_c))$-module with a regular connection, setting $D_c=\{t_c=0\}$. We denote by $\mathscr{L}$ the local system on $Z\smallsetminus(D\cup D_c)$ determined by $\mathscr{R}$ (we know that $\mathscr{L}$ has monodromy equal to identity around $\tau' = 0$, but this will not be used in the following). Denoting by $j' : Z \smallsetminus (D \cup D_c) \hookrightarrow Z \smallsetminus D$ the inclusion, we have $p^{-1}\mathscr{F} = \boldsymbol{R}j'_*\mathscr{L}[1]$. We cannot directly apply Proposition 8.15 as above, because $\mathscr{R}$ is localized along $D_c$, so we will first consider the real blow-up $\varpi_c : \widetilde{Z}_c \to \widetilde{Z}$ of $D_c$ in $\widetilde{Z}$ (i.e., we will work in polar coordinates in both variables $\tau'$ and $t_c$).

By Proposition 8.15, the complex $\mathrm{DR}^{\mathrm{mod}\,(D\cup D_c)}(\mathscr{E}^{(c+t_c)/\tau'} \otimes \mathscr{R})$ on $\widetilde{Z}_c$ has cohomology in degree 0 at most, and by (a variant of) Proposition 8.9, we have $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^{(c+t_c)/\tau'} \otimes \mathscr{R}) = \boldsymbol{R}\varpi_{c,*}\,\mathrm{DR}^{\mathrm{mod}\,(D\cup D_c)}(\mathscr{E}^{(c+t_c)/\tau'} \otimes \mathscr{R})$.

Notice now that the open set in $\widetilde{Z}_c$ where $e^{-(c+t_c)/\tau'}$ is exponentially decreasing is nothing but $\varpi'^{-1}L'_{\leqslant 0}$ (see Diagram (7.9)). Therefore, with obvious notation, we have

$$\beta_!\boldsymbol{R}(\gamma \circ \alpha)_*\boldsymbol{R}j_{c,*}\mathscr{L} = \boldsymbol{R}\varpi_{c,*}\beta_{c,!}\boldsymbol{R}(\gamma_c \circ \alpha_c)_*\mathscr{L}.$$

Since both statements (2a) and (2b) hold on $\widetilde{Z}_c$ by the argument given in the first part of Step two, they hold on $\widetilde{Z}$ after applying $\boldsymbol{R}\varpi_{c,*}$.

(3) Third step. We conclude from Step two and (7.19) that we have an isomorphism

$$\boldsymbol{R}\widehat{q}_*\,\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau})[1] \simeq ({}^{\mathrm{F}}\mathscr{F})_{\leqslant 0}[-1].$$

In order to prove the surjectivity of $\mathscr{H}^0(7.4_{\leqslant 0})$ one can restrict to $S^1_{\widehat{\infty}}$, as it holds on $\widehat{\mathbb{A}}^1$. By Proposition 7.11, we conclude that $\boldsymbol{R}\widehat{q}_*\,\mathrm{DR}^{\mathrm{mod}\,\widehat{\infty}}(p^+\mathscr{M} \otimes \mathscr{E}^{t\tau})[1]$ has cohomology in degree zero only, and we have already seen that so has the left-hand term of $(7.4_{\leqslant 0})$. We are thus reduced to showing that the fibers of these sheaves have the same dimension at any $\widehat{\theta} \in S^1_{\widehat{\infty}}$. This follows from Proposition 7.11 and its extension to any index $c/\tau$, showing that this dimension of the right-hand term of $\mathscr{H}^0((7.4_{\leqslant 0}))_{\widehat{\theta}}$ is the sum of dimensions of $\phi_{t-c}\mathscr{F}$, where $c$ varies in the open subset where $0 \leqslant_{\widehat{\theta}} c/\tau$. That the dimension of the left-hand term is computed similarly follows from (b) of §7.a.

(4) In conclusion, we have proved Theorem 7.6 and, at the same time, the fact that the Stokes filtration of Proposition 7.15 is the Stokes filtration of ${}^{\mathrm{F}}M$ at infinity, and more precisely that, through the Riemann-Hilbert correspondence given by ${}^{p}\mathrm{DR}_{\mathscr{J}\text{ét}}$, the image of ${}^{\mathrm{F}}M$ is the triangle of Proposition 7.15. In other words, when $M$ has only regular singularities, the Stokes-de Rham functor (i.e., the de Rham functor enriched with the Stokes filtration at infinity) exchanges the Laplace transform with the topological Laplace transform. □

**7.f. Compatibility of Laplace transformation with duality.** — Let $M$ be a $\mathbb{C}[t]\langle\partial_t\rangle$-module of finite type (an object of $\mathrm{Mod}_f(\mathbb{C}[t]\langle\partial_t\rangle)$), or more generally an object of $D^b_f(\mathbb{C}[t]\langle\partial_t\rangle)$ (bounded derived category of $\mathbb{C}[t]\langle\partial_t\rangle$-modules with finite type cohomology). The Laplace transformation considered in §7.a is in fact a transformation from $\mathrm{Mod}_f(\mathbb{C}[t]\langle\partial_t\rangle)$ (resp. $D^b_f(\mathbb{C}[t]\langle\partial_t\rangle)$) to $\mathrm{Mod}_f(\mathbb{C}[\tau]\langle\partial_\tau\rangle)$ (resp. $D^b_f(\mathbb{C}[\tau]\langle\partial_\tau\rangle)$), and is not restricted to holonomic objects. Let $\boldsymbol{D}$ be the duality functor in these categories. In particular, for a holonomic left $\mathbb{C}[t]\langle\partial_t\rangle$-module $M$,

$\boldsymbol{D}M$ is the complex having cohomology in degree 0 only, this cohomology being the holonomic left $\mathbb{C}[t]\langle\partial_t\rangle$-module $M^\vee$ naturally associated to the right holonomic $\mathbb{C}[t]\langle\partial_t\rangle$-module $\operatorname{Ext}^1_{\mathbb{C}[t]\langle\partial_t\rangle}(M,\mathbb{C}[t]\langle\partial_t\rangle)$, where the right $\mathbb{C}[t]\langle\partial_t\rangle$-module structure comes from the right structure on $\mathbb{C}[t]\langle\partial_t\rangle$.

According to [**49**], there is a canonical isomorphism of functors ${}^{\mathrm{F}}\circ\boldsymbol{D}(\bullet)\xrightarrow{\sim}\iota^+\boldsymbol{D}\circ{}^{\mathrm{F}}(\bullet)$, where $\iota$ denotes the involution $\mathbb{C}[\tau]\langle\partial_\tau\rangle\xrightarrow{\sim}\mathbb{C}[\tau]\langle\partial_\tau\rangle$ sending $\tau$ to $-\tau$ and $\partial_\tau$ to $-\partial_\tau$. This isomorphism can be algebraically described in two ways, depending on the choice of the definition of the Laplace transformation (see §7.a, (1) and (2)). Both definitions coincide (see [**49**, App. 2.4, p. 224]). Let us recall them.

(1) Let us choose a resolution $L^\bullet$ of $M$ by free left $\mathbb{C}[t]\langle\partial_t\rangle$-modules. Then ${}^{\mathrm{F}}L^\bullet$ is a resolution of ${}^{\mathrm{F}}M$ by free left $\mathbb{C}[\tau]\langle\partial_\tau\rangle$-modules, and clearly, by using such a resolution, $\operatorname{Ext}^1_{\mathbb{C}[\tau]\langle\partial_\tau\rangle}({}^{\mathrm{F}}M,\mathbb{C}[\tau]\langle\partial_\tau\rangle)=\iota^{+\mathrm{F}}\operatorname{Ext}^1_{\mathbb{C}[t]\langle\partial_t\rangle}(M,\mathbb{C}[t]\langle\partial_t\rangle)$ as right $\mathbb{C}[\tau]\langle\partial_\tau\rangle$-modules (see [**49**, Lem. V.3.6, p. 86], [**74**, §V.2.b]).

(2) Let us use the definition ${}^{\mathrm{F}}M=\widehat{p}_+(p^+M\otimes E^{t\tau})$. However, since we will have to use the commutation of direct image with duality, it is necessary to work with proper maps (7.1), and we can work either in the algebraic or the analytic setting. We will choose the latter for future use, and we denote by $\mathscr{M}$ the $\mathscr{D}_{\mathbb{P}^1}$-module localized at $\infty$ associated with $M$. Since $p$ is smooth, we have $p^+\boldsymbol{D}(\bullet)=\boldsymbol{D}p^+(\bullet)$ (see [**11**, Prop. VII.9.13], taking into account the difference in notation). In other words, denoting by $(p^+\mathscr{M})^\vee$ the left $\mathscr{D}_{\mathbb{P}^1\times\widehat{\mathbb{P}}^1}$-module associated with $\operatorname{Ext}^2_{\mathscr{D}_{\mathbb{P}^1\times\widehat{\mathbb{P}}^1}}(p^+\mathscr{M},\mathscr{D}_{\mathbb{P}^1\times\widehat{\mathbb{P}}^1})$, we have $p^+(\mathscr{M}^\vee)=(p^+\mathscr{M})^\vee$. We now get $p^+(\boldsymbol{D}\mathscr{M})\otimes\mathscr{E}^{t\tau}=\boldsymbol{D}(p^+\mathscr{M})\otimes\mathscr{E}^{t\tau}=\boldsymbol{D}(p^+\mathscr{M}\otimes\mathscr{E}^{-t\tau})$. Notice also that $\boldsymbol{D}\mathscr{M}$ may not be localized at $\infty$ (as was $\mathscr{M}$ by definition), but we have $p^+(\boldsymbol{D}\mathscr{M})\otimes\mathscr{E}^{t\tau}=p^+\big((\boldsymbol{D}\mathscr{M})(*\infty)\big)\otimes\mathscr{E}^{t\tau}$, since $\mathscr{E}^{t\tau}$ is localized along $D_\infty$. Therefore, $\widehat{p}_+(p^+(\boldsymbol{D}\mathscr{M})\otimes\mathscr{E}^{t\tau})$ is the $\mathscr{D}_{\widehat{\mathbb{P}}^1}$-module localized at $\widehat{\infty}$ associated with ${}^{\mathrm{F}}(\boldsymbol{D}M)$.

By the relative duality for the proper map $\widehat{p}_+$ (see e.g. [**11**, Prop. VII.9.6], [**79**], [**49**, App. 2, Th. 3.3]), we have an isomorphism

$$\widehat{p}_+\boldsymbol{D}(p^+\mathscr{M}\otimes\mathscr{E}^{-t\tau})\xrightarrow{\sim}\boldsymbol{D}\widehat{p}_+(p^+\mathscr{M}\otimes\mathscr{E}^{-t\tau}),$$

from which we get (by applying $\Gamma(\widehat{\mathbb{P}}^1,\bullet)$)

$$ {}^{\mathrm{F}}(\boldsymbol{D}M)\xrightarrow{\sim}\iota^+\boldsymbol{D}{}^{\mathrm{F}}M. \qquad \square$$

The localized module $\mathbb{C}[\tau,\tau^{-1}]\otimes_{\mathbb{C}[\tau]}{}^{\mathrm{F}}M$ is a free $\mathbb{C}[\tau,\tau^{-1}]$-module of finite rank with connection. We denote by ${}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(M)$ the associated Stokes-filtered local system on $S^1_{\widehat{\infty}}$.

Applying the Riemann-Hilbert functor of Definition 5.6 together with the duality isomorphism of Proposition 5.14, we get an isomorphism

$$ {}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(\boldsymbol{D}M)\xrightarrow{\sim}\iota^{-1}[{}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(M)]^\vee. \tag{7.21}$$

## 7.g. Compatibility of topological Laplace transformation with Poincaré-Verdier duality

We will define an analogue of the previous isomorphism by working at the topological level. We will compare both constructions in §7.h. It will be clearer to distinguish between the projection $p:\mathbb{P}^1\times\widehat{\mathbb{P}}^1$ and its restriction over $\mathbb{A}^1$, that we now denote by $p_o:\mathbb{A}^1\times\widehat{\mathbb{P}}^1\to\mathbb{A}^1$ (or $\mathbb{A}^1\times\widehat{\mathbb{A}}^1\to\mathbb{A}^1$).

Let $\mathscr{F}$ be a perverse sheaf of $\boldsymbol{k}$-vector spaces on $\mathbb{A}^1$, with singularity set $C$. The pull-back $\mathscr{G}:=p_o^{-1}\mathscr{F}[1]$ on $\mathbb{A}^1\times\widehat{\mathbb{A}}^1$ is also perverse. Moreover, there is a functorial isomorphism $p_o^{-1}(\boldsymbol{D}\mathscr{F})[1]\xrightarrow{\sim}(p_o^!\boldsymbol{D}\mathscr{F})[-1]=\boldsymbol{D}(p_o^{-1}\mathscr{F}[1])$ (see e.g. [**35**, Prop. 3.3.2]) which is compatible with bi-duality.

Let $\iota$ denote the involution $\tau\mapsto-\tau$ on $\widehat{\mathbb{A}}^1$. It induces an involution on the spaces $\mathbb{A}^1\times\widehat{\mathbb{A}}^1$, $\widehat{X}$, $Z$, $X\times\widehat{X}$ and $\widetilde{Z}$ in a unique way. We all denote them by $\iota$. We will use the notation $\alpha_\iota$ for the conjugate $\iota\circ\alpha\circ\iota$, etc.

**Proposition 7.22.** — *The topological Laplace transformation, from $\boldsymbol{k}$-perverse sheaves to Stokes-$\boldsymbol{k}$-perverse sheaves, is compatible with duality (up to $\iota$) and bi-duality.*

Given a $\boldsymbol{k}$-perverse sheaf $\mathscr{F}$ on $\mathbb{A}^1$, we denote by ${}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(\mathscr{F})$ the Stokes-filtered local system $({}^{\mathrm{F}}\mathscr{F}_\leqslant)_{|S^1_{\widehat\infty}}$. Then, according to Lemma 4.15, the proposition gives an isomorphism

$$
{}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(\boldsymbol{D}\mathscr{F})\xrightarrow{\sim}[{}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(\mathscr{F})]^\vee, \tag{7.23}
$$

where $(\mathscr{L},\mathscr{L}_\bullet)^\vee$ is defined by Proposition 2.15(2).

*Proof of Proposition* 7.22. — We first wish to prove the existence of isomorphisms

$$
{}^{\mathrm{F}}(\boldsymbol{D}\mathscr{F})_{<0}\simeq\iota^{-1}\boldsymbol{D}({}^{\mathrm{F}}\mathscr{F}_{\leqslant0}),\quad {}^{\mathrm{F}}(\boldsymbol{D}\mathscr{F})_{\leqslant0}\simeq\iota^{-1}\boldsymbol{D}({}^{\mathrm{F}}\mathscr{F}_{<0}) \tag{7.24}
$$

which are exchanged by duality up to bi-duality isomorphisms. Note that we could also write $\iota^{-1}{}^{\mathrm{F}}(\boldsymbol{D}\mathscr{F})_{<0}$ as ${}^{\overline{\mathrm{F}}}(\boldsymbol{D}\mathscr{F})_{<0}$ etc., where $\overline{\mathrm{F}}$ denotes the inverse Laplace transformation, which has kernel $e^{-t\tau}$.

These isomorphisms are obtained by applying to similar isomorphisms on $\widetilde{Z}$ the direct image $\boldsymbol{R}\widetilde{\widehat{q}}_*$, which is known to commute with $\boldsymbol{D}$ in a way compatible with bi-duality since $\widetilde{\widehat{q}}$ is proper, according to Poincaré-Verdier duality (see e.g. [**35**, Prop. 3.1.10]).

We will thus first work locally on $\widetilde{Z}$. Let $\mathscr{G}$ be a $\boldsymbol{k}$-perverse sheaf on $\mathbb{A}^1\times\widehat{\mathbb{A}}^1$ with singular set $S=p_o^{-1}(C)$. By Poincaré-Verdier duality, we have $\boldsymbol{D}(\beta_!\boldsymbol{R}(\gamma\circ\alpha)_*\mathscr{G})=\boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\boldsymbol{D}\mathscr{G}$ in a way compatible with bi-duality (see (7.9) for the notation). We denote by $\overline{\delta}$ the closed inclusion $L''_{<0}\hookrightarrow\widetilde{Z}$ and similarly $\overline{\delta}_\iota:\iota(L''_{<0})\hookrightarrow\widetilde{Z}$.

Similarly, $\boldsymbol{D}((\beta\circ\gamma)_!\boldsymbol{R}\alpha_*\mathscr{G})=\boldsymbol{R}(\beta\circ\gamma)_*\alpha_!\boldsymbol{D}\mathscr{G}$ in a way compatible with bi-duality, and we denote by $\delta$ the closed inclusion $L''_{\leqslant0}\hookrightarrow\widetilde{Z}$ and similarly $\delta_\iota:\iota(L''_{\leqslant0})\hookrightarrow\widetilde{Z}$.

Both $\mathscr{G}$ and $\boldsymbol{D}\mathscr{G}$ have the same singular set, and we will later apply the following lemma to $\boldsymbol{D}\mathscr{G}$.

***Lemma 7.25.*** — *If $\mathscr{G}$ satisfies the previous assumptions, we have*

- $\overline{\delta}_\iota^{-1}\boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G}=0$,
- $\delta_\iota^{-1}\boldsymbol{R}(\beta\circ\gamma)_*\alpha_!\mathscr{G}=0$.

*Proof.* — We start with the first equality. By definition, the cohomology of $\boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G}$ is zero on $L'_{\leqslant 0}$, so it is enough to prove that the pull-back of $\boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G}$ to $\iota(L''_{<0})\cap L''_{\leqslant 0}$ is zero. Notice that this set does not cut $\widetilde{E}\cup\widetilde{F}$ by definition of $L''_{\leqslant 0}$. Notice also that, now that we are far from $E$ and $F$, there is no difference between $L''_{<0}$ and $L''_{\leqslant 0}$, so we will forget $\gamma$ in the notation. We will distinguish three cases:

(1) Smooth points of $D_\infty\cup D_{\widehat{\infty}}$ not in $\overline{S}$ (closure of $S$).
(2) The point $(\infty,\widehat{\infty})$.
(3) The points of $(D_\infty\cup D_{\widehat{\infty}})\cap\overline{S}$ (they are smooth points of $D_{\widehat{\infty}}$ by assumption).

(1) Near such a point, $\iota(L''_{\leqslant 0})\cap L''_{\leqslant 0}$ is topologically a product of a similar dimension-one intersection with a disc and $\mathscr{G}$ is a local system (up to a shift). We will therefore treat the dimension-one analogue. We consider a local system $\mathscr{L}$ on $(0,\varepsilon)\times S^1$, where $S^1$ has coordinate $e^{i\theta}$, we set $L''_{\leqslant 0}=\{(0,\theta)\mid\theta\in[\pi/2,3\pi/2]\bmod 2\pi\}$ and $\iota(L''_{\leqslant 0})=L''_{\leqslant 0}+\pi$. Moreover, $\alpha:(0,\varepsilon)\times S^1\hookrightarrow(0,\varepsilon)\times S^1\cup L'_{\leqslant 0}$ and $\beta:(0,\varepsilon)\times S^1\cup L'_{\leqslant 0}\hookrightarrow[0,\varepsilon)\times S^1$ are the inclusions. We wish to show that $\boldsymbol{R}\beta_*\alpha_!\mathscr{L}$ is zero at $(0,\pi/2)$ and $(0,3\pi/2)$. The cohomology of the germ of $\boldsymbol{R}\beta_*\alpha_!\mathscr{L}$ at $\pi/2$ (say) is the cohomology of an open disc with an open interval $I$ added on its boundary, with coefficient in a sheaf which is constant on the open disc and zero on this open interval. This is also the relative cohomology $H^*(\Delta\cup I,I;\boldsymbol{k}^d)$ ($d=\operatorname{rk}\mathscr{L}$). So this is clearly zero.

(2) The pull-back in $\widetilde{Z}$ of a neighbourhood of $(\infty,\overline{\infty})$ takes the form $[0,\varepsilon)^2\times S^1_\infty\times S^1_{\widehat{\infty}}$, with coordinates $(\theta',\widehat{\theta}')$ on $S^1_\infty\times S^1_{\widehat{\infty}}$. The set $L''_{\leqslant 0}$ is defined by $\theta'+\widehat{\theta}'\in[\pi/2,3\pi/2]\bmod 2\pi$ and $\iota(L''_{\leqslant 0})$ by $\theta'+\widehat{\theta}'\in[-\pi/2,\pi/2]\bmod 2\pi$ (since $\iota$ consists in changing $\widehat{\theta}'$ to $\widehat{\theta}'+\pi$). On the other hand, $\mathscr{G}$ is a local system (up to a shift) on $(0,\varepsilon)^2\times S^1_\infty\times S^1_{\widehat{\infty}}$. Up to taking new coordinates $(\theta'+\widehat{\theta}',\theta'-\widehat{\theta}')$, one is reduced to the same computation as in (1), up to the cartesian product by intervals, which has no effect on the result.

(3) As in the proof on Page 96, it is enough to consider the case where $\mathscr{G}$ is supported in $\overline{S}$, which is treated as in (1), and the case where $\mathscr{G}$ is the maximal extension of a local system (up to a shift) away from $D_c$ ($c\in C'$). In this case, the situation is a product of that considered in (1) and that of a local system on an open punctured disc $\Delta^*$. The argument of (1) applies here also.

Let us now consider the second equality, for which we argue similarly. The cohomology of $\boldsymbol{R}(\beta\circ\gamma)_*\alpha_!\mathscr{G}$ is zero on $L'_{<0}$, so it is enough to prove that the pull-back of $\boldsymbol{R}(\beta\circ\gamma)_*\alpha_!\mathscr{G}$ to $\iota(L''_{\leqslant 0})\cap L''_{<0}$ is zero. This set does not cut $\widetilde{E}\cup\widetilde{F}$, so $\gamma=\mathrm{Id}$ on this set, and we are reduced to the previous computation. □

*End of the proof of Proposition* 7.22. — Let $\mathscr{G}$ be as above. As in Definition 7.10, we set ${}^{\mathscr{F}}\!\mathscr{G}_{\leqslant 0} = \beta_! \boldsymbol{R}(\gamma\circ\alpha)_*\mathscr{G}$ and ${}^{\mathscr{F}}\!\mathscr{G}^{\ll 0} = (\beta\circ\gamma)_!\boldsymbol{R}\alpha_*\mathscr{G}$. We therefore get two functors ${}^{\mathscr{F}}(\bullet)_{\leqslant 0}$ and ${}^{\mathscr{F}}(\bullet)^{\ll 0}$. We will show that they are compatible with $\iota$-twisted duality.

***Lemma 7.26.*** — *There exist unique isomorphisms*

$$ {}^{\mathscr{F}}(\iota^{-1}\boldsymbol{D}\mathscr{G})^{\ll 0} \xrightarrow{\lambda_{\leqslant 0}} \iota^{-1}\boldsymbol{D}({}^{\mathscr{F}}\!\mathscr{G}_{\leqslant 0}), \quad {}^{\mathscr{F}}(\iota^{-1}\boldsymbol{D}\mathscr{G})_{\leqslant 0} \xrightarrow{\lambda^{\ll 0}} \iota^{-1}\boldsymbol{D}({}^{\mathscr{F}}\!\mathscr{G}^{\ll 0}) $$

*which extend the identity on $\mathbb{A}^1\times\widehat{\mathbb{A}}^1$. They are functorial, and fit into a bi-duality commutative diagram*

$$\begin{tikzcd}
\iota^{-1}\boldsymbol{D}\,{}^{\mathscr{F}}(\iota^{-1}\boldsymbol{D}\bullet)_{\leqslant 0} & & {}^{\mathscr{F}}(\boldsymbol{D}\boldsymbol{D}\bullet)^{\ll 0} \arrow[ll, "\lambda_{\leqslant 0}\cdot\iota^{-1}\boldsymbol{D}"', "\sim"] \\
\boldsymbol{D}\big[\boldsymbol{D}({}^{\mathscr{F}}(\bullet)^{\ll 0})\big] \arrow[u, "\iota^{-1}\boldsymbol{D}(\lambda^{\ll 0})", "\wr"'] & & \\
{}^{\mathscr{F}}(\bullet)^{\ll 0} \arrow[u, "\mathrm{bid}\cdot{}^{\mathscr{F}}(\bullet)^{\ll 0}", "\wr"'] \arrow[rr, equal] & & {}^{\mathscr{F}}(\bullet)^{\ll 0} \arrow[uu, "\wr", "{}^{\mathscr{F}}(\mathrm{bid})^{\ll 0}"']
\end{tikzcd} \tag{7.27}$$

*and a simlar one by exchanging $\leqslant 0$ and $\ll 0$ (and we use the identity $\iota^{-1}\boldsymbol{D} = \boldsymbol{D}\iota^{-1}$).*

*Proof.* — We will use the cartesian square of open inclusions:

$$\begin{tikzcd}
\iota(L'_{<0}) \arrow[rr, hook, "\beta_\iota\circ\gamma_\iota"] & & \widetilde{Z} \\
\mathbb{A}^1\times\widehat{\mathbb{A}}^1 = \iota(L'_{<0})\cap L'_{\leqslant 0} \arrow[u, hook, "\alpha_\iota"] \arrow[rr, hook, "\gamma\circ\alpha"] & & L'_{\leqslant 0} \arrow[u, hook, "\beta"']
\end{tikzcd}$$

which implies (see [**35**, Prop. 2.5.11]) $(\gamma\circ\alpha)_!\alpha_\iota^{-1} = \beta^{-1}(\beta_\iota\circ\gamma_\iota)_!$ and $\alpha_{\iota,!}(\gamma\circ\alpha)^{-1} = (\beta_\iota\circ\gamma_\iota)^{-1}\beta_!$, and, by openness, $\boldsymbol{R}(\gamma\circ\alpha)_*\alpha_\iota^{-1} = \beta^{-1}\boldsymbol{R}(\beta_\iota\circ\gamma_\iota)_*$ and $(\beta_\iota\circ\gamma_\iota)^{-1}\boldsymbol{R}\beta_* = \boldsymbol{R}\alpha_{\iota,*}(\gamma\circ\alpha)^{-1}$.

For any $\mathscr{G}$ as above, the previous remark and the first assertion of Lemma 7.25 implies that the following natural adjunction morphism

$$(\beta_\iota\circ\gamma_\iota)_!\boldsymbol{R}\alpha_{\iota,*}\mathscr{G} = (\beta_\iota\circ\gamma_\iota)_!(\beta_\iota\circ\gamma_\iota)^{-1}\boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G} \longrightarrow \boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G}$$

is an isomorphism. We have by adjunction (see [**35**, (2.6.14) & Th. 3.15]):

$$\begin{aligned}
\operatorname{Hom}((\beta_\iota\circ\gamma_\iota)_!\boldsymbol{R}\alpha_{\iota,*}\mathscr{G}, \boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G}) &= \operatorname{Hom}(\beta^{-1}(\beta_\iota\circ\gamma_\iota)_!\boldsymbol{R}\alpha_{\iota,*}\mathscr{G}, (\gamma\circ\alpha)_!\mathscr{G}) \\
&= \operatorname{Hom}((\gamma\circ\alpha)_!\mathscr{G}, (\gamma\circ\alpha)_!\mathscr{G}) \\
&= \operatorname{Hom}(\mathscr{G},\mathscr{G}),
\end{aligned}$$

where the last equality follows from $(\gamma\circ\alpha)^! = (\gamma\circ\alpha)^{-1}$ (since $\gamma\circ\alpha$ is open). Therefore, we can also express the previous adjunction morphism as the adjunction

$$(\beta_\iota\circ\gamma_\iota)_!\boldsymbol{R}\alpha_{\iota,*}\mathscr{G} \longrightarrow \boldsymbol{R}\beta_*\beta^{-1}(\beta_\iota\circ\gamma_\iota)_!\boldsymbol{R}\alpha_{\iota,*}\mathscr{G} = \boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\mathscr{G}.$$

Note that it is directly seen to be an isomorphism by dualizing the second assertion of Lemma 7.25.

Applying these isomorphisms to $\boldsymbol{D}\mathscr{G}$ instead of $\mathscr{G}$, the $D^b(\widetilde{Z},\boldsymbol{k})$-isomorphism

$$(\beta_\iota\circ\gamma_\iota)_!\boldsymbol{R}\alpha_{\iota,*}\boldsymbol{D}\mathscr{G}\xrightarrow{\sim}\boldsymbol{R}\beta_*(\gamma\circ\alpha)_!\boldsymbol{D}\mathscr{G}=\boldsymbol{D}(\beta_!\boldsymbol{R}(\gamma\circ\alpha)_*\mathscr{G})$$

that we deduce is by definition $\lambda_{\leqslant 0}$. Uniqueness follows from the identity of Hom above. Now, $\lambda^{\ll 0}$ is defined in order that (7.27) commutes. It can be defined as $\lambda_{\leqslant 0}$ by using the two other possible adjunction morphisms, by the same uniqueness argument. □

The proof of (7.24) now follows: on the one hand, $\boldsymbol{D}(p_o^{-1}\mathscr{F}[1])\simeq p_o^{-1}(\boldsymbol{D}\mathscr{F})[1]$ since $p_o$ is smooth of real relative dimension two, and, on the other hand, Poincaré-Verdier duality can be applied to $\boldsymbol{R}\widetilde{\widehat{q}}_*$ since $\widetilde{\widehat{q}}_*$ is proper; one concludes by noticing that $\iota^{-1}p_o^{-1}\mathscr{F}=p_o^{-1}\mathscr{F}$, since $p_o\circ\iota=p_o$.

The proof of the proposition is obtained by applying the same reasoning to each $c\in\mathbb{C}$ and ${}^{\mathrm{F}}\mathscr{F}_{\leqslant c/\tau'}$, ${}^{\mathrm{F}}\mathscr{F}_{<c/\tau'}$. □

**7.h. Comparison of both duality isomorphisms.** — The purpose of this section is to show that, through the Riemann-Hilbert correspondence $\mathscr{F}={}^p\mathrm{DR}^{\mathrm{an}}M$, the duality isomorphisms (7.21) and (7.23) correspond each other. We will neglect questions of signs, so the correspondence we prove has to be understood "up to sign".

Let us make this more precise. By the compatibility of the Riemann-Hilbert correspondence with Laplace and topological Laplace transformations, shown in §7.e(4), we have a natural isomorphism

$${}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)({}^p\mathrm{DR}^{\mathrm{an}}M)\simeq{}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(M).$$

By using the local duality theorem $\boldsymbol{D}\,{}^p\mathrm{DR}^{\mathrm{an}}M\simeq{}^p\mathrm{DR}^{\mathrm{an}}\boldsymbol{D}M$ (see [**66**] and the references given therein), (7.21) gives rise to an isomorphism

$$\text{(7.28)}\qquad {}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)(\boldsymbol{D}{}^p\mathrm{DR}^{\mathrm{an}}M)\xrightarrow{\sim}[{}^{\mathrm{F}}(\mathscr{L},\mathscr{L}_\bullet)({}^p\mathrm{DR}^{\mathrm{an}}M)]^\vee.$$

Since both source and target of (7.23) and (7.28) are identical, proving that both isomorphisms coincide amounts to proving that their restriction to the corresponding local systems coincide, through the previously chosen identifications. These identifications will be easier to follow if we restrict to the local systems, that is, to $\tau\in\mathbb{C}^*$.

From now on, the notation in the diagram (7.1) will be understood with $\mathbb{C}^*_\tau$ replacing $\widehat{\mathbb{P}}^1$, and we will also consider the restriction $p_o$ of $p$ to $\mathbb{A}^1\times\mathbb{C}^*_\tau$. We denote by $\mathscr{M}$ the $\mathscr{D}_{\mathbb{P}^1}$-module localized at $\infty$ corresponding to $M$ and we set $\mathscr{F}={}^p\mathrm{DR}^{\mathrm{an}}M$. With this understood, we have

$$({}^{\mathrm{F}}M)^{\mathrm{an}}=\widehat{p}_+(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}),\quad {}^{\mathrm{F}}\mathscr{F}=\boldsymbol{R}\widehat{p}_*\boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_*p_o^{-1}\mathscr{F}).$$

We now consider the functorial isomorphism

$$\text{(7.29)}\qquad \boldsymbol{D}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau})\simeq\boldsymbol{D}(p^+\mathscr{M})\otimes\mathscr{E}^{-t\tau}\simeq p^+(\boldsymbol{D}\mathscr{M})\otimes\mathscr{E}^{-t\tau}.$$

Let us denote by $M^{\mathrm{an}}$ the restriction of $\mathscr{M}$ to $\mathbb{A}^1$. We have a natural isomorphism of $\mathscr{O}^{\mathrm{an}}_{\mathbb{A}^1\times\widehat{\mathbb{A}}^1}$-modules with connection: $(\mathscr{O}, d) \xrightarrow{\sim} (\mathscr{E}^{t\tau}, \nabla)$, sending 1 to $e^{-t\tau}\cdot 1$. Up to this isomorphism, and its dual, the restriction of (7.29) to $\mathbb{A}^1 \times \mathbb{C}^*_\tau$ is the corresponding isomorphism

$$\boldsymbol{D}(p_o^+ M^{\mathrm{an}}) \simeq p_o^+(\boldsymbol{D}M^{\mathrm{an}}).$$

On the other hand, the isomorphism associated to (7.29), obtained by using the local duality theorem:

$$\boldsymbol{D}\,{}^p\mathrm{DR}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}) \xrightarrow{\sim} {}^p\mathrm{DR}\boldsymbol{D}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}) \xrightarrow{\sim} {}^p\mathrm{DR}(p^+(\boldsymbol{D}\mathscr{M})\otimes\mathscr{E}^{-t\tau}) \tag{7.30}$$

restricts similarly to

$$\boldsymbol{D}^p\mathrm{DR}(p_o^+ M^{\mathrm{an}}) \xrightarrow{\sim} {}^p\mathrm{DR}\boldsymbol{D}(p_o^+ M^{\mathrm{an}}) \simeq {}^p\mathrm{DR}(p_o^+(\boldsymbol{D}M^{\mathrm{an}})). \tag{7.31}$$

Recall now that (7.20) (restricted over $\mathbb{C}^*_\tau$) induces, by applying $\boldsymbol{R}\varpi_*$, an isomorphism

$${}^p\mathrm{DR}(p^+\mathscr{M}\otimes\mathscr{E}^{t\tau}) \xrightarrow{\sim} \boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_* p_o^{-1}\mathscr{F}).$$

Hence (7.30) gives rise to an isomorphism

$$\boldsymbol{D}\boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_* p_o^{-1}\mathscr{F}[1]) \xrightarrow{\sim} \iota^{-1}\boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_* p_o^{-1}(\boldsymbol{D}\mathscr{F})[1]), \tag{7.32}$$

by using the local duality $\boldsymbol{D}\,{}^p\mathrm{DR}M^{\mathrm{an}} \simeq {}^p\mathrm{DR}\boldsymbol{D}M^{\mathrm{an}}$.

On the other hand, the isomorphisms of Lemma 7.26 applied with $\mathscr{G} = p_o^{-1}\mathscr{F}$ also give an isomorphism

$$\boldsymbol{D}\,\boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_* p_o^{-1}\mathscr{F}[1]) \xrightarrow{\sim} \iota^{-1}\boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_* p_o^{-1}(\boldsymbol{D}\mathscr{F})[1]), \tag{7.33}$$

since here $\gamma = \mathrm{Id}$ and there is no distinction between ${}^{\mathcal{F}}\mathscr{G}_{\leqslant}0$ and ${}^{\mathcal{F}}\mathscr{G}^{\ll 0}$.

**Lemma 7.34.** — *The isomorphisms* (7.32) *and* (7.33) *coincide (up to a nonzero constant).*

We first start by proving:

**Lemma 7.35.** — *The isomorphisms* (7.32) *and* (7.33) *are completely determined by their restriction to* $\mathbb{A}^1 \times \mathbb{C}^*_\tau$.

*Proof.* — We will use the following notation: $\Delta_\infty$ is a disc in $\mathbb{P}^1$ centered at $\infty$, $\varpi : \widetilde{\Delta}_\infty \to \Delta_\infty$ is the oriented real blow-up, $\Delta^*_\infty$ is the punctured disc. The open subset $L'_{\leqslant 0} \subset \partial\Delta_\infty \times \mathbb{C}^*_\tau$ is defined as in §7.c. In particular, the fiber of $\varpi : L'_{\leqslant 0} \to \{\infty\} \times \mathbb{C}^*_\tau$ is an open half-circle in $\varpi^{-1}(\infty, \tau)$. As above, we denote by $\widetilde{\jmath} = \beta\circ\alpha$ the corresponding decomposition of the inclusion $\widetilde{\jmath} : \Delta^*_\infty \times \mathbb{C}^*_\tau \hookrightarrow \widetilde{\Delta}_\infty \times \mathbb{C}^*_\tau$, and we still denote by $i_\partial$ the inclusion $\partial\Delta_\infty \times \mathbb{C}^*_\tau \hookrightarrow \Delta_\infty \times \mathbb{C}^*_\tau$. Let $\mathscr{G}$ be a locally constant sheaf on $\Delta^*_\infty \times \mathbb{C}^*_\tau$.

**Lemma 7.36.** — *With this notation, any automorphism of* $\boldsymbol{R}\varpi_*\beta_!\boldsymbol{R}\alpha_*\mathscr{G}$ *is uniquely determined from its restriction to* $\Delta^*_\infty \times \mathbb{C}^*_\tau$.

*Proof.* — We notice that $\boldsymbol{R}\varpi_*\beta_!\boldsymbol{R}\alpha_*\mathscr{G}$ is a sheaf, equal to $\varpi_*\beta_!\alpha_*\mathscr{G}$: indeed, this is clearly so for $\beta_!\alpha_*\mathscr{G}$, whose restriction to $\partial\Delta_\infty\times\mathbb{C}^*_\tau$ is a local system on $L'_{\leqslant 0}$ extended by 0; since $\varpi$ is proper, it is enough to check that the push-forward of the latter sheaf is also a sheaf, which is clear.

By adjunction, we have

$$\begin{aligned}\operatorname{Hom}(\boldsymbol{R}\varpi_*\beta_!\boldsymbol{R}\alpha_*\mathscr{G},\boldsymbol{R}\varpi_*\beta_!\boldsymbol{R}\alpha_*\mathscr{G}) &= \operatorname{Hom}(\varpi^{-1}\boldsymbol{R}\varpi_*\beta_!\boldsymbol{R}\alpha_*\mathscr{G},\beta_!\boldsymbol{R}\alpha_*\mathscr{G})\\ &= \operatorname{Hom}(\varpi^{-1}\varpi_*\beta_!\alpha_*\mathscr{G},\beta_!\alpha_*\mathscr{G}).\end{aligned}$$

On the one hand, $i_\partial^{-1}\varpi^{-1}\varpi_*\beta_!\alpha_*\mathscr{G}$ is a local system $\mathscr{H}'$ on $\partial\Delta_\infty\times\mathbb{C}^*_\tau$, and on the other hand $i_\partial^{-1}\beta_!\alpha_*\mathscr{G}$ is a local system on $L'_{\leqslant 0}$ extended by 0, and is a subsheaf of the local system $\mathscr{H} := i_\partial^{-1}\widetilde{\jmath}_*\mathscr{G}$. If $\lambda$ belongs to $\operatorname{Hom}(\varpi^{-1}\varpi_*\beta_!\alpha_*\mathscr{G},\beta_!\alpha_*\mathscr{G})$, then $i_\partial^{-1}\lambda$ induces a morphism $\mathscr{H}'\to\mathscr{H}$ which vanishes on $L''_{\leqslant 0}$, hence is zero. Therefore, $\lambda$ is uniquely determined by its restriction to $\Delta^*_\infty\times\mathbb{C}^*_\tau$. ☐

In order to end the proof of Lemma 7.35, we notice that $\boldsymbol{R}\varpi_*(\beta_!\boldsymbol{R}\alpha_*p_o^{-1}\mathscr{F}[1])$ is of the form considered in Lemma 7.36 in the neighbourhood of $D_\infty$. ☐

*Proof of Lemma* 7.34. — According to Lemma 7.34, we are reduced to proving the coincidence on $\mathbb{A}^1\times\mathbb{C}^*_\tau$. This is given by Lemma 7.37 below. ☐

***Lemma 7.37.*** — *The functorial duality isomorphism* $\boldsymbol{D}\circ p_o^+(\bullet)\xrightarrow{\sim}p_o^+\circ\boldsymbol{D}(\bullet)$, *in the category of analytic holonomic modules, is compatible (up to a nonzero constant), via the de Rham functor, to the functorial isomorphism* $\boldsymbol{D}\circ p_o^{-1}(\bullet)\xrightarrow{\sim}p_o^{-1}\circ\boldsymbol{D}(\bullet)[2]$ *in the category of perverse sheaves.*

*Proof.* — See [**41**, Cor. 5.6.8]. ☐

***Proposition 7.38.*** — *The isomorphisms* (7.23) *and* (7.28) *coincide (up to nonzero constant).*

*Proof.* — According to the relative duality theorem already mentioned, the isomorphism (7.28) is also obtained first by applying $\widehat{p}_+$ to (7.29) and then applying ${}^p\mathrm{DR}$. By [**79**, Th. 3.13], we can first apply ${}^p\mathrm{DR}$ to (7.29) and then $\boldsymbol{R}\widehat{p}_*$. It is thus obtained by applying $\boldsymbol{R}\widehat{p}_*$ to (7.30) or equivalently to (7.32), and then by using Verdier duality for $\boldsymbol{R}\widehat{p}_*$ (i.e., commuting $\boldsymbol{D}$ and $\boldsymbol{R}\widehat{p}_*$).

On the other hand, (7.23) is obtained by applying $\boldsymbol{R}\widehat{p}_*$ to (7.33) and then by using Verdier duality for $\boldsymbol{R}\widehat{p}_*$. The conclusion follows then from Lemma 7.34. ☐

# PART II

# DIMENSION TWO AND MORE

# LECTURE 8

# REAL BLOW-UP SPACES AND MODERATE DE RHAM COMPLEXES

**Summary.** The purpose of this lecture is to give a global construction of the real blow-up space of a complex manifold along a family of divisors. On this space is defined the sheaf of holomorphic functions with moderate growth, whose basic properties are analyzed. The moderate de Rham complex of a meromorphic connection is introduced, and its behaviour under the direct image by a proper modification is explained. This lecture ends with an example of a moderate de Rham complex having cohomology in degree $\geqslant 1$, making a possible definition of Stokes-perverse sheaves more complicated than in dimension one.

**8.a. Real blow-up.** — Recall that the real blow-up $\widetilde{\mathbb{C}}^\ell$ of $\mathbb{C}^\ell$ along $t_1, \dots, t_\ell$ is the space of polar coordinates in each variable $t_j$, that is, the product $(S^1 \times \mathbb{R}_+)^\ell$ with coordinates $(e^{i\theta_j}, \rho_j)_{j=1,\dots,\ell}$ and $t_j = \rho_j e^{i\theta_j}$. The real blow-up map $\varpi : \widetilde{\mathbb{C}}^\ell \to \mathbb{C}^\ell$ induces a diffeomorphism $\{\rho_1 \cdots \rho_\ell \neq 0\} =: (\widetilde{\mathbb{C}}^\ell)^* \xrightarrow{\sim} (\mathbb{C}^\ell)^* := \{t_1 \cdots t_\ell \neq 0\}$.

*Real blow-up along a divisor.* — Let $X$ be a reduced complex analytic space (e.g. a complex manifold) and let $f : X \to \mathbb{C}$ be a holomorphic function on $X$ with zero set $X_0 = X_0(f)$. The oriented real blow-up of $X$ along $f$, denoted by $\widetilde{X}(f)$, is the closure in $X \times S^1$ of the graph of the map $f/|f| : X^* = X \smallsetminus X_0 \to S^1$. The real blowing-up map $\varpi : \widetilde{X} \to X$ is the map induced by the first projection. The inverse image $\varpi^{-1}(X_0)$, that we denote by $\partial\widetilde{X}$, is a priori contained in $X_0 \times S^1$.

***Lemma 8.1.*** — *We have* $\partial\widetilde{X} = X_0 \times S^1$.

*Proof.* — This is a local question on $X_0$. As $f$ is open, for $x_o \in X_0$ there exists a fundamental system $(U_m)_{m\in\mathbb{N}}$ of open neighbourhoods of $x_o$ and a decreasing family $\Delta_m$ of open discs centered at 0 in $\mathbb{C}$ such that $f : U_m \to \Delta_m$ is onto, as well as $f : U_m^* = U_m \smallsetminus X_0 \to \Delta_m \smallsetminus \{0\}$. It follows that, given any $e^{i\theta_o} \in S^1$, there exists $x_m \in U_m^*$ with $f(x_m)/|f(x_m)| = e^{i\theta_o}$, so $(x_o, e^{i\theta_o}) \in \widetilde{X}$. □

As a consequence, $\widetilde{X}$ is equal to the subset of $X \times S^1$ defined by the (in)equation $fe^{-i\theta} \in \mathbb{R}_+$, so is a real semi-analytic subset of $X \times S^1$.

Let now $D$ be a locally principal divisor in $X$ and let $(U_\alpha)_{\alpha\in A}$ be a locally finite covering of $X$ by open sets $U_\alpha$ such that in each $U_\alpha$, the divisor $D$ is defined by a holomorphic function $f^{(\alpha)}$. The data $[U_\alpha, f^{(\alpha)}]_{\alpha\in A}$ allow one to define, by gluing the real blow-ups $\widetilde{U}_\alpha(f^{(\alpha)})$, a space $\widetilde{X}(D)$. Set $f^{(\alpha)} = u^{(\alpha,\beta)}f^{(\beta)}$ on $U_\alpha\cap U_\beta$. The gluing map is induced by

$$\begin{aligned}(U_\alpha\cap U_\beta)\times(\mathbb{C}^*/\mathbb{R}_+^*) &\longrightarrow (U_\alpha\cap U_\beta)\times(\mathbb{C}^*/\mathbb{R}_+^*)\\ \big(x,(e^{i\theta})\big) &\longmapsto \big(x,(u^{(\alpha,\beta)}e^{i\theta} \bmod \mathbb{R}_+^*)\big).\end{aligned}$$

One checks that the space $\widetilde{X}(D)$ does not depend on the choices made (up to a unique homeomorphism compatible with the projection to $X$).

In a more intrinsic way, let $L(D)$ be the rank-one bundle over $X$ associated with $D$ (with associated sheaf $\mathscr{O}_X(D)$) and let $S^1L(D)$ be the corresponding $S^1$-bundle. Let us fix a section $f:\mathscr{O}_X\to\mathscr{O}_X(D)$. It vanishes exactly along $D$ and induces a holomorphic map $X^*:=X\smallsetminus D\to L(D)\smallsetminus D$, that we compose with the projection $L(D)\smallsetminus D\to S^1L(D)$. Then $\widetilde{X}(D)$ is the closure in $S^1L(D)$ of the image of $X^*$ by this map. From Lemma 8.1 we deduce that $\partial\widetilde{X}(D)=S^1L(D)_{|D}$. If $g=u\cdot f$ with $u\in\Gamma(X,\mathscr{O}_X^*)$, then both constructions give homeomorphic blow-up spaces. More precisely, denoting by $\varpi_f:\widetilde{X}(f)\to X$ the real blowing-up map obtained with the section $f$, the multiplication by $u$ induces the multiplication by $u/|u|$ on $S^1L(D)$, which sends the subspace $\widetilde{X}(f)$ to $\widetilde{X}(g)$. Moreover, this is the unique homeomorphism $\widetilde{X}(f)\xrightarrow{\sim}\widetilde{X}(g)$ making the following diagram

$$\begin{tikzcd}
\widetilde{X}(f) \arrow[r, "\sim"] \arrow[d, "\varpi_f"'] & \widetilde{X}(g) \arrow[d, "\varpi_g"] \\
X \arrow[r, equal] & X
\end{tikzcd}$$

commute. This explains the notation and terminology for *the* real blow-up space of $X$ along $D$.

*Real blow-up along a family of divisors.* — Let now $(D_j)_{j\in J}$ be a locally finite family of locally principal divisors in $X$ and let $f_j$ be sections $\mathscr{O}_X\to\mathscr{O}_X(D_j)$. The fibre product over $X$ of the $\widetilde{X}(D_j)$ (each defined with $f_j$), when restricted over $X_J^*:=X\smallsetminus\bigcup_j D_j$, is isomorphic to $X_J^*$. We then define the real blow-up $\widetilde{X}(D_{j\in J})$ as the closure of $X_J^*$ in this fibre product. If $J$ is finite, $\widetilde{X}(D_{j\in J})$ is the closure in the direct sum bundle $\bigoplus_j S^1L(D_j)$ of the image of the section $(f_j/|f_j|)_{j\in J}$ on $X^*$. It is defined up to unique homeomorphism compatible with the projection to $X$. We usually regard $X_J^*$ as an open analytic submanifold in $\widetilde{X}(D_{j\in J})$.

The closure $\widetilde{X}(D_{j\in J})$ of $X^*$ can be strictly smaller than the fibre product of the $\widetilde{X}(D_j)$ (e.g. consider $\widetilde{X}(D,D)$ for twice the same divisor, and more generally when the $D_j$ have common components). Here is an example when both are the same.

***Lemma 8.2.*** — *Assume that $X$ and each $D_j$ is smooth and that the family $(D_j)_{j\in J}$ defines a normal crossing divisor $D = \bigcup_j D_j$ in $X$. Then $\widetilde{X}(D_{j\in J})$ is equal to the fibre product (over $X$) of the $\widetilde{X}(D_j)$ for $j \in J$ and we have a natural proper surjective map $\widetilde{X}(D_{j\in J}) \to \widetilde{X}(D)$.*

*Proof.* — The first assertion is checked locally. For instance, in the case of two divisors crossing normally, we are reduced to checking that

$$\big[(S^1\times\mathbb{R}_+)\times\mathbb{C}\big]\underset{\mathbb{C}\times\mathbb{C}}{\times}\big[\mathbb{C}\times(S^1\times\mathbb{R}_+)\big]\simeq(S^1\times\mathbb{R}_+)\times(S^1\times\mathbb{R}_+). \qquad \square$$

***Corollary 8.3.*** — *Under the assumptions of Lemma 8.2, if $\mathscr{F}^*$ is a local system on $X^*$ and $\widetilde{\jmath}: X^* \hookrightarrow \widetilde{X}(D_{j\in J})$ denotes the open inclusion, then $\boldsymbol{R}\widetilde{\jmath}_*\mathscr{F}^* = \widetilde{\jmath}_*\mathscr{F}^*$ is a local system on $\widetilde{X}$.*

*Proof.* — The question is local and, according to the lemma, we can locally regard $\widetilde{\jmath}$ as being the inclusion $(\mathbb{R}_+^*)^\ell\times\mathbb{R}^\ell\times\mathbb{C}^{n-\ell}\hookrightarrow\mathbb{R}_+^\ell\times\mathbb{R}^\ell\times\mathbb{C}^{n-\ell}$, and we are mainly reduced to consider the inclusion of the open octant $(\mathbb{R}_+^*)^\ell$ into the closed octant $\mathbb{R}_+^\ell$. Then the assertion is clear. $\square$

*Morphisms between real blow-up spaces.* — For any locally principal divisor $D$ in $X$ and any integer $n \geqslant 1$, there is a natural morphism $\mathscr{O}(D)\to\mathscr{O}(nD)$, inducing $L(D)\to L(nD)$ and $\widetilde{X}(D)\to\widetilde{X}(nD)$, which is the identity on $X^*$.

More generally, let $(D_{j\in J})$ be a finite family of locally principal divisors and let $(n_{ij})$ ($i\in I$, $j\in J$) be a finite family of nonnegative integers. Set $E_i=\sum_j n_{ij}D_j$, so that in particular the support of $(E_{i\in I})$ is contained in the support of $(D_{j\in J})$. Then the identity morphism $\mathrm{Id}: X\to X$ lifts as a morphism $\widetilde{X}(D_{j\in J})\to\widetilde{X}(E_{i\in I})$. Indeed, for any $i\in I$, one has a natural morphism $\bigoplus_j L(D_j)\to L(E_i)$, and taking the direct sums of such morphisms when $i$ varies induces the desired lifting of Id.

In particular, if $J'$ is a subset of $J$, there is a natural projection map between the fibre products, which induces a proper surjective map $\widetilde{X}(D_{j\in J})\to\widetilde{X}(D_{j\in J'})$.

Similarly, defining locally $D=\bigcup_j D_j$ by the product of the local equations of the $D_j$, we have a proper surjective map $\widetilde{X}(D_{j\in J})\to\widetilde{X}(D)$.

Given a morphism $\pi: X'\to X$ and a family $(D_{j\in J})$ of divisors of $X$, let $(E_{j\in J})$ be the pull-back family in $X'$. Then there is a natural morphism $\widetilde{\pi}:\widetilde{X}'(E_{j\in J})\to X(D_{j\in J})$. In particular, if we are given a family of divisors $(E'_{i\in I})$ in $X'$ such that $E_j=\sum_i n_{ji}E'_i$ for any $j$ ($n_{ji}\in\mathbb{N}$), we get a natural morphism $\widetilde{\pi}:\widetilde{X}'(E'_{i\in I})\to\widetilde{X}(D_{j\in J})$.

For example, if $\pi$ is chosen such that the divisor $E=\pi^*(\sum_j D_j)$ has simple normal crossings, we can choose for $(E'_{i\in I})$ the family of reduced irreducible components of $E$.

## 8.b. The sheaf of functions with moderate growth on the real blow-up space.

— We consider as above a locally finite family $(D_j)_{j\in J}$ of effective divisors in a smooth complex manifold $X$, and we set $D=\bigcup_j|D_j|$, where $|D_j|$ denotes the support of $D_j$. Let $\mathscr{O}_X(*D)$ denotes the sheaf of meromorphic functions on $X$ with poles along $D$ at most. It can also be defined as the subsheaf of $j_*\mathscr{O}_{X^*}$ (with $j : X^* = X \smallsetminus D \hookrightarrow X$ the open inclusion) consisting of holomorphic functions having moderate growth along $D$.

We define a similar sheaf on $\widetilde{X} := \widetilde{X}(D_{j\in J})$, that we denote by $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$: Given an open set $\widetilde{U}$ of $\widetilde{X}$, a section $f$ of $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$ on $\widetilde{U}$ is a holomorphic function on $U^* := \widetilde{U} \cap X^*$ such that, for any compact set $K$ in $\widetilde{U}$, in the neighbourhood of which $D$ is defined by $g_K \in \mathscr{O}_X(K)$, there exists constants $C_K > 0$ and $N_K \geqslant 0$ such that $|f| \leqslant C_K |g_K|^{-N_K}$ on $K$.

***Remark 8.4* (Rapid decay).** — We will also use the sheaf $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ of holomorphic functions having rapid decay along $\partial\widetilde{X}$: Given an open set $\widetilde{U}$ of $\widetilde{X}$, a section $f$ of $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ on $\widetilde{U}$ is a holomorphic function on $U^* := \widetilde{U} \cap X^*$ such that, for any compact set $K$ in $\widetilde{U}$, in the neighbourhood of which $D$ is defined by $g_K \in \mathscr{O}_X(K)$, and for any $N \in \mathbb{N}$, there exists constants $C_{K,N} > 0$ such that $|f| \leqslant C_K |g_K|^N$ on $K$.

***Proposition 8.5* (Faithful flatness, see [72, Prop. 2.8]).** — *If* $\dim X \leqslant 2$*, the sheaves* $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$ *and* $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ *are flat over* $\varpi^{-1}\mathscr{O}_X(*D)$ *or* $\varpi^{-1}\mathscr{O}_X$*, faithfully over* $\varpi^{-1}\mathscr{O}_X(*D)$*.*

***Remark 8.6*.** — T. Mochizuki has recently shown similar results in higher dimension. More precisely, relying on the basic theorems in [**44**, Chap. VI], he proves the statement for $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ (and more general sheaves defined with rapid decay condition). Concerning moderate growth, the trick is to use, instead of the sheaf of holomorphic functions with moderate growth along $D$, which is very big, the subsheaf of such functions of the Nilsson class, as in [**16**, p. 45] (sheaf denoted there by $\mathscr{A}_{\log}$) or in [**49**, p. 61] (sheaf denoted there by $\mathscr{O}^{\mathrm{Nils}}$). This trick is useful when $D$ has normal crossings. The flatness of this subsheaf is also a consequence of basic flatness results.

*Proof*. — We will give the proof for $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$, the proof for $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ being completely similar. Let us fix $x_o \in D$ and $\widetilde{x}_o \in \varpi^{-1}(x_o) \subset \partial\widetilde{X}$. The case where $\dim X = 1$ is clear, because $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$ has no $\mathscr{O}_X$-torsion.

*Faithful flatness*. — Notice first that, if flatness is proved, the faithful flatness over $\varpi^{-1}\mathscr{O}_X(*D)$ is easy: If $\mathscr{M}_{x_o}$ has finite type over $\mathscr{O}_{X,x_o}(*D)$ and $\mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{mod}\,D} \otimes_{\mathscr{O}_{X,x_o}(*D)} \mathscr{M}_{x_o} = 0$ then, extending locally $\mathscr{M}_{x_o}$ as a $\mathscr{O}_X(*D)$-coherent module $\mathscr{M}$, we obtain that $\mathscr{M}$ vanishes on some multi-sectorial neighbourhood of $\widetilde{x}_o$ away from $\varpi^{-1}(D)$. Being $\mathscr{O}_X(*D)$-coherent, it vanishes on some neighbourhood of $x_o$ away from $D$. It is therefore equal to zero.

*Flatness*. — Proving flatness is a matter of proving that, given $f_1, \dots, f_p \in \mathscr{O}_{X,x_o}(*D)$, if $a_1, \dots, a_p \in \mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{mod}\,D}$ (resp. in $\mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{rd}\,D}$) are such that $a_1 f_1 + \cdots + a_p f_p = 0$, then $(a_1, \dots, a_p)$ is a linear combination with coefficients in $\mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{mod}\,D}$ (resp. in $\mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{rd}\,D}$) of relations between $f_1, \dots, f_p$ with coefficients in $\mathscr{O}_{X,x_o}(*D)$. Notice then that it is equivalent to prove flatness over $\mathscr{O}_X$, because any local equation of $D$ is invertible in $\mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{mod}\,D}$ (resp. in $\mathscr{A}_{\widetilde{X},\widetilde{x}_o}^{\mathrm{rd}\,D}$), so we will assume below that $f_1,\dots,f_p\in\mathscr{O}_{X,x_o}$. We argue by induction on $p$, starting with $p=2$.

*Case where* $p=2$. — At this point, we do not need to assume that $\dim X\leqslant 2$. We can then assume that $f_1$ and $f_2$ have no common irreducible component. Let us denote by $g$ a local equation of $D$ at $x_o$. Let $\widetilde{U}$ be an neighbourhood of $\widetilde{x}_o$ in $\widetilde{X}$ such that we have a relation $a_1f_1=a_2f_2$ on $U^*=\widetilde{U}\smallsetminus\partial\widetilde{X}$. By Hartogs, there exists then a holomorphic function $\lambda$ on $U^*$ such that $a_1=\lambda f_2$ and $a_2=\lambda f_1$. We wish to show that $\lambda$ belongs to $\Gamma(\widetilde{U},\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D})$.

Let us choose a proper modification $e:Z\to X$, with $Z$ smooth, such that $f_1\circ e\cdot g\circ e$ defines a divisor $E$ with normal crossing in some neighbourhood of $e^{-1}(x_o)$. It is then enough to show that $\lambda\circ e$ has moderate growth along the real blow-up $\partial\widetilde{Z}$ of the irreducible components of $e^{-1}(D)$ in $Z$, in the neighbourhood of $\widetilde{e}^{-1}(\widetilde{x}_o)$. By compactness of this set, we can work locally near a point $\widetilde{z}_o\in\widetilde{e}^{-1}(\widetilde{x}_o)$. Let us fix local coordinates $z$ at $z_o=\varpi_Z(\widetilde{z}_o)\in Z$ adapted to $E$. We denote by $z'$ the coordinates defining $e^{-1}(D)$ and set $z=(z',z'')$, so that $f_1\circ e$ is the monomial $z'^{m'}z''^{m''}$. The assumption is that, for any compact neighbourhood $K$ of $\widetilde{z}_o$ in $\widetilde{Z}$, there exists a constant $C_K$ and a negative integer $N_K$ such that, on $K^*$, $|\lambda\circ e|\cdot|z'|^{m'}|z''|^{m''}\leqslant C_K|z'|^{N_K}$. Notice that such a $K$ can be chosen as the product of a compact polydisc in the variables $z''$ with a compact multi-sector in the variables $z'$. Up to changing $N_K$, this reduces to $|\lambda\circ e|\cdot|z''|^{m''}\leqslant C_K|z'|^{N_K}$. Fixing $|z''_i|=r''_i>0$ and small enough, and using Cauchy's formula, we obtain $|\lambda\circ e|\leqslant C'_K|z'|^{N_K}$.

*Case where* $p\geqslant 3$. — We argue by induction on $p$. Assume that we have a relation $a_1f_1+\cdots+a_pf_p=0$ as above.

Firstly, we can reduce to the case where $f_1,\dots,f_{p-1}$ do not have a non trivial common factor $\delta$: we deduce a relation $(a_1f'_1+\cdots+a_{p-1}f'_{p-1})\delta+a_pf_p=0$, and by the $p=2$ case, we deduce a relation $a_1f'_1+\cdots+a_{p-1}f'_{p-1}+a'_pf_p$ of the same kind.

In such a case, since $\dim X=2$, we have $\dim V(f_1,\dots,f_{p-1})=0$, so locally $V(f_1,\dots,f_{p-1})=\{x_o\}$, and there is a relation $\sum_{i=1}^{p-1}h_if_i=1$ in $\mathscr{O}_{X,x_o}(*D)$. We deduce the relation $\sum_{i=1}^{p-1}(a_i+a_pf_ph_i)f_i=0$, and the inductive step expresses the vector of coefficients $(a_i+a_pf_ph_i)$ in terms of relations between $f_1,\dots,f_{p-1}$ in $\mathscr{O}_{X,x_o}(*D)$. The conclusion then follows for $f_1,\dots,f_p$, by using the supplementary relation $(h_1f_p)f_1+\cdots+(h_{p-1}f_p)f_{p-1}-f_p=0$. ☐

If $\pi:X\to X'$ is a proper modification which is an isomorphism $X\smallsetminus D\xrightarrow{\sim}X'\smallsetminus E'$, and if $E_i=\sum_j n_{ij}D_j$ for each $i$ ($n_{ij}\in\mathbb{N}$), it induces $\widetilde{\pi}:\widetilde{X}(D_{j\in J})\to\widetilde{X}'(E'_{i\in I})$. (With the first assumption, the inclusion $E\subset D$ is an equality.) Then $\widetilde{\pi}_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D})=\mathscr{A}_{\widetilde{X}'}^{\mathrm{mod}\,E'}$. Similarly, if $\varpi:\widetilde{X}(D_{j\in J})\to X$ is the natural projection, we have $\varpi_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D})=\mathscr{O}_X(*D)$.

***Proposition 8.7.*** — *With the previous assumption, assume moreover that the divisors* $\sum_j D_j$ *and* $\sum_i E'_i$ *are normal crossing divisors. Then* $\boldsymbol{R}\widetilde{\pi}_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}) = \mathscr{A}_{\widetilde{X}'}^{\mathrm{mod}\, E'}$*, that is,* $\boldsymbol{R}^k\widetilde{\pi}_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}) = 0$ *for* $k \geqslant 1$*.*

*Proof.* — Since $\sum_j D_j$ is a normal crossing divisor, we can apply the Dolbeault-Grothendieck theorem on $\widetilde{X}(D)$ (see [**73**, Prop. II.1.1.7]) and get a $c$-soft resolution of $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}$ by the Dolbeault complex of moderate currents $\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\, D,(0,\bullet)}$ on $\widetilde{X}$ (which are $(0,\bullet)$-forms on $\widetilde{X}$ with coefficients in the sheaf $\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\, D}$ of distributions on $X \smallsetminus D$ with moderate growth along $D$). Therefore, $\boldsymbol{R}\widetilde{\pi}_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}) = \widetilde{\pi}_*\,\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\, D,(0,\bullet)}$.

Recall that, on an open set $\widetilde{U}$, $\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\, D}(\widetilde{U})$ is dual to the space of $C^\infty$ functions with compact support in $\widetilde{U}$ which have rapid decay along $\widetilde{U}\cap\partial\widetilde{X}$. Since $\pi$ is a modification, for any compact set $K$ in $\widetilde{U}$, setting $K' = \widetilde{\pi}(K)$, the pull-back of forms $\widetilde{\pi}^*$ identifies $C^\infty$ forms on $\widetilde{X}'$ with support in $K'$ and having rapid decay along $\partial\widetilde{X}'$ with the corresponding forms on $\widetilde{X}$ with support in $K$ and rapid decay along $\partial\widetilde{X}$, and this identification is compatible with the differential, as well as with $\partial$ and $\overline{\partial}$. Dually, the integration along the fibres of $\widetilde{\pi}$ of currents identifies the complexes $\widetilde{\pi}_*\,\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\, D,(0,\bullet)}$ with $\mathfrak{Db}_{\widetilde{X}'}^{\mathrm{mod}\, E',(0,\bullet)}$.

Lastly, the latter complex is a resolution of $\mathscr{A}_{\widetilde{X}'}^{\mathrm{mod}\, E'}$ by Dolbeault-Grothendieck, since $\sum_i E'_i$ is a normal crossing divisor. ☐

***Remark 8.8.*** — Other variants of this proposition can be obtained with a similar proof. For instance, we have $\boldsymbol{R}\varpi_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}) = \varpi_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}) = \mathscr{O}_X(*D)$. More generally, with the assumptions in Proposition 8.7, let $I_1$ and $J_1$ be subsets of $I$ and $J$ respectively such that each $E_i$ $(i \in I_1)$ is expressed as a linear combination with coefficients in $\mathbb{N}$ of $(D_{j\in J_1})$. Let $\widetilde{\pi}_1 : \widetilde{X}(D_{j\in J_1}) \to \widetilde{X}'(E'_{i\in I_1})$ be the morphism induced by $\pi$ between the partial real blow-up spaces. Then, with obvious notation, $\boldsymbol{R}\widetilde{\pi}_*(\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D_1}(*D)) = \mathscr{A}_{\widetilde{X}'}^{\mathrm{mod}\, E'_1}(*E')$. A particular case is $I_1 = \varnothing$, $J_1 = \varnothing$, giving $\boldsymbol{R}\pi_*\mathscr{O}_X(*D) = \mathscr{O}_{X'}(*E')$.

## 8.c. The moderate de Rham complex.

— We keep the setting of §8.b. The sheaf $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D}$ is stable by derivations of $X$ (in local coordinates) and there is a natural de Rham complex on $\widetilde{X}(D_{j\in J})$:

$$\mathrm{DR}^{\mathrm{mod}\, D}(\mathscr{O}_X) := \{\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D} \xrightarrow{d} \mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\, D} \otimes \varpi^{-1}\Omega_X^1 \longrightarrow \cdots\}$$

When restricted to $X^*$, this complex is nothing but the usual holomorphic de Rham complex.

Let $\mathscr{M}$ be a holonomic $\mathscr{D}_X$-module which is localized along $D$, that is, such that $\mathscr{M} = \mathscr{O}_X(*D) \otimes_{\mathscr{O}_X} \mathscr{M}$. In particular, $\mathscr{M}$ is also a coherent $\mathscr{D}_X(*D)$-module. We can also regard $\mathscr{M}$ as a $\mathscr{O}_X(*D)$-module equipped with a flat connection $\nabla$. If moreover $\mathscr{M}$ is $\mathscr{O}_X(*D)$-coherent, we call it a meromorphic connection with poles along $D$ (according to [**52**, Prop. 1.1], it is then locally stably free as a $\mathscr{O}_X(*D)$-module).

We associate with $\mathscr{M}$ the *moderate de Rham complex*

$$\mathrm{DR}^{\bmod D}(\mathscr{M}) := \{\mathscr{A}_{\widetilde{X}}^{\bmod D} \otimes \varpi^{-1}(\mathscr{M}) \xrightarrow{\nabla} \mathscr{A}_{\widetilde{X}}^{\bmod D} \otimes \varpi^{-1}(\mathscr{M} \otimes \Omega_X^1) \to \cdots\}$$

which coincides with $\mathrm{DR}(\mathscr{M})$ on $X^*$.

**Proposition 8.9.** — *Let $\pi : X \to X'$ be a proper morphism between complex manifolds. Assume the following:*

(1) *There exist locally finite families of divisors $(E'_{i\in I})$ of $X'$ and $(D_{j\in J})$ of $X$ such that $E_i := \pi^* E'_i = \sum_j n_{ij} D_j$ with $n_{ij} \in \mathbb{N}$,*

(2) *$\pi : X \smallsetminus D \to X' \smallsetminus E'$ is an isomorphism.*

*Let $\mathscr{M}$ be a holonomic $\mathscr{D}_X$-module which is localized along $D$. If $\dim X \geqslant 3$, assume moreover that $\mathscr{M}$ is smooth on $X \smallsetminus D$, i.e., is a meromorphic connection with poles along $D$ at most. Let $\pi_+\mathscr{M}$ the direct image of $\mathscr{M}$ (as a $\mathscr{D}_{X'}(*E)$-module). Then*

$$\mathrm{DR}^{\bmod E'}(\pi_+\mathscr{M}) \simeq \boldsymbol{R}\widetilde{\pi}_* \, \mathrm{DR}^{\bmod D}(\mathscr{M}).$$

**Remark 8.10.** — We have variants corresponding to those in Remark 8.8.

*Preliminaries on meromorphic connections and proper modifications.* — Let us first recall classical facts concerning direct images of meromorphic connections by a proper modification. The setting is the following. We denote by $\pi : X \to X'$ a proper modification between complex manifolds, and we assume that there are reduced divisors $D \subset X$ and $E' \subset X'$ such that $\pi : X \smallsetminus D \to X' \smallsetminus E'$ is an isomorphism (so that in particular $\pi^{-1}(E') = D$). We do not assume now that $D$ or $E'$ are normal crossing divisors. We denote by $\mathscr{O}_X(*D)$ and $\mathscr{O}_{X'}(*E')$ the corresponding sheaves of meromorphic functions.

**Lemma 8.11.** — *We have $\boldsymbol{R}\pi_*\mathscr{O}_X(*D) = \mathscr{O}_{X'}(*E')$.*

*Proof.* — The statement is similar to that of Proposition 8.7, but will be proved with less assumptions. If $D$ and $E'$ have normal crossings, one can adapt the proof of Proposition 8.7, according to the Dolbeault-Grothendieck lemma using currents with moderate growth. In general, one can argue differently as follows.

Since $\pi$ is proper, one has $\boldsymbol{R}^k\pi_*\mathscr{O}_X(*D) = \varinjlim_j \boldsymbol{R}^k\pi_*\mathscr{O}_X(jD)$. The left-hand term is equal to $\mathscr{O}_{X'}(*E') \otimes_{\mathscr{O}'_X} \boldsymbol{R}^k\pi_*\mathscr{O}_X(*D)$. Then, $\mathscr{O}_{X'}(*E') \otimes_{\mathscr{O}'_X} \varinjlim_j \boldsymbol{R}^k\pi_*\mathscr{O}_X(jD) = \varinjlim_j \mathscr{O}_{X'}(*E') \otimes_{\mathscr{O}'_X} \boldsymbol{R}^k\pi_*\mathscr{O}_X(jD)$ (see e.g. [**23**, p. 10]) and the right-hand term is zero if $k \geqslant 1$, since $\boldsymbol{R}^k\pi_*\mathscr{O}_X(jD)$ is then $\mathscr{O}_{X'}$-coherent and supported on $E'$. $\square$

For any $k \in \mathbb{N}$ we have a natural morphism

$$\Omega^k_{X'}(*E') \longrightarrow \pi_*\big(\mathscr{O}_X(*D) \underset{\pi^{-1}\mathscr{O}_{X'}}{\otimes} \pi^{-1}\Omega^k_{X'}(*E')\big).$$

It follows from the previous lemma that this morphism is an isomorphism. Indeed, on the one hand, the projection formula and the previous lemma give

$$\begin{aligned}\boldsymbol{R}\pi_*\big(\mathscr{O}_X(*D)\overset{\boldsymbol{L}}{\underset{\pi^{-1}\mathscr{O}_{X'}}{\otimes}}\pi^{-1}\Omega^k_{X'}(*E')\big) &\simeq \boldsymbol{R}\pi_*\mathscr{O}_X(*D)\overset{\boldsymbol{L}}{\underset{\pi^{-1}\mathscr{O}_{X'}}{\otimes}}\Omega^k_{X'}(*E')\\ &\simeq \mathscr{O}_{X'}(*E')\overset{\boldsymbol{L}}{\underset{\pi^{-1}\mathscr{O}_{X'}}{\otimes}}\Omega^k_{X'}(*E')\simeq\Omega^k_{X'}(*E').\end{aligned}$$

On the other hand, since $\Omega^k_{X'}$ is $\mathscr{O}_{X'}$-locally free, one can eliminate the '$\boldsymbol{L}$' in the first line above, and conclude

$$\begin{aligned}\Omega^k_{X'}(*E') &\simeq \mathscr{H}^0\boldsymbol{R}\pi_*\big(\mathscr{O}_X(*D)\overset{\boldsymbol{L}}{\underset{\pi^{-1}\mathscr{O}_{X'}}{\otimes}}\pi^{-1}\Omega^k_{X'}(*E')\big)\\ &= \pi_*\big(\mathscr{O}_X(*D)\underset{\pi^{-1}\mathscr{O}_{X'}}{\otimes}\pi^{-1}\Omega^k_{X'}(*E')\big).\end{aligned}$$

The cotangent map $T^*\pi$ is a morphism $\mathscr{O}_X\otimes_{\pi^{-1}\mathscr{O}_{X'}}\pi^{-1}\Omega^1_{X'}\to\Omega^1_X$. It induces an isomorphism $\mathscr{O}_X(*D)\otimes_{\pi^{-1}\mathscr{O}_{X'}}\pi^{-1}\Omega^1_{X'}(*E')\to\Omega^1_X(*D)$. Applying $\pi_*$ and using the previous remark, we get an isomorphism $\pi_*T^*\pi:\Omega^1_{X'}(*E')\xrightarrow{\sim}\pi_*\Omega^1_X(*D)$, and $\boldsymbol{R}^k\pi_*\Omega^1_X(*D)=0$ if $k\geqslant 1$. Since $T^*\pi$ is compatible with differentials, we get a commutative diagram

$$\begin{tikzcd}\mathscr{O}_{X'}(*E') \arrow[r, "d"] \arrow[d, "\wr"'] & \Omega^1_{X'}(*E') \arrow[d, "\wr"', "\pi_*T^*\pi"] \\ \pi_*\mathscr{O}_X(*D) \arrow[r, "\pi_*d"] & \pi_*\Omega^1_X(*D)\end{tikzcd} \tag{8.12}$$

where the upper $d$ is the differential on $X'$ and the lower $d$ is that on $X$. Arguing similarly of all $\Omega^k$ and the corresponding differentials gives an isomorphism of complexes

$$\mathrm{DR}\,\mathscr{O}_{X'}(*E')\xrightarrow{\sim}\pi_*\,\mathrm{DR}\,\mathscr{O}_X(*D)\simeq\boldsymbol{R}\pi_*\,\mathrm{DR}\,\mathscr{O}_X(*D).$$

Let now $\mathscr{M}$ be a holonomic $\mathscr{D}_X$-module which is localized along $D$ and let us also regard it as a $\mathscr{O}_X(*D)$-module with a flat connection $\nabla$.

***Lemma 8.13***

(1) *The direct image $\pi_+\mathscr{M}$ of $\mathscr{M}$ as a $\mathscr{D}_X$-module, once localized along $E'$, has cohomology in degree $0$ at most and this cohomology is a holonomic $\mathscr{D}_{X'}$-module localized along $E'$.*

(2) *We have $\boldsymbol{R}^k\pi_*\mathscr{M}=0$ for $k\geqslant 1$. Moreover, $\pi_*\mathscr{M}$ is equal to the $\mathscr{O}_{X'}(*D')$-module underlying $\pi_+\mathscr{M}(*E')$, and the connection on the latter is equal to the composition of $\pi_*\nabla:\pi_*\mathscr{M}\to\pi_*(\Omega^1_X(*D)\otimes\mathscr{M})$ with the isomorphism*

$$\Omega^1_{X'}(*E')\otimes\pi_*\mathscr{M}\simeq\pi_*(\pi^{-1}\Omega^1_{X'}(*E')\otimes\mathscr{M})\xrightarrow{\sim}\pi_*(\Omega^1_X(*D)\otimes\mathscr{M}) \tag{8.13 $*$}$$

*induced by $\pi_*T^*\pi$.*

*Proof*

(1) Since $\mathscr{M}$ has a coherent $\mathscr{O}_X$-module $\mathscr{N}$ generating $\mathscr{M}$ as a $\mathscr{D}_X$-module (see [**53**], see also [**54**, Th. 3.1]), it is known (see [**31, 48**]) that $\pi_+\mathscr{M}$ has holonomic cohomology. Moreover, $\mathscr{H}^k\pi_+\mathscr{M}$ is supported on $E'$ if $k \geqslant 1$. By flatness of $\mathscr{O}_{X'}(*E')$ over $\mathscr{O}_{X'}$, we have $\mathscr{H}^k(\mathscr{O}_{X'}(*E') \otimes_{\mathscr{O}_{X'}} \pi_+\mathscr{M}) = \mathscr{O}_{X'}(*E') \otimes_{\mathscr{O}_{X'}} \mathscr{H}^k(\pi_+\mathscr{M})$. By a theorem of Kashiwara [**32**, Prop. 2.9], $\mathscr{O}_{X'}(*E') \otimes_{\mathscr{O}_{X'}} \mathscr{H}^k(\pi_+\mathscr{M})$ is a holonomic $\mathscr{D}_{X'}$-module and it is localized along $E'$ by definition. If $k \geqslant 1$, it is thus equal to zero.

(2) The proof is similar to that of Lemma 8.11. We use that $\mathscr{M}$ has a $\mathscr{O}_X(*D)$-generating coherent $\mathscr{O}_X$-submodule: In the case where $\mathscr{M}$ is a meromorphic connection, this follows from [**52**] (see also [**54**]). In the general case, one first uses that $\mathscr{M}$ has a coherent $\mathscr{O}_X$-module $\mathscr{N}$ generating $\mathscr{M}$ as a $\mathscr{D}_X$-module (see [**53**], see also [**54**, Th. 3.1]). Using Bernstein's theory for a function locally defining $D$, one then shows that, locally on $D$, there exists $\ell$ such that $(F_\ell\mathscr{D}_X)\mathscr{N}$ generates $\mathscr{N}$ as a $\mathscr{O}_X(*D)$-module, where $F_\bullet\mathscr{D}_X$ is the filtration by the order. Since $\pi$ is proper, one can find a suitable $\ell$ valid on the inverse image by $\pi$ of any compact set in $X'$. In this way, we get the vanishing of $R^k\pi_*\mathscr{M}$ for $k \geqslant 1$.

Using the isomorphism (8.13 $*$) induced by $\pi_*T^*\pi$, we regard $\pi_*\nabla : \pi_*\mathscr{M} \to \pi_*(\Omega^1_X(*D) \otimes \mathscr{M})$ as a morphism $\pi_+\nabla : \pi_*\mathscr{M} \to \Omega^1_{X'}(*E') \otimes \pi_*\mathscr{M}$, and (8.12) shows that it is a connection on $\pi_*\mathscr{M}$. We denote the resulting object by $\pi_+(\mathscr{M}, \nabla)$. Checking that it is equal to $\mathscr{O}_{X'}(*D') \otimes_{\mathscr{O}_X} \mathscr{H}^0\pi_+\mathscr{M}$ (in the $\mathscr{D}_X$-module sense) is then straightforward. By definition, we have a commutative diagram

$$\begin{tikzcd}
\pi_*\mathscr{M} \arrow[r, "\pi_+\nabla"] \arrow[d, equal] & \Omega^1_{X'}(*E') \otimes \pi_*\mathscr{M} \arrow[d, "\pi_*(T^*\pi \otimes \mathrm{Id})", "\wr"'] \\
\pi_*\mathscr{M} \arrow[r, "\pi_*\nabla"] & \pi_*(\Omega^1_X(*D) \otimes \mathscr{M})
\end{tikzcd} \tag{8.14}$$

□

Let us now go in the other direction. Given a holonomic $\mathscr{D}_{X'}$-module $\mathscr{M}'$ localized along $E'$, the inverse image $\pi^+\mathscr{M}' := \mathscr{D}_{X\to X'} \otimes^{\boldsymbol{L}}_{\pi^{-1}\mathscr{D}_{X'}} \pi^{-1}\mathscr{M}'$ has holonomic cohomology (see [**32**]) and $\mathscr{H}^k\pi^+\mathscr{M}'$ is supported on $D = \pi^{-1}(E')$. It follows that $\mathscr{H}^k(\mathscr{O}_X(*D) \otimes_{\mathscr{O}_X} \pi^+\mathscr{M}') = 0$ if $k \neq 0$ and $\mathscr{H}^0(\mathscr{O}_X(*D) \otimes_{\mathscr{O}_X} \pi^+\mathscr{M}') = \mathscr{O}_X(*D) \otimes_{\mathscr{O}_X} \mathscr{H}^0(\pi^+\mathscr{M}')$ is holonomic and localized along $D$, and its underlying $\mathscr{O}_X(*D)$-submodule is $\pi^*\mathscr{M}' := \mathscr{O}_X(*D) \otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')} \pi^{-1}\mathscr{M}'$. Denoting by $\nabla'$ the connection on $\mathscr{M}'$, denoted by $\pi^+\nabla'$, is defined as $d \otimes \mathrm{Id} + T^*\pi(\mathrm{Id} \otimes \pi^{-1}\nabla')$, where the second term is the composition of

$$\mathrm{Id} \otimes \nabla' : \mathscr{O}_X(*D) \otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')} \pi^{-1}\mathscr{M}' \longrightarrow \mathscr{O}_X(*D) \otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')} \pi^{-1}(\Omega^1_{X'} \otimes \mathscr{M}')$$

with

$$\begin{aligned}T^*\pi\otimes\mathrm{Id} : (\mathscr{O}_X(*D)\otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')}\pi^{-1}\Omega^1_{X'}(*E'))\otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')}\pi^{-1}\mathscr{M}'&\\ \longrightarrow \Omega^1_X(*D)\otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')}\pi^{-1}\mathscr{M}'.&\end{aligned}$$

We set $\pi^+(\mathscr{M}',\nabla')=(\pi^*\mathscr{M},\pi^+\nabla')$, and this is nothing but $\mathscr{O}_X(*D)\otimes_{\mathscr{O}_X}\mathscr{H}^0(\pi^+\mathscr{M}')$ as a holonomic $\mathscr{D}_X$-module localized along $D$.

The natural morphism of $\mathscr{O}_X(*D)$-modules

$$\pi^*\pi_*\mathscr{M} := \mathscr{O}_X(*D)\otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')}\pi^{-1}\pi_*\mathscr{M}\longrightarrow\mathscr{M}$$

induced by the adjunction $\pi^{-1}\pi_*\mathscr{M}\to\mathscr{M}$ is compatible with the connections, so that it induces a morphism

$$\pi^+\pi_+(\mathscr{M},\nabla)\longrightarrow(\mathscr{M},\nabla)$$

which can be regarded as the adjunction morphism at the level of $\mathscr{D}_X(*D)$-modules. It is therefore an isomorphism, since the kernel and cokernel are localized holonomic $\mathscr{D}_X$-modules which are supported on $D$.

Similarly, the adjunction morphism $\mathrm{Id}\to\pi_*\pi^{-1}$ together with the projection formula induces an isomorphism

$$(\mathscr{M}',\nabla')\xrightarrow{\ \sim\ }\pi_+\pi^+(\mathscr{M}',\nabla').$$

In conclusion, the functors $\pi_+$ and $\pi^+$ are quasi-inverse one to the other.

*Proof of Proposition* 8.9. — One can extend the previous results by replacing $\mathscr{O}_X(*D)$ with $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}$ and $\mathscr{O}_{X'}(*E')$ with $\mathscr{A}^{\mathrm{mod}\,E'}_{\widetilde X'}$. In order to do this, we now assume that $D$ and $E'$ have normal crossings. By the first assumption in the proposition, we have a morphism $\widetilde\pi:\widetilde X\to\widetilde X'$ lifting $\pi$. By the preliminaries above, we can write $\mathscr{M}=\mathscr{O}_X(*D)\otimes_{\pi^{-1}\mathscr{O}_{X'}(*E')}\pi^{-1}\mathscr{M}'$. Therefore, $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}\otimes_{\varpi^{-1}\mathscr{O}_X}\mathscr{M}=\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}\otimes_{\widetilde\pi^{-1}\varpi'^{-1}\mathscr{O}_{X'}}\widetilde\pi^{-1}\varpi'^{-1}\mathscr{M}'$. Since $\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}$ (resp. $\mathscr{A}^{\mathrm{mod}\,E'}_{\widetilde X'}$) is flat over $\varpi^{-1}\mathscr{O}_X(*D)$ (resp. $\varpi'^{-1}\mathscr{O}_{X'}(*E')$) ($\dim X\leqslant 2$) or since $\mathscr{M}$ is locally stably free over $\mathscr{O}_X(*D)$ ($\dim X\geqslant 3$), it follows from Proposition 8.7 and the projection formula that

$$\begin{aligned}\boldsymbol{R}\widetilde\pi_*(\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}\otimes_{\varpi^{-1}\mathscr{O}_X(*D)}\mathscr{M})&=\boldsymbol{R}\widetilde\pi_*(\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}\otimes^{\boldsymbol{L}}_{\varpi^{-1}\mathscr{O}_X(*D)}\mathscr{M})\\&=\mathscr{A}^{\mathrm{mod}\,E'}_{\widetilde X'}\otimes^{\boldsymbol{L}}_{\varpi'^{-1}\mathscr{O}_{X'}}\varpi'^{-1}\mathscr{M}'\\&=\mathscr{A}^{\mathrm{mod}\,E'}_{\widetilde X'}\otimes_{\varpi'^{-1}\mathscr{O}_{X'}}\varpi'^{-1}\mathscr{M}',\end{aligned}$$

and therefore the latter term is equal to $\widetilde\pi_*(\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}\otimes_{\varpi^{-1}\mathscr{O}_X}\mathscr{M})$. Arguing similarly after tensoring with $\Omega^k$ gives that each term of the complex $\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$ is $\widetilde\pi_*$-acyclic and that $\widetilde\pi_*\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$ and $\mathrm{DR}^{\mathrm{mod}\,E'}\mathscr{M}'$ are isomorphic termwise. Moreover, the connection $\pi_*\nabla$ on $\widetilde\pi_*(\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde X}\otimes_{\varpi^{-1}\mathscr{O}_X}\mathscr{M})$ coincides, via a diagram similar to (8.14) to the connection $\pi_+\nabla$ on $\mathscr{A}^{\mathrm{mod}\,E'}_{\widetilde{X}'}\otimes_{\varpi'^{-1}\mathscr{O}_{X'}}\varpi'^{-1}\mathscr{M}'$. Extending this isomorphism to the de Rham complexes gives

$$\mathrm{DR}^{\mathrm{mod}\,E'}\mathscr{M}'\xrightarrow{\sim}\widetilde{\pi}_*\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}\simeq\boldsymbol{R}\widetilde{\pi}_*\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}.\qquad\square$$

**8.d. Examples of moderate de Rham complexes.** — We consider the local setting where $(X,0)$ is a germ of complex manifold, $D$ is a divisor with normal crossing in $X$ (defined by $t_1\cdots t_\ell=0$ in some coordinate system $(t_1,\dots,t_n)$ and, setting $L=\{1,\dots,\ell\}$, $\widetilde{X}=\widetilde{X}(D_{i\in L})$, with $D_i=\{t_i=0\}$. We will give examples of computation of moderate de Rham complexes $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)$, with $\varphi\in\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$, and $\mathscr{E}^\varphi:=(\mathscr{O}_X(*D),d+d\varphi)$. In the following, we assume that $\varphi\neq0$ in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$.

***Proposition 8.15.*** — *Assume that there exists $\boldsymbol{m}\in\mathbb{N}^\ell$ such that $\varphi=t^{-\boldsymbol{m}}u(t)\bmod\mathscr{O}_{X,0}$, with $u\in\mathscr{O}_{X,0}$ and $u(0)\neq0$. Then $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)$ has cohomology in degree $0$ at most.*

*Proof.* — This is a direct consequence of theorems in asymptotic analysis due to Majima [**42**]. See a proof in [**25**, Appendix, Th. A.1]. $\square$

On the other hand, if $u(0)=0$, the result can fail, as shown by the following example.

***Example 8.16.*** — *Assume that $X=\mathbb{C}^2$ with coordinates $x,y$, $D=\{y=0\}$ and $\varphi=x^2/y$. Let $\varpi:\widetilde{X}(D)\to X$ be the real blowing-up of $D$ in $X$. Then $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)$ has a nonzero $\mathscr{H}^1$.*

*Proof.* — By Proposition 8.15, $\mathscr{H}^1\,\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)$ is supported on $\varpi^{-1}(0,0)\simeq S^1_y$, since $\varphi$ satisfies the assumption of this proposition away from $x=0$. In order to compute this sheaf, we will use a blowing-up method similar to that used in Lecture 7, in order to reduce to local computations where Proposition 8.15 applies. In the following, we will fix $\theta_o\in S^1_y$ and we will compute the germ $\mathscr{H}^1\,\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)$ at $\theta_o$.

Let $\pi:X'\to X$ be the complex blow-up of the origin in $X$. It is covered by two affine charts, $X'_1$ with coordinates $(x,v)$ with $\pi(x,v)=(x,xv)$ and $X'_2$ with coordinates $(u,y)$ with $\pi(u,y)=(yu,y)$. On $X'_1\cap X'_2$, we have $v=1/u$, and these are the coordinates on the exceptional divisor $E\simeq\mathbb{P}^1$. We still denote by $D$ the strict transform of $D$, which is defined by $v=0$ in $X'_1$ (and does not meet $X'_2$). We have a natural map $\widetilde{\pi}:\widetilde{X}'(D,E)\to\widetilde{X}(D)$, and by Proposition 8.9, we have $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)=\boldsymbol{R}\widetilde{\pi}_*\,\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})$, where $D'=D\cup E=\pi^{-1}(D)$ and $\varphi'=\varphi\circ\pi$. Restricting to $\arg y=\theta_o$ gives, by proper base change,

$$\mathscr{H}^1\,\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^\varphi)_{\theta_o}=\boldsymbol{H}^1\big(\widetilde{\pi}^{-1}(\theta_o),\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})\big).$$

*Chart* $X_2'$. — We have $\widetilde{X}_2' = \widetilde{X}_2'(E)$ and, on $X_2'$, $\varphi' = yu^2 = 0 \bmod \mathscr{O}_{X_2'}$. Therefore, according to Proposition 8.15, $\mathrm{DR}^{\mathrm{mod}\,E}(\mathscr{E}^{\varphi'})$ has cohomology in degree 0 only on $\widetilde{X}_2'$, and $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,E}(\mathscr{E}^{\varphi'})$ is the constant sheaf $\mathbb{C}$. Let us note that $\partial\widetilde{X}_2' = \mathbb{A}^1 \times S^1_y$, so the restriction to $\arg y = \theta_o$ of $\mathrm{DR}^{\mathrm{mod}\,E}(\mathscr{E}^{\varphi'})$ is the constant sheaf $\mathbb{C}_{\mathbb{A}^1}$.

*Chart* $X_1'$. — We have $\widetilde{X}_1' = \widetilde{X}_1'(D,E)$ and $\partial\widetilde{X}'_{1|E} = S^1_x \times S^1_v \times [0,\infty)$, where $|v|$ runs in $[0,\infty)$. The map $\widetilde{\pi} : \partial\widetilde{X}'_{1|E} \to S^1_y$ is the composed map

$$\begin{aligned} S^1_x \times S^1_v \times [0,\infty) \longrightarrow S^1_x \times S^1_v &\longrightarrow S^1_y \\ (\alpha,\beta) &\longmapsto \alpha\beta, \end{aligned}$$

so, in this chart, $\widetilde{\pi}^{-1}(\theta_o) \simeq S^1 \times [0,\infty)$ and we can assume $S^1 = S^1_v$ so that we can identify $\widetilde{\pi}^{-1}(\theta_o)$ in $\partial\widetilde{X}'$ to a closed disc having $S^1_v$ as boundary. We also have $\varphi' = x/v$.

**Claim.** — *The complex* $\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})$ *has cohomology in degree* 0 *at most on* $\widetilde{X}_1'(D,E)$.

Assuming this claim is proved, it is not difficult to compute $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})_{\theta_o}$. In the chart $X_2'$, this has been done previously, so we are reduced to compute it on $S^1_v$ (boundary of the closed disc $\widetilde{\pi}^{-1}(\theta_o)$). The $\mathscr{H}^0$ is zero unless $\arg x - \arg v = \theta_o - 2\arg v \in (\pi/2, 3\pi/2) \bmod 2\pi$.

*Conclusion.* — The complex $\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})_{\theta_o}$ on $\widetilde{\pi}^{-1}(\theta_o)$ has cohomology in degree 0 only, and is the constant sheaf on $\Delta \cup I_1 \cup I_2$, extended by 0 to $\overline{\Delta}$, where $\Delta$ is an open disc in $\mathbb{C}$ and $I_1$ and $I_2$ are two opposite (and disjoint) open intervals of length $\pi/2$ on $\partial\Delta$. It is now an exercise to show that $\dim \boldsymbol{H}^1\big(\widetilde{\pi}^{-1}(\theta_o), \mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})\big) = 1$. ☐

*Proof of the claim.* — Let us sketch it. The question is local in the $(x,v)$-chart, on $S^1_x \times S^1_v$. We blow up the origin in this chart $X_1'$ and get an exceptional divisor $E' \simeq \mathbb{P}^1$ with coordinates $v_1 = 1/u_1$. On the blown-up space $X_1''$, in the chart $X_{11}''$ with coordinates $(x, v_1)$, the blowing-up map $\pi'$ is $(x,v_1) \mapsto (x, v = xv_1)$ and we have $\varphi'' = x/v = 1/v_1$. In the chart $X_{12}''$ with coordinates $(u_1, v)$, the blowing-up map $\pi'$ is $(u_1,v) \mapsto (x = u_1v, v)$ and we have $\varphi'' = x/v = u_1$.

On $\widetilde{X}''_{12|E'}$, $\widetilde{\pi}'$ is the composed map

$$\begin{aligned} S^1_{u_1} \times S^1_v \times [0,\infty) \longrightarrow S^1_{u_1} \times S^1_v &\longrightarrow S^1_x \times S^1_v \\ (\alpha,\beta) &\longmapsto (\alpha\beta,\beta), \end{aligned}$$

where $|u_1|$ varies in $[0,\infty)$. On this space, $\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi''})$ is the constant sheaf $\mathbb{C}$.

Similarly, on $\widetilde{X}''_{11|E'}$, $\widetilde{\pi}'$ is the composed map

$$\begin{aligned} S^1_x \times S^1_{v_1} \times [0,\infty) \longrightarrow S^1_x \times S^1_{v_1} &\longrightarrow S^1_x \times S^1_v \\ (\alpha,\gamma) &\longmapsto (\alpha,\alpha\gamma), \end{aligned}$$

and $|v_1|$ varies in $[0,\infty)$. On this space, $\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi''})$ has cohomology in degree 0 at most, after Proposition 8.15, and is the constant sheaf $\mathbb{C}$, except on $|v_1|=0$, $\arg v_1 \notin (\pi/2, 3\pi/2) \bmod 2\pi$, where it is zero.

Let us fix a point $(\alpha^o,\beta^o)\in S^1_x\times S^1_v$. We have $\widetilde{\pi}'^{-1}(\alpha^o,\beta^o)\simeq[0,+\infty]$, and $\mathscr{H}^k\,\mathrm{DR}^{\mathrm{mod}\,D'}(\mathscr{E}^{\varphi'})_{(\alpha^o,\beta^o)}=H^k([0,+\infty],\mathscr{F}_{(\alpha^o,\beta^o)})$, where $\mathscr{F}_{(\alpha^o,\beta^o)}$ is the constant sheaf $\mathbb{C}$ if $\gamma^o:=\beta^o\alpha^{o-1}\notin(\pi/2,3\pi/2)\bmod 2\pi$, and the constant sheaf on $(0,+\infty]$ extended by 0 at 0 otherwise.

In the first case, we have $H^k=0$ for any $k\geqslant 1$, and in the second case we have $H^k=0$ for any $k\geqslant 0$. $\square$

***Remark 8.17***. — The previous proof was adapted for using Proposition 8.9. One can use instead similar arguments for finite mappings by considering the map $(x,y)\mapsto(z=x^2,y)$. The direct image of $\mathscr{E}^{x^2/y}$ (as a $\mathscr{D}$-module) by this map is an extension of the regular holonomic $\mathscr{D}$-module $\mathbb{C}[y,y^{-1}]\delta(z)$ by $\mathscr{E}^{z/y}$. The latter is responsible for the non-vanishing of $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{E}^{x^2/y})$.

# LECTURE 9

# STOKES-FILTERED LOCAL SYSTEMS ALONG A DIVISOR WITH NORMAL CROSSINGS

**Summary.** We construct the sheaf $\mathcal{I}$ to be considered as the index sheaf for Stokes filtrations. This is a sheaf on the real blow-up space of a complex manifold along a family of divisors. We will consider only divisors with normal crossings. The global construction of $\mathcal{I}$ needs some care, as the trick of considering a ramified covering cannot be used globally. The important new notion is that of goodness. It is needed to prove abelianity and strictness in this setting, generalizing the results of Lecture 3.

**9.a. The sheaf $\mathcal{I}$ on the real blow-up (smooth divisor case).** — Let $X$ be a smooth complex manifold and let $D$ be a smooth divisor in $X$. Let $\varpi : \widetilde{X}(D) \to X$ be "the" real blow-up space of $X$ along $D$ (associated with the choice of a section $f : \mathscr{O}_X \to \mathscr{O}_X(D)$ of $L(D)$ defining $D$, see §8.a). We also denote by $\widetilde{\imath}, \widetilde{\jmath}$ the inclusions $\partial\widetilde{X}(D) \hookrightarrow \widetilde{X}(D)$ and $X^* \hookrightarrow \widetilde{X}(D)$.

We consider the sheaf $\widetilde{\jmath}_*\mathscr{O}_{X^*}$ and its subsheaf $(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$ of locally bounded functions on $\widetilde{X}$. We will construct $\mathcal{I}$ as a subsheaf of the quotient sheaf $\widetilde{\jmath}_*\mathscr{O}_{X^*}/(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$ (which is supported on $\partial\widetilde{X}$). It is the union, over $d \in \mathbb{N}^*$, of the subsheaves $\mathcal{I}_d$ that we define below.

Let us start with $\mathcal{I}_1$. There is a natural inclusion $\varpi^{-1}\mathscr{O}_X(*D) \hookrightarrow \widetilde{\jmath}_*\mathscr{O}_{X^*}$, and $\varpi^{-1}\mathscr{O}_X = \varpi^{-1}\mathscr{O}_X(*D) \cap (\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$, since a meromorphic function which is bounded in some sector centered on an open set of $D$ is holomorphic. We then set $\widetilde{\mathcal{I}}_1 = \varpi^{-1}\mathscr{O}_X(*D) \subset \widetilde{\jmath}_*\mathscr{O}_{X^*}$ and $\mathcal{I}_1 = \varpi^{-1}(\mathscr{O}_X(*D)/\mathscr{O}_X) \subset \widetilde{\jmath}_*\mathscr{O}_{X^*}/(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$.

Locally on $D$, we can define ramified coverings $\rho_d : X_d \to X$ of order $d$ along $D$, for any $d$. Let $\widetilde{\rho}_d : \widetilde{X}_d \to \widetilde{X}$ be the corresponding covering. The subsheaf $\widetilde{\mathcal{I}}_d \subset \widetilde{\jmath}_*\mathscr{O}_{X^*}$ of $d$-multivalued meromorphic functions on $\widetilde{X}$ is defined as the intersection of the subsheaves $\widetilde{\jmath}_*\mathscr{O}_{X^*}$ and $\widetilde{\rho}_{d,*}\widetilde{\mathcal{I}}_{\widetilde{X}_d,1}$ of $\widetilde{\rho}_{d,*}\widetilde{\jmath}_{d,*}\mathscr{O}_{X_d^*} = \widetilde{\jmath}_*\rho_{d,*}\mathscr{O}_{X_d^*}$. As above, we have $\widetilde{\mathcal{I}}_d \cap (\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}} = \widetilde{\jmath}_*\mathscr{O}_{X^*} \cap \widetilde{\rho}_{d,*}\varpi_d^{-1}\mathscr{O}_{X_d}$. The sheaf $\mathcal{I}_d$ is then defined as the quotient sheaf $\widetilde{\mathcal{I}}_d/\widetilde{\mathcal{I}}_d \cap (\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$. This is a subsheaf of $\widetilde{\jmath}_*\mathscr{O}_{X^*}/(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$.

The locally defined subsheaves $\widetilde{\mathcal{I}}_d$ (and thus $\mathcal{I}_d$) glue together as a subsheaf of $\widetilde{\jmath}_*\mathscr{O}_{X^*}$, since the local definition does not depend on the chosen local ramified $d$-covering. Similarly, $\mathcal{I}_d$ exists as a subsheaf of $\widetilde{\jmath}_*\mathscr{O}_{X^*}/(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$ all over $D$.

***Definition 9.1* (Case of a smooth divisor).** — The sheaf $\widetilde{\mathcal{I}}$ (resp. $\mathcal{I}$) is the union of the subsheaves $\widetilde{\mathcal{I}}_d$ (resp. $\mathcal{I}_d$) of $\widetilde{\jmath}_*\mathscr{O}_{X^*}$ (resp. $\widetilde{\jmath}_*\mathscr{O}_{X^*}/(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$) for $d\in\mathbb{N}^*$.

***Definition 9.2* ($\mathcal{I}$ as a sheaf of ordered abelian groups).** — The sheaf $\widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{O}_{X^*}$ is naturally ordered by setting $(\widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{O}_{X^*})_{\leqslant 0}=\log\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}(D)}$ (see §8.c). In this way, $\widetilde{\mathcal{I}}$ inherits an order: $\widetilde{\mathcal{I}}_{\leqslant 0}=\widetilde{\mathcal{I}}\cap\log\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}(D)}$. This order is not altered by adding a local section of $(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$, and thus defines an order on $\mathcal{I}$.

***Lemma 9.3*.** — *For any local ramified covering $\rho_d:(X_d,D)\to(X,D)$ of order $d$ along $D$, $\widetilde{\rho}_d^{-1}\widetilde{\mathcal{I}}_d$ is identified with $\varpi_d^{-1}\mathscr{O}_{X_d}(*D)$ and $\widetilde{\rho}_d^{-1}\mathcal{I}_d$ with $\varpi_d^{-1}(\mathscr{O}_{X_d}(*D)/\mathscr{O}_{X_d})$. This identification is compatible with order.*

*Proof*. — The proof is completely similar to that given in Remark 2.23. ☐

### 9.b. The sheaf $\mathcal{I}$ on the real blow-up (normal crossing case)

Let us now consider a family $(D_{j\in J})$ of smooth divisors of $X$ whose union $D$ has only normal crossings, and the corresponding real blow-up map $\varpi:\widetilde{X}(D_{j\in J})\to X$. We will consider multi-integers $\boldsymbol{d}\in(\mathbb{N}^*)^J$. The definition of the sheaves $\widetilde{\mathcal{I}}_{\boldsymbol{d}}$ and $\mathcal{I}_{\boldsymbol{d}}$ is similar to that in dimension one.

Let us set $\mathbf{1}=(1,\dots,1)$ ($\#J$ terms) and $\widetilde{\mathcal{I}}_{\mathbf{1}}=\varpi^{-1}\mathscr{O}_X(*D)\subset\widetilde{\jmath}_*\mathscr{O}_{X^*}$. Fixing $x_o\in D$, let us denote by $D_1,\dots,D_\ell$ the components of $D$ going through $x_o$, and let $\widetilde{x}_o\in\varpi^{-1}(x_o)\simeq(S^1)^\ell$. Then a local section of $\varpi^{-1}\mathscr{O}_X(*D)$ near $\widetilde{x}_o$ is locally bounded in the neighbourhood of $\widetilde{x}_o$ if and only if it is holomorphic in the neighbourhood of $x_o$. In other words, as in the smooth case, $\varpi^{-1}\mathscr{O}_X(*D)\cap(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}=\varpi^{-1}(\mathscr{O}_X)$.

We locally define $\widetilde{\mathcal{I}}_{\boldsymbol{d}}$ near $x_o$, by using a ramified covering $\rho_{\boldsymbol{d}}$ of $(X,x_o)$ along $(D,x_o)$ of order $\boldsymbol{d}$, by the formula $\widetilde{\mathcal{I}}_{\boldsymbol{d}}:=\widetilde{\rho}_{\boldsymbol{d},*}[\varpi_{\boldsymbol{d},*}\mathscr{O}_{X_{\boldsymbol{d}}}(*D)]\cap\widetilde{\jmath}_*\mathscr{O}_{X^*}$, and $\mathcal{I}_{\boldsymbol{d}}$ by $\mathcal{I}_{\boldsymbol{d}}:=\widetilde{\mathcal{I}}_{\boldsymbol{d}}/\widetilde{\mathcal{I}}_{\boldsymbol{d}}\cap(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$.

The locally defined subsheaves $\widetilde{\mathcal{I}}_{\boldsymbol{d}}$ glue together all over $D$ as a subsheaf $\widetilde{\mathcal{I}}_{\boldsymbol{d}}$ of $\widetilde{\jmath}_*\mathscr{O}_{X^*}$. We also set globally $\mathcal{I}_{\boldsymbol{d}}=\widetilde{\mathcal{I}}_{\boldsymbol{d}}/\widetilde{\mathcal{I}}_{\boldsymbol{d}}\cap(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$.

***Definition 9.4*.** — The subsheaf $\widetilde{\mathcal{I}}\subset\widetilde{\jmath}_*\mathscr{O}_{X^*}$ is the union of the subsheaves $\widetilde{\mathcal{I}}_{\boldsymbol{d}}$ for $\boldsymbol{d}\in(\mathbb{N}^*)^J$. The sheaf $\mathcal{I}$ is the subsheaf $\widetilde{\mathcal{I}}/\widetilde{\mathcal{I}}\cap(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$ of $\widetilde{\jmath}_*\mathscr{O}_{X^*}/(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$.

***Definition 9.5*.** — The order on $\widetilde{\mathcal{I}}$ is given by $\widetilde{\mathcal{I}}_{\leqslant 0}:=\widetilde{\mathcal{I}}\cap\log\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}(D_{j\in J})}$. It is stable by the addition of an element of $(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$ and defines an order on $\mathcal{I}$.

***Lemma 9.6*.** — *For any local ramified covering $\rho_{\boldsymbol{d}}:(X_{\boldsymbol{d}},D)\to(X,D)$ of order $\boldsymbol{d}$ along $(D_{j\in J})$, $\widetilde{\rho}_{\boldsymbol{d}}^{-1}\widetilde{\mathcal{I}}_{\boldsymbol{d}}$ is identified with $\varpi_{\boldsymbol{d}}^{-1}\mathscr{O}_{X_{\boldsymbol{d}}}(*D)$ and $\widetilde{\rho}_{\boldsymbol{d}}^{-1}\mathcal{I}_{\boldsymbol{d}}$ with $\varpi_{\boldsymbol{d}}^{-1}\big(\mathscr{O}_{X_{\boldsymbol{d}}}(*D)/\mathscr{O}_{X_{\boldsymbol{d}}}\big)$. These identifications are compatible with order.*

*Proof.* — Same proof as in Remark 2.23. □

***Remark 9.7.*** — For any subset $I \subset J$, let $D_I$ denote the intersection $\bigcap_{j\in I} D_j$ and set $D_I^\circ = D_I \smallsetminus \bigcup_{j\in J\smallsetminus I} D_j$. Set also $Y_I = \varpi^{-1}(D_I^\circ) \subset \partial\widetilde{X}(D_{j\in J})$. The family $\mathscr{Y} = (Y_I)_{I\subset J}$ is a stratification of $\partial\widetilde{X}$ which satisfies the property (1.44). Moreover, the sheaf $\mathcal{I}$ is Hausdorff with respect to $\mathscr{Y}$ (this is seen easily locally on $D$).

**9.c. Goodness.** — The order on sections of $\mathcal{I}$ is best understood for *purely monomial sections* of $\mathcal{I}$. Let us use the following local notation. We consider the case where $X = \Delta^\ell\times\Delta^{n-\ell}$ with base point $0 = (0_\ell, 0_{n-\ell})$, and $D_i = \{t_i = 0\}$ $(i \in L := \{1,\dots,\ell\})$ and $D = \bigcup_{i=1}^\ell D_i$. The real blow-up map $\varpi_L : \widetilde{X}(D_{i\in L}) = (S^1)^\ell \times [0,1)^\ell \times \Delta^{n-\ell} \to X = \Delta^\ell \times \Delta^{n-\ell}$ is defined by sending $(e^{i\theta_j}, \rho_j)$ to $t_j = \rho_j e^{i\theta_j}$ $(j = 1,\dots,\ell)$.

- In the non-ramified case (i.e., we consider sections of $\mathcal{I}_{\boldsymbol{1}}$), a germ $\eta$ at $\theta \in (S^1)^\ell = (S^1)^\ell \times 0_\ell \times 0_{n-\ell} \subset (S^1)^\ell \times [0,1)^\ell \times \Delta^{n-\ell}$ of section of $\mathcal{I}_{\boldsymbol{1}}$ is nothing but a germ at $0 \in X$ of section of $\mathscr{O}_X(*D)/\mathscr{O}_X$.

***Definition 9.8.*** — We say that $\eta$ is *purely monomial* at 0 if $\eta = 0$ or $\eta$ is the class of $t^{-\boldsymbol{m}}u_{\boldsymbol{m}}$, with $\boldsymbol{m} \in \mathbb{N}^\ell \smallsetminus \{0\}$, $u_{\boldsymbol{m}} \in \mathscr{O}_{X,0}$ and $u_{\boldsymbol{m}}(0) \neq 0$. We then set $\boldsymbol{m} = \boldsymbol{m}(\eta)$ with the convention that $\boldsymbol{m}(0) = 0$, so that $\boldsymbol{m}(\eta) = 0$ iff $\eta = 0$.

For every $\theta \in (S^1)^\ell$, we have:

$$\eta \leqslant_\theta 0 \iff \eta = 0 \text{ or } \arg u_{\boldsymbol{m}}(0) - \textstyle\sum_j m_j\theta_j \in (\pi/2, 3\pi/2) \mod 2\pi. \tag{9.9}$$

If $\eta \neq 0$ and $\eta$ is purely monomial at 0, it is purely monomial on some open set $Y = (S^1)^\ell \times V$ (with $V$ an open neighbourhood of $0_{n-\ell}$ in $\Delta^{n-\ell}$, embedded as $0_\ell\times V \subset [0,1)^\ell\times\Delta^{n-\ell}$), and $Y_{\eta\leqslant 0}$ is defined by the inequation $\arg u_{\boldsymbol{m}}(v) - \sum_j m_j\theta_j \in (\pi/2, 3\pi/2) \bmod 2\pi$, where $v$ varies in the parameter space $V$. For $v$ fixed, it is the inverse image by the fibration map $(S^1)^n \to S^1$, $(e^{i\theta_1},\dots,e^{i\theta_n}) \mapsto e^{i(m_1\theta_1+\cdots+m_n\theta_n)}$, of a set of the kind defined at the end of Example 1.6. This set rotates smoothly when $v$ varies in $V$. Similarly, the boundary $\mathrm{St}(\eta, 0)$ of $Y_{\eta\leqslant 0}$ in $Y$ is the pull-back by the previous map of a subset of $S^1 \times V$ which is a finite covering of $V$. It has codimension one in $Y$.

- The order on $\mathcal{I}_{\boldsymbol{d}}$ is described similarly after a ramification which is cyclic of order $d_i$ around the component $D_i$ of $D$.

For non-purely monomial elements, we still have the following (e.g. in the non-ramified case).

***Proposition 9.10.*** — *For every $\varphi, \psi \in \Gamma(U, \mathscr{O}_X(*D)/\mathscr{O}_X)$, the set $Y_{\psi\leqslant\varphi}$ is subanalytic in $Y$ and its boundary $\mathrm{St}(\varphi,\psi)$ has (real) codimension $\geqslant 1$ in $Y$.*

***Lemma 9.11.*** — *Let $\eta \in \Gamma(U, \mathscr{O}_X(*D)/\mathscr{O}_X)$ and let $x \in D$. Then there exists a projective modification $\varepsilon : U' \to U$ of some open neighbourhood $U$ of $x$ in $X$, which* *is an isomorphism away from $D$, such that $D' := |\varepsilon^{-1}(D)|$ is a reduced divisor with normal crossings and smooth components, and $\eta\circ\varepsilon$ is locally purely monomial on $D'$.*

*Proof of Proposition* 9.10. — Set $\eta = \psi - \varphi = t^{-m}u_m$ with $m \in \mathbb{N}^\ell \smallsetminus \{0\}$ and $u_m$ holomorphic. We have $Y_{\psi\leqslant\varphi} = Y_{\eta\leqslant 0}$. The purely monomial case ($u_m \neq 0$ on $U$) has been treated above. The statement is local subanalytic on $U$, and for any point of $U$ we replace $U$ by a subanalytic open neighbourhood of this point, that we still denote by $X$, on which Lemma 9.11 applies. If we set $D' = \bigcup_j D'_j$, we have natural real analytic proper maps (see §8.a) $\widetilde{X}'(D'_{j\in J}) \to \widetilde{X}$. The set $Y \smallsetminus Y_{\eta\leqslant 0}$ is the push-forward by this map of the set $Y' \smallsetminus Y'_{\eta\circ\varepsilon\leqslant 0}$, where $Y' \subset \widetilde{X}'$ is the inverse image of $Y$. Using the purely monomial case considered previously, one shows that $Y' \smallsetminus Y'_{\eta\circ\varepsilon\leqslant 0}$ is closed and semi-analytic in $Y'$. The subanalyticity of $Y_{\eta\leqslant 0}$ follows then from Hironaka's theorem [**28**] on the proper images of sub- (or semi-) analytic sets, and stability by complements and closure. The statement for $\mathrm{St}(\varphi,\psi)$ also follows. ☐

*Sketch of the proof of Lemma* 9.11. — By using the resolution of singularities in the neighbourhood of $x \in U$, we can find a projective modification $\varepsilon_1 : U_1 \to U$ such that the union of the divisors of zeros and and of the poles of $\eta$ form a divisor with normal crossings and smooth components in $U_1$ (so that, locally in $U_1$ and with suitable coordinates, $\eta\circ\varepsilon_1$ takes the form of a monomial with exponents in $\mathbb{Z}$). The problem is now reduced to the following question: given a divisor with normal crossings and smooth components $D_1$ in $U_1$, attach to each smooth component an integer (the order of the zero or minus the order of the pole of $\eta\circ\varepsilon_1$), so as to write $D_1 = D_1^+ \cup D_1^0 \cup D_1^-$ with respect to the sign of the integer; to any projective modification $\varepsilon_2 : U_2 \to U_1$ such that $D_2 := \varepsilon_2^{-1}(D_1)$ remains a divisor with normal crossings and smooth components, one can associate in a natural way a similar decomposition; we then look for the existence of such an $\varepsilon_2$ such that $D_2^-$ and $D_2^+$ do not intersect.

We now denote $U_1$ by $X$ and $D_1$ by $D = \bigcup_{j\in J} D_j$. The divisor $D$ is naturally stratified, and we consider minimal (that is, closed) strata. To each such stratum is attached a subset $L$ of $J$ consisting of indices $j$ for which $D_j$ contains the stratum. Because of the normal crossing condition, the cardinal of this subset is equal to the codimension of the stratum $D_L$. We set $D(L) = \bigcup_{j\in L} D_j$. We will construct the modification corresponding to this stratum with a toric argument. Let us set $\ell = \#L$ and, for each $j \in L$, let us denote by $\mathscr{I}_j$ the ideal of $D_j$ in $\mathscr{O}_X$.

As is usual in toric geometry (see [**14, 67, 21**] for instance), we consider the space $\mathbb{R}^\ell$ equipped with its natural lattice $N := \mathbb{Z}^\ell$ and the dual space $(\mathbb{R}^\vee)^\ell$ equipped with the dual lattice $M$. To each rational cone $\sigma$ in the first octant $(\mathbb{R}_+)^\ell$ we consider the dual cone $\sigma^\vee \in (\mathbb{R}^\vee)^\ell$ and its intersection with $M$. This allows us to define a sheaf of subalgebras $\sum_{m\in\sigma^\vee\cap M} \mathscr{I}_1^{m_1}\cdots\mathscr{I}_\ell^{m_\ell}$ of $\mathscr{O}_X(*D(L))$. Locally on $D_L$, if $D_j$ is defined by $\{x_j = 0\}$, this is $\mathscr{O}_X \otimes_{\mathbb{C}[x_1,\dots,x_\ell]} \mathbb{C}[\sigma^\vee \cap M]$, hence this sheaf of subalgebras corresponds to an affine morphism $X_\sigma \to X$. Similarly, to any a fan $\Sigma$ in the first octant $(\mathbb{R}_+)^\ell$ one associates a morphism $X_\Sigma \to X$, which is a projective modification if the fan completely subdivides the octant. Moreover, if each cone of the fan is strictly simplicial, the space $X_\Sigma$ is smooth and the pull-back of the divisor $D(L)$ has normal crossings with smooth components.

We will now choose the fan $\Sigma$. To each basis vector $e_j$ of $\mathbb{R}^\ell$ is attached a multiplicity $\nu_j \in \mathbb{Z}$, namely the order of $\eta \circ \varepsilon_1$ along $D_j$. We consider the trace $H$ on $(\mathbb{R}_+)^\ell$ of the hyperplane $\{(n_1, \dots, n_\ell) \in \mathbb{R}^\ell \mid \sum_j \nu_j n_j = 0\}$ and we choose a strictly simplicial fan in $(\mathbb{R}_+)^\ell$ such that $H$ is a union of cones of this fan. For each basis vector $(n_1, \dots, n_\ell) \in \mathbb{N}^\ell$ of a ray (dimension-one cone) of this fan, the multiplicity of the pull-back of $\eta \circ \varepsilon_1$ along the divisor corresponding to this ray is given by $\sum_j \nu_j n_j$. Therefore, for each $\ell$-dimensional cone of the fan, the multiplicities at the rays all have the same sign (or are zero).

The proof of the Lemma now proceeds by decreasing induction on the maximal codimension $\ell$ of closed strata of $D$ which are contained both in $D_+$ and $D_-$. In the space $\mathbb{R}^J$ we consider the various subspaces $\mathbb{R}^L$ $(L \subset J)$ corresponding to these closed strata of $D$. We subdivide each octant $(\mathbb{R}_+)^L$ by a strict simplical fan as above. We also assume that the fans coincide on the common faces of distinct subspaces $\mathbb{R}^L$. We denote by $\Sigma$ the fan we obtain in this way. In order to obtain such a $\Sigma$, one can construct a strict simplicial fan completely subdividing $(\mathbb{R}_+)^J$ which is compatible with the hyperplane $\sum_{j\in J} \nu_j n_j = 0$ and with the various octants $(\mathbb{R}_+)^L$ corresponding to codimension $\ell$ strata of $D$ contained in $D_+ \cap D_-$, and then restrict it to the union of these octants $(\mathbb{R}_+)^L$. We also consider the sheaves $\sum_{m\in\sigma^\vee\cap M_J} \prod_{j\in J} \mathscr{I}_j^{m_j} \subset \mathscr{O}_X(*D)$ for $\sigma \in \Sigma$, and get a projective modification $\varepsilon_\Sigma : X_\Sigma \to X$. By construction, the maximal codimension of closed strata of $\varepsilon_\Sigma^{-1}(D)$ contained in $\varepsilon_\Sigma^{-1}(D)_+ \cap \varepsilon_\Sigma^{-1}(D)_-$ is $\leqslant \ell - 1$. □

For a finite set $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$, the notion of pure monomiality is replaced by *goodness* (see also Remark 11.5(4) below).

***Definition 9.12* (Goodness).** — We say that a finite subset $\Phi$ of $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ is *good* if $\#\Phi = 1$ or for any $\varphi \neq \psi$ in $\Phi$, $\varphi - \psi$ is purely monomial, that is, the divisor of $\varphi - \psi$ is supported on $D = \{t_1 \cdots t_\ell = 0\}$ and is $\leqslant 0$, i.e., there exists $\boldsymbol{m} = \boldsymbol{m}(\varphi - \psi) \in \mathbb{N}^\ell \setminus \{0\}$ such that $\varphi - \psi = t^{-\boldsymbol{m}} u(t)$ with $u(0) \neq 0$.

***Remark 9.13*.** — Let us give some immediate properties of local goodness (see [**73**, I.2.1.4]).

(1) Any subset of a good set is good, and any subset consisting of one element (possibly not purely monomial) is good. If $\Phi$ is good, then its pull-back $\Phi_{\boldsymbol{d}}$ by the ramification $X_{\boldsymbol{d}} \to X$ is good for any $\boldsymbol{d}$. Conversely, if $\Phi_{\boldsymbol{d}}$ is good for some $\boldsymbol{d}$, then $\Phi$ is good.

(2) More generally, for $f : X' \to X$ as in §9.e, if $\Phi \subset \mathscr{O}_{X,x_o}(*D)/\mathscr{O}_{X,x_o}$ is good, then for any $x'_o \in f^{-1}(x_o)$, the subset $(f^*\Phi)_{x'_o} \subset \mathscr{O}_{X',x'_o}(*D')/\mathscr{O}_{X',x'_o}$ is good.

(3) Any germ $\varphi_0$ of $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ defines in a unique way a germ $\varphi_x \in \mathscr{O}_{X,x}(*D)/\mathscr{O}_{X,x}$ for $x \in D \cap U$, $U$ some open neighbourhood of 0. Indeed, choose a lifting $\varphi_0^*$ in $\mathscr{O}_{X,0}(*D)$. It defines in a unique way a section $\varphi^*$ of $\Gamma(U, \mathscr{O}_U(*D))$ for $U$ small enough. Its germ at $x \in U \cap D$ is denoted $\varphi_x^*$. Its image in $\mathscr{O}_{X,x}(*D)/\mathscr{O}_{X,x}$ is $\varphi_x$. Given two liftings $\varphi_0^*$ and $\varphi_0^\star$, their difference is in $\mathscr{O}_{X,0}$. Choose $U$ so that $\varphi^*$ and $(\varphi^* - \varphi^\star)$ are respectively sections of $\Gamma(U, \mathscr{O}_U(*D))$ and $\Gamma(U, \mathscr{O}_U)$. Then these two liftings give the same $\varphi_x$.

Similarly, any finite subset $\Phi$ of $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ defines in a unique way a finite subset, still denoted by $\Phi$, of $\mathscr{O}_{X,x}(*D)/\mathscr{O}_{X,x}$ for any $x$ close enough to 0.

Then, if $\Phi$ is is good at 0, it is good at any point of $D$ in some open neighbourhood of 0. Note however that a difference $\varphi - \psi$ which is non-zero at 0 can be zero along some components of $D$, a phenomenon which causes $\mathcal{I}^{\text{ét}}$ to be non-Hausdorff.

(4) A subset $\Phi$ is good at 0 if and only if for some (or any) $\eta \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ the translated subset $\Phi + \eta$ of it is good.

(5) For a good set $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$, and for any *fixed* $\varphi_o \in \Phi$, the subset $\{\boldsymbol{m}(\varphi - \varphi_o) \mid \varphi \in \Phi\} \subset \mathbb{N}^\ell$ is totally ordered. Its maximum does not depend on the choice of $\varphi_o \in \Phi$, it is denoted by $\boldsymbol{m}(\Phi)$ and belongs to $\mathbb{N}^\ell$. We have $\boldsymbol{m}(\Phi) = 0$ iff $\#\Phi = 1$. We have $\boldsymbol{m}(\Phi + \eta) = \boldsymbol{m}(\Phi)$ for any $\eta \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$.

(6) Assume $\Phi$ is good. Let us fix $\varphi_o \in \Phi$ and let $\boldsymbol{m} := \boldsymbol{m}(\Phi)$. The set $\{\boldsymbol{m}(\varphi - \varphi_o) \mid \varphi \in \Phi\}$ is totally ordered, and we denote by $\boldsymbol{\ell} = \boldsymbol{\ell}_{\varphi_o}$ its submaximum. For any $\psi \neq \varphi_o$ in $\Phi$ such that $\boldsymbol{m}(\psi - \varphi_o) = \boldsymbol{m}$, we denote by $[\psi - \varphi_o]_{\boldsymbol{\ell}}$ the class of $\psi - \varphi_o$ in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(\sum \ell_i D_i)$. Let $\varphi$ be such that $\boldsymbol{m}(\varphi - \varphi_o) = \boldsymbol{m}$. Then the set

$$\Phi_{[\varphi-\varphi_o]_{\boldsymbol{\ell}}} := \{\psi \in \Phi \mid [\psi - \varphi_o]_{\boldsymbol{\ell}} = [\varphi - \varphi_o]_{\boldsymbol{\ell}}\} \subset \Phi$$

is good at 0, and $\boldsymbol{m}(\Phi_{[\varphi-\varphi_o]_{\boldsymbol{\ell}}}) < \boldsymbol{m} = \boldsymbol{m}(\Phi)$.

Let $\widetilde{\Sigma}_L \subset \mathcal{I}^{\text{ét}}_{|Y_L}$ be a finite covering of $Y_L$ (see §9.b). There exists $\boldsymbol{d}$ such that the pull-back $\widetilde{\Sigma}_{L,\boldsymbol{d}}$ of $\widetilde{\Sigma}_L$ by the ramification $\widetilde{X}_{\boldsymbol{d}} \to \widetilde{X}$ is a trivial covering of $(S^1)^\ell \times D_L$. Hence there exists a finite set $\Phi_{\boldsymbol{d}} \subset \mathscr{O}_{X_{\boldsymbol{d}}}(*D)/\mathscr{O}_{X_{\boldsymbol{d}}}$ such that the restriction of $\widetilde{\Sigma}_{L,\boldsymbol{d}}$ over $(S^1)^\ell \times \{0\}$ is equal to $\Phi_{\boldsymbol{d}} \times (S^1)^\ell \times \{0\}$. We say that $\widetilde{\Sigma}_L$ is *good* at $0 \in D_L$ if the corresponding subset $\Phi_{\boldsymbol{d}}$ is good for some (or any) $\boldsymbol{d}$ making the covering trivial. By the previous remark, if $\widetilde{\Sigma}_L$ is good at 0, it is good in some neighbourhood of $0 \in D_L$. Moreover, if $f : X' \to X$ is as in §9.e, if $\widetilde{\Sigma}_L$ is good at 0 then $\widetilde{f}^*(\widetilde{\Sigma}_L)$ is good at each point of $f^{-1}(0)$.

Let us now consider a stratified $\mathcal{I}$-covering $\widetilde{\Sigma} \subset \mathcal{I}^{\text{ét}}$ of $\widetilde{X}$ (see Definition 1.47).

***Lemma 9.14.*** — *If $\widetilde{\Sigma}_L$ is good at $0 \in D_L$, each $\widetilde{\Sigma}_I$ is good on some neighbourhood of $0$.*

*Proof.* — Assume first that $\widetilde{\Sigma}_L \to Y_L = (S^1)^L \times \Delta^{n-\ell}$ is trivial and thus $\widetilde{\Sigma}_L = \Phi \times Y_L$ for some finite good set $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. Then, if $\Delta^n$ is small enough, we have $\widetilde{\Sigma} = \bigcup_{\varphi\in\Phi} \varphi(\widetilde{\Delta}^n)$ and the assertion follows from Remark 9.13(3).

In general, one first performs a suitable ramification around the components of $D$ to reduce to the previous case. $\square$

Lastly, let us consider the global setting, where $X$ is a complex manifold and $(D_{j\in J})$ is a family of smooth divisors on $X$ which intersect normally, and the sheaf of ordered abelian groups $\mathcal{I}$ on $\partial\widetilde{X}(D_{j\in J})$ is as in Definitions 9.4 and 9.5. For any nonempty subset $I$ of $J$, we set $D_I=\bigcap_{i\in I}D_i$ and $D_I^\circ=D_I\smallsetminus\bigcup_{j\in J\smallsetminus I}D_j$. The family $(D_I^\circ)_{\varnothing\neq I\subset J}$ is a Whitney stratification of $D=\bigcup_{j\in J}D_j$.

***Definition 9.15*** **(Global goodness).** — Let us consider a stratified $\mathcal{I}$-covering $\widetilde{\Sigma}\subset\mathcal{I}^{\text{ét}}$. We say that it is *good* if each $\widetilde{\Sigma}_I$ is good at each point of $D_I^\circ$.

**9.d. Stokes filtrations on local systems.** — As above, $(D_{j\in J})$ is a family of smooth divisors on $X$ which intersect normally, and the sheaf of ordered abelian groups $\mathcal{I}$ on $\partial\widetilde{X}(D_{j\in J})$ is as in Definitions 9.4 and 9.5.

***Definition 9.16*** **(Stokes-filtered local system).** — Let $\mathscr{L}$ be a local system of $\boldsymbol{k}$-vector spaces on $\partial\widetilde{X}(D_{j\in J})$. A *Stokes filtration* of $\mathscr{L}$ is a $\mathcal{I}$-filtration of $\mathscr{L}$, in the sense of Definition 1.48. We denote by $(\mathscr{L},\mathscr{L}_\bullet)$ a Stokes-filtered local system.

***Definition 9.17*** **(Goodness).** — We say that a Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$ is *good* if its associated stratified $\mathcal{I}$-covering $\widetilde{\Sigma}(\mathscr{L})\subset\mathcal{I}^{\text{ét}}$, which is the union of the supports of the various $\operatorname{gr}\mathscr{L}_{|Y_I}$ (with $Y_I=\varpi^{-1}(D_I^\circ)$), is good.

***Theorem 9.18.*** — *Let us fix a good stratified $\mathcal{I}$-covering $\widetilde{\Sigma}\subset\mathcal{I}^{\text{ét}}$ and let $(\mathscr{L},\mathscr{L}_\bullet)$, $(\mathscr{L}',\mathscr{L}'_\bullet)$ be Stokes-filtered local systems on $\widetilde{X}(D_{j\in J})$ such that their associated $\mathcal{I}$-stratified coverings $\widetilde{\Sigma}(\mathscr{L}),\widetilde{\Sigma}(\mathscr{L}')$ are contained in $\widetilde{\Sigma}$. Let $\lambda:(\mathscr{L},\mathscr{L}_\bullet)\to(\mathscr{L}',\mathscr{L}'_\bullet)$ be a morphism of local systems which is compatible with the Stokes filtrations. Then $\lambda$ is strict.*

***Corollary 9.19.*** — *If $\widetilde{\Sigma}$ is good, the category of Stokes-filtered local systems satisfying $\widetilde{\Sigma}(\mathscr{L})\subset\widetilde{\Sigma}$ is abelian.*

This will be a consequence of the following generalization of Theorem 3.5.

***Proposition 9.20.*** — *Assume that $\widetilde{\Sigma}(\mathscr{L}),\widetilde{\Sigma}(\mathscr{L}')\subset\widetilde{\Sigma}$ for some good stratified $\mathcal{I}$-covering $\widetilde{\Sigma}\subset\mathcal{I}^{\text{ét}}$. Let $\lambda$ be a morphism of local systems compatible with the Stokes filtrations. Then, in the neighbourhood of any point of $\partial\widetilde{X}(D_{j\in J})$ there exist gradations of the Stokes filtrations such that the morphism is diagonal with respect to them. In particular, it is strict, and the natural $\mathcal{I}$-filtrations on the local systems* $\operatorname{Ker}\lambda$, $\operatorname{Im}\lambda$ *and* $\operatorname{Coker}\lambda$ *are good Stokes-filtered local systems. Their associated stratified $\mathcal{I}$-coverings satisfy*

$$\widetilde{\Sigma}(\operatorname{Ker}\lambda)\subset\widetilde{\Sigma}(\mathscr{L}),\quad\widetilde{\Sigma}(\operatorname{Coker}\lambda)\subset\widetilde{\Sigma}(\mathscr{L}'),\quad\widetilde{\Sigma}(\operatorname{Im}\lambda)\subset\widetilde{\Sigma}(\mathscr{L})\cap\widetilde{\Sigma}(\mathscr{L}').$$

*Preliminary reductions.* — As in the one-dimensional case, one reduces to the non-ramified case by a suitable $\boldsymbol{d}$-cyclic covering. Moreover, the strictness property is checked on the germs at any point of $\partial\widetilde{X}$, so we can work in the local setting of §9.b and restrict the filtered local system to the torus $(S^1)^\ell$. We will moreover forget about the term $\Delta^{n-\ell}$ and assume that $\ell = n$.

In the following, we will give the proof for a $\mathcal{I}$-local system on the torus $(S^1)^n$ in the non-ramified case, that is, $\mathcal{I}$ is the constant sheaf with fibre $\mathcal{P}_n = \mathbb{C}\{t_1, \dots, t_n\}[(t_1 \cdots t_n)^{-1}]/\mathbb{C}\{t_1, \dots, t_n\}$. As this sheaf satisfies the Hausdorff property, many of the arguments used in the proof of Theorem 3.5 can be extended in a straightforward way for Proposition 9.20. Nevertheless, we will give the proof with details, as the goodness condition is new here. Let us first notice, as a consequence of the definition of $\mathcal{I}$:

***Corollary 9.21.*** — *All the results of Lecture* 2 *apply to Stokes filtrations on a torus.*

*Proof.* — The $\mathbb{R}$-constructibility of $\mathscr{L}_{\leqslant\varphi}$ follows from Proposition 9.10. Lemma 2.8 has to be replaced with [**35**, Lemma 8.4.7(i)]. □

*Level structure of a Stokes filtration.* — For every $\boldsymbol{\ell} \in \mathbb{N}^n$, we define the notion of *Stokes filtration of level* $\geqslant \boldsymbol{\ell}$ on $\mathscr{L}$, by replacing the set of indices $\mathcal{P}_n = \mathbb{C}[t, t^{-1}]/\mathbb{C}[t]$ ($t = t_1, \dots, t_n$) by the set $\mathcal{P}_n(\boldsymbol{\ell}) := \mathbb{C}[t, t^{-1}]/t^{-\boldsymbol{\ell}}\mathbb{C}[t]$ (with $t^{-\boldsymbol{\ell}} := t_1^{-\boldsymbol{\ell}_1} \cdots t_n^{-\boldsymbol{\ell}_n}$). We denote by $[\cdot]_{\boldsymbol{\ell}}$ the map $\mathbb{C}[t, t^{-1}]/\mathbb{C}[t] \to \mathbb{C}[t, t^{-1}]/t^{-\boldsymbol{\ell}}\mathbb{C}[t]$. The constant sheaf $\mathcal{I}(\boldsymbol{\ell})$ is ordered as follows: for every connected open set $U$ of $(S^1)^n$ and $[\varphi]_{\boldsymbol{\ell}}, [\psi]_{\boldsymbol{\ell}} \in \mathcal{P}_n(\boldsymbol{\ell})$, we have $[\psi]_{\boldsymbol{\ell}} \leqslant_U [\varphi]_{\boldsymbol{\ell}}$ if, for some (or any) representatives $\varphi, \psi$ in $\mathbb{C}[t, t^{-1}]$, $e^{|t|^{\boldsymbol{\ell}}(\psi-\varphi)}$ has moderate growth along $D$ in a neighbourhood of $U$ in $X$ intersected with $X^*$. In particular, a Stokes filtration as defined previously has level $\geqslant 0$.

***Lemma 9.22.*** — *The natural morphism* $\mathcal{I} \to \mathcal{I}(\boldsymbol{\ell})$ *is compatible with the order.*

*Proof.* — Let $U$ be a connected open set in $(S^1)^n$ and let $K$ be a compact set in $U$. Let $\eta \in \mathcal{P}_n$ (or a representative of it). We have to show that if $e^\eta$ has moderate growth along $D$ on $\mathrm{nb}(K) \smallsetminus D$, then so does $e^{|t|^{\boldsymbol{\ell}}\eta}$. Let $\varepsilon : X' \to X$ be a projective modification as in the proof of Proposition 9.10, let $\widetilde{\varepsilon}$ be the associated morphism of real blowing up spaces, and let $K' \subset Y'$ be the inverse image of $K$ in $Y'$. Then $e^\eta$ has moderate growth along $D$ on $\mathrm{nb}(K) \smallsetminus D$ iff $e^{\eta\circ\varepsilon}$ has moderate growth along $D'$ on $\mathrm{nb}(K') \smallsetminus D' \simeq \mathrm{nb}(K) \smallsetminus D$. It is therefore enough to prove the lemma when $\eta$ is purely monomial, and the result follows from (9.9). □

Given a Stokes filtration $(\mathscr{L}, \mathscr{L}_\bullet)$ (of level $\geqslant 0$), we set

$$\mathscr{L}_{\leqslant[\varphi]_{\boldsymbol{\ell}}} = \sum_\psi \beta_{[\psi]_{\boldsymbol{\ell}}\leqslant[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}_{\leqslant\psi},$$

where the sum is taken in $\mathscr{L}$. Then

$$\mathscr{L}_{<[\varphi]_{\boldsymbol{\ell}}} := \sum_{[\psi]_{\boldsymbol{\ell}}} \beta_{[\psi]_{\boldsymbol{\ell}}<[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}_{\leqslant[\psi]_{\boldsymbol{\ell}}} = \sum_\psi \beta_{[\psi]_{\boldsymbol{\ell}}<[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}_{\leqslant\psi}.$$

We can also pre-$\mathcal{I}$-filter $\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}$ by setting, for $\psi\in\mathcal{P}_n$,

$$(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L})_{\leqslant\psi}=(\mathscr{L}_{\leqslant\psi}\cap\mathscr{L}_{\leqslant[\varphi]_{\boldsymbol{\ell}}}+\mathscr{L}_{<[\varphi]_{\boldsymbol{\ell}}})/\mathscr{L}_{<[\varphi]_{\boldsymbol{\ell}}}.$$

***Proposition 9.23.*** — *Assume $(\mathscr{L},\mathscr{L}_\bullet)$ is a Stokes filtration (of level $\geqslant 0$) and let $\Phi$ be the finite set of its exponential factors.*

(1) *For each $\boldsymbol{\ell}\in\mathbb{N}^n$, $\mathscr{L}_{\leqslant[\bullet]_{\boldsymbol{\ell}}}$ defines a Stokes filtration $(\mathscr{L},\mathscr{L}_{[\bullet]_{\boldsymbol{\ell}}})$ of level $\geqslant\boldsymbol{\ell}$ on $\mathscr{L}$, $\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}$ is locally isomorphic to $\bigoplus_{\psi,\,[\psi]_{\boldsymbol{\ell}}=[\varphi]_{\boldsymbol{\ell}}}\operatorname{gr}_\psi\mathscr{L}$, and the set of exponential factors of $(\mathscr{L},\mathscr{L}_{[\bullet]_{\boldsymbol{\ell}}})$ is $\Phi(\boldsymbol{\ell}):=\operatorname{image}(\Phi\to\mathcal{P}_n(\boldsymbol{\ell}))$.*

(2) *For every $[\varphi]_{\boldsymbol{\ell}}\in\Phi(\boldsymbol{\ell})$, $(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L},(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L})_\bullet)$ is a Stokes filtration and its set of exponential factors is the pull-back of $[\varphi]_{\boldsymbol{\ell}}$ by $\Phi\to\Phi(\boldsymbol{\ell})$.*

(3) *Let us set*

$$\bigl(\operatorname{gr}_{\boldsymbol{\ell}}\mathscr{L},(\operatorname{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet\bigr):=\bigoplus_{[\psi]_{\boldsymbol{\ell}}\in\Phi(\boldsymbol{\ell})}\bigl(\operatorname{gr}_{[\psi]_{\boldsymbol{\ell}}}\mathscr{L},(\operatorname{gr}_{[\psi]_{\boldsymbol{\ell}}}\mathscr{L})_\bullet\bigr).$$

*Then $(\operatorname{gr}_{\boldsymbol{\ell}}\mathscr{L},(\operatorname{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet)$ is a Stokes-filtered local system (of level $\geqslant 0$) which is locally isomorphic to $(\mathscr{L},\mathscr{L}_\bullet)$.*

*Proof.* — Similar to that of Proposition 3.7. ☐

***Remark 9.24.*** — Similarly to Remark 3.8, we note that, as a consequence of the last statement of the proposition, given a fixed Stokes-filtered local system $(\mathscr{G}_{\boldsymbol{\ell}},\mathscr{G}_{\boldsymbol{\ell},\bullet})$ graded at the level $\boldsymbol{\ell}\geqslant 0$, the pointed set of isomorphism classes of Stokes-filtered local systems $(\mathscr{L},\mathscr{L}_\bullet)$ equipped with an isomorphism $f_{\boldsymbol{\ell}}:(\operatorname{gr}_{\boldsymbol{\ell}}\mathscr{L},(\operatorname{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet)\xrightarrow{\sim}(\mathscr{G}_{\boldsymbol{\ell}},\mathscr{G}_{\boldsymbol{\ell},\bullet})$ is in bijection with the pointed set $H^1\bigl((S^1)^\ell,\mathscr{A}ut^{<0}(\mathscr{G}_{\boldsymbol{\ell}},\mathscr{G}_{\boldsymbol{\ell},\bullet})\bigr)$.

*Proof of Proposition* 9.20. — Let $\Phi$ be a good finite set in $\mathcal{P}_n$ such that $\#\Phi\geqslant 2$. As in Remark 9.13(5), let us set $\boldsymbol{m}=\boldsymbol{m}(\Phi)=\max\{\boldsymbol{m}(\varphi-\psi)\mid\varphi\neq\psi\in\Phi\}$ and let us fix $\varphi_o\in\Phi$ for which there exists $\varphi\in\Phi$ such that $\boldsymbol{m}(\varphi-\varphi_o)=\boldsymbol{m}$. The subset $\{\boldsymbol{m}(\varphi-\varphi_o)\mid\varphi\in\Phi\}$ is totally ordered, its maximum is $\boldsymbol{m}$, and we denote by $\boldsymbol{\ell}$ its submaximum (while $\boldsymbol{m}$ is independent of $\varphi_o$, $\boldsymbol{\ell}$ may depend on the choice $\varphi_o$). In the following, we will work in $\Phi-\varphi_o$ in order to have $\varphi_o=0$.

Let $\varphi\in\Phi-\varphi_o$. If $\boldsymbol{m}(\varphi)=\boldsymbol{m}$, then the image of $\varphi$ in $(\Phi-\varphi_o)(\boldsymbol{\ell})$ is nonzero, otherwise $\varphi$ is zero. For every $\varphi\in\Phi-\varphi_o$, the subset $(\Phi-\varphi_o)_{[\varphi]_{\boldsymbol{\ell}}}:=\{\psi\in\Phi-\varphi_o\mid[\psi]_{\boldsymbol{\ell}}=[\varphi]_{\boldsymbol{\ell}}\}$ is good (any subset of a good set is good) and $\boldsymbol{m}((\Phi-\varphi_o)_{[\varphi]_{\boldsymbol{\ell}}})\leqslant\boldsymbol{\ell}<\boldsymbol{m}$. Indeed, if $[\varphi]_{\boldsymbol{\ell}}=0$, then any $\psi\in(\Phi-\varphi_o)_{[\varphi]_{\boldsymbol{\ell}}}$ can be written as $t^{-\boldsymbol{\ell}}u_\psi(t)$ with $u(t)\in\mathbb{C}[t]$, and the difference of two such elements is written $t^{-\boldsymbol{\ell}}v_\psi(t)$ with $v(t)\in\mathbb{C}[t]$. On the other hand, if $[\varphi]_{\boldsymbol{\ell}}\neq 0$, $\psi$ is written as $\varphi+t^{-\boldsymbol{\ell}}u_\psi(t)$ and the same argument applies.

***Corollary 9.25* (of Prop. 9.23).** — *Let $(\mathscr{L},\mathscr{L}_\bullet)$ be a good Stokes filtration, let $\Phi''$ be a good finite subset of $\mathcal{P}_n$ containing $\Phi(\mathscr{L},\mathscr{L}_\bullet)$, set $\boldsymbol{m}=\boldsymbol{m}(\Phi'')$, fix $\varphi_o$ as above and let $\boldsymbol{\ell}$ be the corresponding submaximum element of $\{\boldsymbol{m}(\varphi-\varphi_o)\mid\varphi\in\Phi''\}$.*

*Then, for every $[\varphi]_{\boldsymbol{\ell}}\in(\Phi''-\varphi_o)(\boldsymbol{\ell})$, $(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}[-\varphi_o],(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}[-\varphi_o])_\bullet)$ is a good Stokes filtration and $\boldsymbol{m}_{\max}(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}[-\varphi_o],(\operatorname{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}[-\varphi_o])_\bullet)\leqslant\boldsymbol{\ell}<\boldsymbol{m}$.* ☐

Let us fix $\theta_o \in (S^1)^n$ and $\alpha_1, \dots, \alpha_n \in \mathbb{N}^*$ such that $\gcd(\alpha_1, \dots, \alpha_n) = 1$. The map $\theta \mapsto (\alpha_1\theta + \theta_{o,1}, \dots, \alpha_n\theta + \theta_{o,n})$ embeds $S^1$ in $(S^1)^n$. In the following, $S^1_{\boldsymbol{\alpha},\theta_o}$ denotes this circle.

Let $\Phi \subset \mathcal{P}_n$ be good finite set. Let us describe the Stokes hypersurfaces $\mathrm{St}(\varphi, \psi)$ with $\varphi \neq \psi \in \Phi$. Since $\varphi - \psi$ is purely monomial, it is written $u_{\boldsymbol{m}}(t)t^{-\boldsymbol{m}}$ with $\boldsymbol{m} = (m_1, \dots, m_n) \in \mathbb{N}^n \smallsetminus \{0\}$ and $u_{\boldsymbol{m}}(0) \neq 0$. Then

$$\mathrm{St}(\varphi, \psi) = \Big\{(\theta_1, \dots, \theta_n) \in (S^1)^n \mid \textstyle\sum_j m_j\theta_j - \arg u_{\boldsymbol{m}}(0) = \pm\pi/2 \bmod 2\pi\Big\},$$

so in particular it is the union of translated subtori of codimension one. As a consequence, the circle $S^1_{\boldsymbol{\alpha},\theta_o}$ intersects transversally every Stokes hypersurface. We call *Stokes points* with respect to $\Phi$ the intersection points when $\varphi, \psi$ vary in $\Phi$.

**Lemma 9.26.** — *Let $I$ be any open interval of $S^1_{\boldsymbol{\alpha},\theta_o}$ such that, for any $\varphi, \psi \in \Phi$, $\mathrm{card}(I \cap \mathrm{St}(\varphi, \psi)) \leqslant 1$. Then there exists an open neighbourhood $\mathrm{nb}(I)$ such that the decompositions* (1.39) *hold on* $\mathrm{nb}(I)$.

*Proof.* — A proof is then similar to that of Lemma 3.11 gives that $H^1(I, \mathscr{L}_{<\psi|I}) = 0$ for any $\psi$. We can then lift for any $\psi \in \Phi$ a basis of global sections of $\mathrm{gr}_\psi \mathscr{L}_{|I}$ as a family sections of $\mathscr{L}_{\leqslant\psi|I}$, which are defined on some $\mathrm{nb}(I)$. The images of these sections in $\mathrm{gr}_\psi \mathscr{L}_{|\mathrm{nb}(I)}$ restrict to the given basis of $\mathrm{gr}_\psi \mathscr{L}_{|I}$ and thus form a basis of $\mathrm{gr}_\psi \mathscr{L}_{|\mathrm{nb}(I)}$ if $\mathrm{nb}(I)$ is simply connected, since $\mathrm{gr}_\psi \mathscr{L}$ is a locally constant sheaf. We therefore get a section $\mathrm{gr}_\psi \mathscr{L}_{|\mathrm{nb}(I)} \to \mathscr{L}_{\leqslant\psi|\mathrm{nb}(I)}$ of the projection $\mathscr{L}_{\leqslant\psi|\mathrm{nb}(I)} \to \mathrm{gr}_\psi \mathscr{L}_{|\mathrm{nb}(I)}$.

For every $\varphi \in \Phi$, we have a natural inclusion $\beta_{\psi\leqslant\varphi}\mathscr{L}_{\leqslant\psi} \hookrightarrow \mathscr{L}_{\leqslant\varphi}$, and we deduce a morphism $\bigoplus_{\psi\in\Phi} \beta_{\psi\leqslant\varphi} \mathrm{gr}_\psi \mathscr{L}_{|\mathrm{nb}(I)} \to \mathscr{L}_{\leqslant\varphi|\mathrm{nb}(I)}$, which is seen to be an isomorphism on stalks at points of $I$, hence on a sufficiently small $\mathrm{nb}(I)$, according to the local decomposition (1.39). The same result holds then for any $\eta \in \mathcal{P}_n$ instead of $\varphi$, since $\mathscr{L}_{\leqslant\eta} = \sum_{\varphi\in\Phi} \beta_{\varphi\leqslant\eta}\mathscr{L}_{\leqslant\varphi}$ and similarly for the graded pieces. ☐

**Corollary 9.27.** — *In the setting of Corollary* 9.25, *let us set $\boldsymbol{m} = (m_1, \dots, m_n)$ and $m = \sum_i m_i\alpha_i$. Let $I$ be any open interval of $S^1$ of length $\pi/m$ with no Stokes points as boundary points. Then, if* $\mathrm{nb}(I)$ *is a sufficiently small tubular neighbourhood of $I$,* $(\mathscr{L}, \mathscr{L}_\bullet)_{|\mathrm{nb}(I)} \simeq (\mathrm{gr}_{\boldsymbol{\ell}} \mathscr{L}, (\mathrm{gr}_{\boldsymbol{\ell}} \mathscr{L})_\bullet)_{|\mathrm{nb}(I)}$.

*Proof.* — By the choice of $\boldsymbol{m}$ and the definition of $m$, $I$ satisfies the assumption of Lemma 9.26 for both $(\mathscr{L}, \mathscr{L}_\bullet)$ and $(\mathrm{gr}_{\boldsymbol{\ell}} \mathscr{L}, (\mathrm{gr}_{\boldsymbol{\ell}} \mathscr{L})_\bullet)$, hence, when restricted to $\mathrm{nb}(I)$, both are isomorphic to the trivial Stokes filtration determined by $\mathrm{gr}\mathscr{L}$ restricted to $\mathrm{nb}(I)$. ☐

*End of the proof of Proposition* 9.20. — Let $\lambda : (\mathscr{L}, \mathscr{L}_\bullet) \to (\mathscr{L}', \mathscr{L}'_\bullet)$ be a morphism of Stokes-filtered local systems on $(S^1)^n$ with set of exponential factors contained in $\Phi''$. The proof that $\lambda$ is strict and that $\mathrm{Ker}\,\lambda$, $\mathrm{Im}\,\lambda$ and $\mathrm{Coker}\,\lambda$ (equipped with the naturally induced pre-$\mathcal{P}_n$-filtrations) are Stokes filtrations follows from the local decomposition of the morphism, the proof of which is be done by induction on $\boldsymbol{m} = \boldsymbol{m}(\Phi'')$, with $\Phi'' = \Phi \cup \Phi'$. The result is clear if $\boldsymbol{m} = 0$ (so $\Phi'' = \{0\}$), as both Stokes filtrations have only one jump. The remaining part of the inductive step is completely similar to the end of the proof of Theorem 3.5 by working on $\mathrm{nb}(I)$ instead of $I$, and we will not repeat it. We obtain that, for any such $I$, $\lambda_{|\mathrm{nb}(I)}$ is graded, so this ends the proof of the proposition. □

**9.e. Behaviour by pull-back.** — Let $f : X' \to X$ be a morphism of complex manifolds. Let $(D_{j\in J})$ be a family of smooth divisors in $X$ whose union $D$ is a divisor with normal crossings. We set $D' = f^{-1}(D)$, and we assume that $D' = \bigcup_{j'\in J'} D'_{j'}$ is also a divisor with normal crossings and smooth components $D'_{j'\in J'}$. We will usually denote by $f : (X', D') \to (X, D)$ a mapping satisfying such properties.

According to §8.a, there is a natural morphism $\widetilde{f} : \widetilde{X}'(D'_{j'\in J'}) \to \widetilde{X}(D_{j\in J})$ lifting $f : X' \to X$. There are natural inclusions $\widetilde{\jmath} : X \smallsetminus D = X^* \hookrightarrow \widetilde{X}(D_{j\in J})$ and $\widetilde{\jmath}' : X' \smallsetminus D' = X'^* \hookrightarrow \widetilde{X}'(D'_{j'\in J'})$, and we have $\widetilde{f} \circ \widetilde{\jmath}' = \widetilde{\jmath} \circ f$.

Let us describe such a mapping in a local setting: the space $(X, D)$ is the polydisc $\Delta^n$ with coordinates $(x_1, \dots, x_n)$ and $D = \{x_1 \cdots x_\ell = 0\}$, and similarly for $(X', D')$, and $f(0) = 0$ in these coordinates. We have coordinates $(\theta_1, \dots, \theta_\ell, \rho_1, \dots, \rho_\ell, x_{\ell+1}, \dots, x_n)$ on $\widetilde{X}$, and similarly for $\widetilde{X}'$. In these local coordinates, we set $f = (f_1, \dots, f_n)$, with

$$f_1(x') = u'_1(x')x'^{\boldsymbol{k}_1}, \dots, f_\ell(x') = u'_\ell(x')x'^{\boldsymbol{k}_\ell}, \tag{9.28}$$

where $u'_j(x')$ are local units, and $\boldsymbol{k}_j = (k_{j,1}, \dots, k_{j,\ell'}) \in \mathbb{N}^{\ell'} \smallsetminus \{0\}$. We also have $f_j(0) = 0$ for $j \geqslant \ell+1$. We note that the stratum $D'_{L'}$ going through the origin in $X'$ (defined by $x'_1 = \cdots = x'_{\ell'} = 0$) is sent to the stratum $D_L$ going to the origin in $X$ (defined by $x_1 = \cdots = x_\ell = 0$), maybe not submersively.

When restricted to $\varpi'^{-1}(D'_{L'})$ defined by the equations $\rho'_{j'} = 0$, $j' = 1, \dots, \ell'$, the map $\widetilde{f}$ takes values in $\varpi^{-1}(D_L)$ and is given by the formula (with $\rho_1 = \cdots = \rho_\ell = 0$):

$$(\theta'_1, \dots, \theta'_{\ell'}, x'_{\ell'+1}, \dots, x'_{n'}) \longmapsto \begin{pmatrix} \sum k_{1,i}\theta'_i + \arg u'_1(0, x'_{\ell'+1}, \dots, x'_{n'}) \\ \vdots \\ \sum k_{\ell,i}\theta'_i + \arg u'_\ell(0, x'_{\ell'+1}, \dots, x'_{n'}) \\ f_{\ell+1}(0, x'_{\ell'+1}, \dots, x'_{n'}) \\ \vdots \\ f_n(0, x'_{\ell'+1}, \dots, x'_{n'}) \end{pmatrix} \tag{9.29}$$

Going back to the global setting, we have a natural morphism $f^* : \widetilde{f}^{-1}\widetilde{\jmath}_*\mathscr{O}_{X^*} \to \widetilde{\jmath}'_*\mathscr{O}_{X'^*}$, which sends $\widetilde{f}^{-1}(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$ to $(\widetilde{\jmath}'_*\mathscr{O}_{X'^*})^{\mathrm{lb}}$. This morphism is compatible with the order: it sends $\widetilde{f}^{-1}\mathscr{A}^{\mathrm{mod}\,D}_{\widetilde{X}(D_{j\in J})}$ to $\mathscr{A}^{\mathrm{mod}\,D'}_{\widetilde{X}'(D'_{j'\in J'})}$.

We consider the sheaves $\widetilde{\mathfrak{I}}$ and $\mathfrak{I}$ on $\widetilde{X}$ and $\widetilde{\mathfrak{I}}'$ and $\mathfrak{I}'$ on $\widetilde{X}'$, relative to the divisors $D$ and $D'$.

***Proposition 9.30***. — *The morphism $f^*$ sends $\widetilde{f}^{-1}\widetilde{\mathfrak{I}}$ to $\widetilde{\mathfrak{I}}'$ and induces a morphism $f^*: \widetilde{f}^{-1}\mathfrak{I} \to \mathfrak{I}'$, which is compatible with the order. Moreover, if $\widetilde{f}: \widetilde{X}' \to \widetilde{X}$ is open, it is injective and strictly compatible with the order.*

***Remark 9.31***. — If $\dim X = 1$, then $\widetilde{f}$ is *open*. Indeed, there are local coordinates in $X'$ where $f$ is expressed as a monomial. The assertion is easy to see in this case.

*Proof of Proposition* 9.30. — Let us prove the first statement. It is clear that $f^*$ sends $\widetilde{f}^{-1}\widetilde{\mathfrak{I}}_{\mathbf{1}}$ to $\widetilde{\mathfrak{I}}'_{\mathbf{1}}$. In general, we note that the assertion is local on $X$ and $X'$ and, given a local ramified covering $\rho_{\boldsymbol{d}}: X_{\boldsymbol{d}} \to X$, there is a commutative diagram

$$\begin{tikzcd}
X'_{\boldsymbol{d}'} \arrow[r, "\rho'_{\boldsymbol{d}'}"] \arrow[d, "g"'] & X' \arrow[d, "f"] \\
X_{\boldsymbol{d}} \arrow[r, "\rho_{\boldsymbol{d}}"] & X
\end{tikzcd}$$

for a suitable $\boldsymbol{d}'$ (this is easily seen in local coordinates in $X$ and $X'$ adapted to $D$ and $D'$). The morphism $\rho'^{-1}_{\boldsymbol{d}'}f^*: \rho'^{-1}_{\boldsymbol{d}'}f^{-1}\mathscr{O}_{X^*} \to \rho'^{-1}_{\boldsymbol{d}'}\mathscr{O}_{X'^*}$ is identified with $g^*: g^{-1}\mathscr{O}_{X^*_{\boldsymbol{d}}} \to \mathscr{O}_{X'^*_{\boldsymbol{d}'}}$ since $\rho_{\boldsymbol{d}}$ and $\rho'_{\boldsymbol{d}'}$ are coverings, and $f^*$ is recovered from $g^*$ as the restriction of $\rho'_{\boldsymbol{d}',*}(g^*)$ to $\mathscr{O}_{X'^*} \subset \rho'_{\boldsymbol{d}',*}\mathscr{O}_{X'^*_{\boldsymbol{d}'}}$.

As we know that $g^*$ sends $\widetilde{g}^{-1}\varpi_{\boldsymbol{d}}^{-1}\mathscr{O}_{X_{\boldsymbol{d}}}(*D)$ to $\varpi'^{-1}_{\boldsymbol{d}'}\mathscr{O}_{X'_{\boldsymbol{d}'}}(*D')$, we conclude by applying $\rho'_{\boldsymbol{d}',*}$ and intersecting with $\widetilde{\jmath}'_*\mathscr{O}_{X'^*}$ that $f^*$ sends $\widetilde{f}^{-1}\widetilde{\mathfrak{I}}_{\boldsymbol{d}}$ to $\widetilde{\mathfrak{I}}'_{\boldsymbol{d}'}$, hence in $\widetilde{\mathfrak{I}}'$.

For the injectivity statement, it is enough to prove that, if $\widetilde{f}: \widetilde{X}' \to \widetilde{X}$ is open, then $f^*: \widetilde{f}^{-1}\widetilde{\jmath}_*\mathscr{O}_{X^*}/\widetilde{f}^{-1}(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}} \to \widetilde{\jmath}'_*\mathscr{O}_{X'^*}/(\widetilde{\jmath}'_*\mathscr{O}_{X'^*})^{\mathrm{lb}}$ is injective. Let $y' \in \partial\widetilde{X}'$ and set $y = \widetilde{f}(y') \in \partial\widetilde{X}$. Note first that $\partial\widetilde{X}' = \widetilde{f}^{-1}(\partial\widetilde{X})$. If $V(y')$ is an open neighbourhood of $y'$ in $\widetilde{X}'$, then $\widetilde{f}(V(y'))$ is an open neighbourhood of $y$ in $\widetilde{X}$ by the openness assumption, and we have

$$\widetilde{f}(V(y')) \smallsetminus \partial\widetilde{X} = \widetilde{f}\big(V(y') \smallsetminus \widetilde{f}^{-1}(\partial\widetilde{X})\big) = \widetilde{f}(V(y') \smallsetminus \partial\widetilde{X}').$$

If $\lambda$ is a local section of $\widetilde{f}^{-1}\widetilde{\jmath}_*\mathscr{O}_{X^*}$ at $y'$, it is defined on such a $V(y')$. If $f^*\lambda$ is bounded on $V(y') \smallsetminus \partial\widetilde{X}'$, then $\lambda$ is a bounded section of $\mathscr{O}_{X^*}$ on the open set $\widetilde{f}(V(y') \smallsetminus \partial\widetilde{X}') = \widetilde{f}(V(y')) \smallsetminus \partial\widetilde{X}$, hence is a local section of $\widetilde{f}^{-1}(\widetilde{\jmath}_*\mathscr{O}_{X^*})^{\mathrm{lb}}$.

Let us show the strictness property. It means that, for any $y' \in \widetilde{f}^{-1}(y)$, $\varphi \circ f \leqslant_{y'} 0$ implies $\varphi \leqslant_y 0$ if $\varphi \in \mathfrak{I}_{\widetilde{X},y}$. Let $h$ be a local equation of $D$ and set $h' = h \circ f$. The relation $\varphi \circ f \leqslant_{y'} 0$ means $|e^{\varphi} \circ f| \leqslant |h'|^{-N} = |h \circ f|^{-N}$ for some $N \geqslant 0$ on $V(y') \smallsetminus \partial\widetilde{X}'$. We then have $|e^{\varphi}| \leqslant |h|^{-N}$ on $\widetilde{f}(V(y')) \smallsetminus \partial\widetilde{X}$, hence $\varphi \leqslant_y 0$ according to the openness of $\widetilde{f}$. $\square$

In general, the map $f^*$ need not be injective. Indeed, (in the local setting) given $\varphi \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$, $f^*\varphi$ may have no poles along $D'$ near $0 \in X'$. More precisely, let $\varphi$ be a local section of $\mathcal{I}$ at $0 \in X$ and let $\widetilde{\Sigma}_\varphi \subset \mathcal{I}^{\text{ét}}$ be the image by $\varphi$ of a small open neighbourhood of 0. Then $f^*$ induces a map from $\widetilde{f}^{-1}\widetilde{\Sigma}_\varphi$ to $\mathcal{I}'^{\text{ét}}$ whose image is equal to $\widetilde{\Sigma}_{f^*\varphi}$.

With a goodness assumption (which is automatically satisfied in dimension one) we recover the injectivity.

***Lemma 9.32***. — *Assume that $\varphi \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ is purely monomial. Then,*

$$f^*\varphi = 0 \Longrightarrow \varphi = 0.$$

*Proof*. — Assume $f^*\varphi = 0$. If $\varphi = 0$, there is nothing to prove. Otherwise, let us set $\varphi = U(x)/x^{\boldsymbol{m}}$, where $U(x)$ is a local unit and $\boldsymbol{m} = (m_1, \dots, m_\ell) \in \mathbb{N}^\ell \smallsetminus \{0\}$. Using the notation above for $f$, we have $f^*\varphi = U'(x')/x'^{\boldsymbol{m}'}$, where $U' = f^*U/u_1'^{m_1} \cdots u_\ell'^{m_\ell}$ is a local unit and $\boldsymbol{m}' = m_1\boldsymbol{k}_1 + \cdots + m_\ell\boldsymbol{k}_\ell$. Then $\boldsymbol{m}' \in \mathbb{N}^{\ell'} \smallsetminus \{0\}$, so $f^*\varphi \neq 0$. $\square$

***Corollary 9.33***. — *Let $\varphi$ be a local section of $\mathcal{I}$ at $0 \in X$ which is purely monomial. Then $f^*$ in injective on $\widetilde{f}^{-1}\Sigma_\varphi$.* $\square$

We also recover a property similar to strictness.

***Lemma 9.34***. — *With the same assumption as in Lemma 9.32, if $f^*\varphi \leqslant 0$ (resp. $f^*\varphi < 0$) at $(\theta_o', 0) \in \partial\widetilde{X}'_{|D_{L'}}$, then $\varphi \leqslant 0$ (resp. $\varphi < 0$) at $(\theta_o, 0) = \widetilde{g}(\theta_o', 0)$.*

*Proof*. — We keep the same notation as above, and it is enough to consider the case $f^*\varphi < 0$, according to Lemma 9.32. The assumption can then be written as

$$\arg U'(0) - \big\langle m_1\boldsymbol{k}_1 + \cdots + m_\ell\boldsymbol{k}_\ell, \theta_o' \big\rangle \in (\pi/2, 3\pi/2) \mod 2\pi,$$

where $\langle\, ,\rangle$ is the standard scalar product on $\mathbb{R}^{\ell'}$. Notice now that $\theta_{o,j} = \langle \boldsymbol{k}_j, \theta_o' \rangle + \arg u_j'(0)$ for $j = 1, \dots, \ell$, so that the previous relation is written as

$$\arg U(0) - \sum m_j\theta_{o,j} \in (\pi/2, 3\pi/2) \mod 2\pi,$$

which precisely means that $\varphi <_{\theta_o} 0$. $\square$

Let now $(\mathscr{L}, \mathscr{L}_\bullet)$ be a Stokes-filtered local system on $\partial\widetilde{X}$. Its pull-back $f^+(\mathscr{L}, \mathscr{L}_\bullet)$ (see Definition 1.34), which is a priori a pre-$\mathcal{I}$-filtered local system, is also a Stokes-filtered local system (see Lemma 1.41), and its associated stratified $\mathcal{I}$-covering is $f^*(\widetilde{f}^{-1}\widetilde{\Sigma}(\mathscr{L}))$ (see Lemma 1.49).

***Proposition 9.35***. — *With the previous assumptions on $f$, let us assume that $(\mathscr{L}, \mathscr{L}_\bullet)$ is good. Then $f^+(\mathscr{L}, \mathscr{L}_\bullet)$ is also good.*

*Proof*. — According to the previous considerations, it remains to check that, if $\widetilde{\Sigma}$ is a good stratified $\mathcal{I}$-covering, then $f^*(\widetilde{f}^{-1}\widetilde{\Sigma})$ is also good, and this reduces to showing that, if $\varphi \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ is purely monomial, then so is $f^*\varphi$, a property that we already saw in the proof of Lemma 9.32. $\square$

**9.f. Partially regular Stokes-filtered local systems.** — In the setting of §9.a, let $(\mathscr{L},\mathscr{L}_\bullet)$ be a Stokes-filtered local system on $\partial\widetilde X(D_{j\in J})$ with associated stratified $\mathcal{I}$-covering $\widetilde\Sigma$ equal to the zero section of $\mathcal{I}^{\text{ét}}_{\partial\widetilde X}$. In such a case, we will say that $(\mathscr{L},\mathscr{L}_\bullet)$ is *regular*. Then $(\mathscr{L},\mathscr{L}_\bullet)$ is the graded Stokes-filtered local system with jump at $\varphi=0$ only. The category of regular Stokes-filtered local systems is then equivalent to the category of local systems on $\partial\widetilde X(D_{j\in J})$.

We now consider the case where $(\mathscr{L},\mathscr{L}_\bullet)$ is *partially regular*, that is, there exists a decomposition $J=J'\cup J''$ such that its associated stratified covering $\widetilde\Sigma$ reduces to the zero section when restricted over $D(J'')\smallsetminus D(J')$ (recall that $D(I)=\bigcup_{j\in I}D_j$). We will set $D'=D(J')$ and $D''=D(J'')$ for simplicity. Near each point of $D''\smallsetminus D'$ the Stokes-filtered local system is regular. We will now analyze its local behaviour near $D''\cap D'$. We will restrict to a local analysis in the non-ramifed case.

According to §8.a, the identity map $X\to X$ lifts as a map $\pi:\widetilde X:=\widetilde X(D_{j\in J})\to\widetilde X':=\widetilde X(D_{j\in J'})$ and we have a commutative diagram

$$\begin{tikzcd}
\mathcal{I}^{\text{ét}}_{\widetilde X} \arrow[d, "\mu"'] & \pi^{-1}\mathcal{I}^{\text{ét}}_{\widetilde X'} \arrow[l, "q"'] \arrow[r, "\widetilde\pi"] \arrow[dl] & \mathcal{I}^{\text{ét}}_{\widetilde X'} \arrow[d, "\mu'"] \\
\widetilde X \arrow[rr, "\pi"] \arrow[dr, "\varpi"'] & & \widetilde X' \arrow[dl, "\varpi'"] \\
& X &
\end{tikzcd}$$

The boundary $\partial\widetilde X$ of $\widetilde X$ is $\varpi^{-1}(D)$ and $\partial\widetilde X'=\varpi'^{-1}(D')$.

We now consider the local setting of §9.c and we set $\ell=\ell'+\ell''$, $L'=\{1,\dots,\ell'\}$ and $L''=\{\ell'+1,\dots,\ell\}$. If $(\mathscr{L},\mathscr{L}_\bullet)$ is a non-ramified Stokes-filtered local system with associated $\mathcal{I}$-covering equal to $\widetilde\Sigma$, we assume that $\widetilde\Sigma_{|D_L}$ is a trivial covering of $Y_L=\varpi^{-1}(D_L)$. Then $\widetilde\Sigma$ is determined by a finite set $\Phi\subset\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. The partial regularity property means that the representatives $\varphi$ of the elements of $\Phi$ are holomorphic away from $D'$, that is, have no poles along $D''$, that is also, $\Phi\subset\mathscr{O}_{X,0}(*D')/\mathscr{O}_{X,0}$, and $\Phi$ defines a trivial stratified $\mathcal{I}_{\widetilde X'}$-covering $\widetilde\Sigma'$. The map $q$ induces an homeomorphism of $\pi^{-1}\widetilde\Sigma'$ onto $\widetilde\Sigma$.

**Proposition 9.36.** — *In this local setting, the category of non-ramified Stokes-filtered local systems on $\partial\widetilde X(D_{j\in L})_{|D_L}$ with associated stratified $\mathcal{I}$-covering contained in $\widetilde\Sigma$ is equivalent to the category of non-ramified Stokes-filtered local systems on $\partial\widetilde X(D_{j\in L'})_{|D_L}$ with associated stratified $\mathcal{I}$-covering contained in $\widetilde\Sigma'$, equipped with commuting automorphisms $T_k$ ($k\in L''$).*

*Proof.* — Let us first consider the following general setting: $\mathscr{F}$ is a $\mathbb{R}$-constructible sheaf on $Z\times(S^1)^k$, where $Z$ is a nice space (e.g. a subanalytic subset of $\mathbb{R}^n$). We denote by $\pi : Z \times (S^1)^k \to Z$ the projection and by $\rho : Z \times \mathbb{R}^k \to Z \times (S^1)^k$ the map $(z, \theta_1, \dots, \theta_k) \mapsto (z, e^{i\theta_1}, \dots, e^{i\theta_k})$. We will also set $\widetilde{\pi} = \rho \circ \pi$.

***Lemma 9.37.*** — *The category of $\mathbb{R}$-constructible sheaves $\mathscr{F}$ on $Z \times (S^1)^k$ whose restriction to each fibre of $\pi$ is locally constant is naturally equivalent to the category of $\mathbb{R}$-constructible sheaves on $Z$ equipped with commuting automorphisms $T_1, \dots, T_k$.*

*Proof.* — Let $\sigma_i$ $(i = 1, \dots, k)$ denote the translation by one in the direction of the $i$th coordinate in $\mathbb{R}^k$. Then the functor $\rho^{-1}$ induces an equivalence between the category of sheaves $\mathscr{F}$ on $Z \times (S^1)^k$ and that of sheaves $\mathscr{G}$ on $Z \times \mathbb{R}^k$ equipped with isomorphisms $\sigma_i^{-1}\mathscr{G} \xrightarrow{\sim} \mathscr{G}$ which commute in a natural way. It induces an equivalence between the corresponding full subcategories of $\mathbb{R}$-constructible sheaves which are locally constant in the fibres of $\pi$ and $\widetilde{\pi}$.

Let $\mathscr{G}$ be a $\mathbb{R}$-constructible sheaf on $Z \times \mathbb{R}^k$. We have a natural (dual) adjunction morphism $\mathscr{G} \to \widetilde{\pi}^! \boldsymbol{R}\widetilde{\pi}_! \mathscr{G} = \widetilde{\pi}^{-1} \boldsymbol{R}\widetilde{\pi}_! \mathscr{G}[k]$ (see [**35**, Prop. 3.3.2] for the second equality), which is an isomorphism if $\mathscr{G}$ is locally constant (hence constant) in the fibres of $\widetilde{\pi}$ (see [**35**, Prop. 2.6.7]). This shows that (via $R^k\widetilde{\pi}_!$ and $\widetilde{\pi}^{-1}$) the category of $\mathbb{R}$-constructible sheaves on $Z \times \mathbb{R}^k$ which are constant in the fibres of $\widetilde{\pi}$ is equivalent to the category of $\mathbb{R}$-constructible sheaves on $Z$. If now $\mathscr{H}$ is a $\mathbb{R}$-constructible sheaf on $Z$ with commuting automorphisms $T_i$ $(i = 1, \dots, k)$, it produces a sheaf $\mathscr{G} = \widetilde{\pi}^{-1}\mathscr{H}$ with commuting isomorphisms $\sigma_i^{-1}\mathscr{G} \simeq \mathscr{G}$ by composing the natural morphism $\sigma_i^{-1}\widetilde{\pi}^{-1}\mathscr{H} \to \widetilde{\pi}^{-1}\mathscr{H}$ with $T_i$. □

Let us end the proof of the proposition. We know that the first category considered in the proposition is equivalent to Stokes-filtered local systems indexed by $\Phi$. Each $\mathscr{L}_{\leqslant\varphi}$ is locally constant in the fibres of $\pi$, due to the local grading property of $(\mathscr{L}, \mathscr{L}_\bullet)$. We can therefore apply Lemma 9.37, since each $\mathscr{L}_{\leqslant\varphi}$ is $\mathbb{R}$-constructible, to get the essential surjectivity. The full faithfulness is obtained in the same way. □

***Remark 9.38.*** — The statement of Proposition 9.36 does not extend as it is in the ramified case. Indeed, even if $\widetilde{\Sigma}$ is regular along $D''$, a ramification may be necessary along $D''$ to trivialize $\widetilde{\Sigma}_{|Y_L}$ (e.g. a local section of $\widetilde{\Sigma}_{|Y_L}$ is written $a(y', y'')/y'^k$ for ramified coordinates $y', y''$, and $a$ is possibly not of the form $a(y', x'')$).

# LECTURE 10

# THE RIEMANN-HILBERT CORRESPONDENCE FOR GOOD MEROMORPHIC CONNECTIONS (CASE OF A SMOOTH DIVISOR)

**Summary.** This lecture is similar to Lecture 5, but we add holomorphic parameters. Moreover, we assume that no jump occurs in the the exponential factors, with respect to the parameters. This is the meaning of the goodness condition in the present setting. We will have to treat the Riemann-Hilbert functor in a more invariant way, and more arguments will be needed in the proof of the main result (equivalence of categories) in order to make it global with respect to the divisor. For the sake of simplicity, we will only consider the case of *germs* of meromorphic connections along a smooth divisor.

We consider the following setting:

- $X$ is a complex manifold and $D$ is a smooth divisor in $X$, $X^* := X \smallsetminus D$,
- $\varpi : \widetilde{X} := \widetilde{X}(D) \to X$ is the oriented real blow-up of $D$ in $X$, so that $\varpi$ is a $S^1$-fibration,
- $j : X^* \hookrightarrow X$ and $\widetilde{\jmath} : X^* \hookrightarrow \widetilde{X}$ denote the open inclusions, and $i : D \hookrightarrow X$ and $\widetilde{\imath} : \partial\widetilde{X} \hookrightarrow \widetilde{X}$ denote the closed inclusions,
- the ordered sheaf $\mathcal{I}$ on $\partial\widetilde{X}$ is as in Definitions 9.1 and 9.2.

Since $\mathcal{I}^{\text{ét}}$ is Hausdorff, the notion of $\mathcal{I}$-filtration that used in this lecture is the notion introduced in Lecture 1.

**10.a. Good formal structure of a meromorphic connection.** — We anticipate here the general definitions of Lecture 11. Let $\mathscr{M}$ be a meromorphic connection on $X$ with poles along $D$ (see §8.c) and let $x_o \in D$. We say that $\mathscr{M}$ has a *good formal structure* at $x_o$ if, after some ramification $\rho_d$ around $D$ in some neighbourhood of $x_o$, the pull-back connection $\rho_d^+\mathscr{M}$ has a good formal decomposition, that is, denoting by $\widehat{D}_d$ the space $D_d = D$ equipped with the sheaf $\mathscr{O}_{\widehat{D}_d} := \varprojlim_k \mathscr{O}_{X_d}/\mathscr{O}_{X_d}(-kD_d)$,

$$\mathscr{O}_{\widehat{D}_d} \otimes_{\mathscr{O}_{X_d}} \rho_d^+\mathscr{M} \simeq \bigoplus_{\varphi\in\Phi_d} (\mathscr{E}^{\varphi} \otimes \widehat{\mathscr{R}}_{\varphi}), \tag{10.1}$$

where $\Phi_d$ is a *good* subset of $\mathscr{O}_{X_d,x_o}(*D_d)/\mathscr{O}_{X_d,x_o}$ (see Definition 9.12), $\mathscr{E}^{\varphi} = (\mathscr{O}_{\widehat{D}_d}, d + d\varphi)$ and $\widehat{\mathscr{R}}_{\varphi}$ is a free $\mathscr{O}_{X_d,x_o}(*D_d)$-module equipped with a connection having regular singularities along $D_d$. We will usually make the abuse of identifying $\Phi_d \subset \mathscr{O}_{X_d,x_o}(*D_d)/\mathscr{O}_{X_d,x_o}$ with a set of representatives in $\Gamma(U,\mathscr{O}_{X_d}(*D_d))/\Gamma(U,\mathscr{O}_{X_d})$ for some open neighbourhood $U$ of $x_o$ in $X$ (for instance choose $U$ Stein so that $H^1(U,\mathscr{O}_X)=0$).

***Remarks 10.2***

(1) In the neighbourhood of $x_o$, each $\widehat{\mathscr{R}}_\varphi$ is obtained, after tensoring with $\mathscr{O}_{\widehat{D}_d}$, from a meromorphic connection with regular singularity, hence is locally free over $\mathscr{O}_{\widehat{D}_d}(*D_d)$. As a consequence, if $\mathscr{M}$ has a good formal structure, it is $\mathscr{O}_X(*D)$-locally free.

(2) In [**63**, Lemma 5.3.1], T. Mochizuki gives a criterion for $\mathscr{M}$ to have a good formal structure at $x_o$: choose a local isomorphism $(X,x_o)\simeq(D,x_o)\times(\mathbb{C},0)$; if there exists a *good* set $\Phi_d \subset \mathscr{O}_{X_d,x_o}(*D_d)/\mathscr{O}_{X_d,x_o}$ defined on some neighbourhood $U\subset D$ of $x_o$ such that, for any $x\in U$, the set of exponential factors of $\rho_d^+\mathscr{M}_{|\{x\}\times(\mathbb{C},0)}$ is $\Phi_{d|\{x\}\times(\mathbb{C},0)}$, then $\mathscr{M}$ has a good formal structure at $x_o$.

In [**1**, Th. 3.4.1], Y. André gives a similar criterion in terms of the Newton polygon of $\mathscr{M}_{|\{x\}\times(\mathbb{C},0)}$ and that of $\mathscr{E}nd(\mathscr{M})_{|\{x\}\times(\mathbb{C},0)}$ (so with weaker conditions a priori than in Mochizuki's criterion). Lastly, in [**37**], K. Kedlaya gives another criterion in terms of an irregularity function.

(3) For any given $\mathscr{M}$ with poles along $D$, the good formal structure property holds generically on $D$ (see [**52**]). Here, we assume that it holds all over $D$.

We now associate to a germ along $D$ of meromorphic connection having a good formal structure a $\mathfrak{I}$-covering $\widetilde{\Sigma}\subset\mathfrak{I}^{\text{ét}}$ in an intrinsic way. Notice that, due to the goodness assumption, the decomposition (10.1) is locally unique. Indeed, one checks that, if $\varphi-\psi$ is purely monomial, there is no nonzero morphism $\widehat{\mathscr{R}}_\psi\to\mathscr{E}^{\varphi-\psi}\otimes\widehat{\mathscr{R}}_\varphi$. Recall now that we have locally a cartesian square

$$\begin{tikzcd}
\rho_d^{-1}\mathfrak{I}_d^{\text{ét}} \arrow[r, "\rho_d^{\text{ét}}"] \arrow[d, "\mu_d"'] & \mathfrak{I}_d^{\text{ét}} \arrow[d, "\mu"] \\
\partial\widetilde{X}_d \arrow[r, "\rho_d"] & \partial\widetilde{X}
\end{tikzcd}$$

and $\rho_d^{-1}\mathfrak{I}_d=\varpi_d^{-1}\mathscr{O}_{X_d}(*D_d)/\mathscr{O}_{X_d}$. The set $\Phi_d$ defines a finite (trivial) covering $\widetilde{\Sigma}_d\subset\rho_d^{-1}\mathfrak{I}_d^{\text{ét}}$ locally on $D_d$, which is invariant under the Galois group of $\rho_d^{\text{ét}}$, and is therefore equal to $(\rho_d^{\text{ét}})^{-1}(\widetilde{\Sigma})$ for some locally defined $\mathfrak{I}$-covering $\widetilde{\Sigma}\subset\mathfrak{I}^{\text{ét}}$. The uniqueness statement above implies that $\widetilde{\Sigma}_d$ is uniquely determined from $\mathscr{M}$, and therefore so is $\widetilde{\Sigma}$, which thus glues globally as a $\mathfrak{I}$-covering $\widetilde{\Sigma}(\mathscr{M})$ all over $D$, since the goodness assumption is made all over $D$. We call $\widetilde{\Sigma}(\mathscr{M})$ the $\mathfrak{I}$-covering associated to $\mathscr{M}$.

**10.b. The Riemann-Hilbert functor.** — The sheaf $\mathscr{A}_{\widetilde{X}}^{\text{mod}\,D}$ is defined in §8.b. Its restriction to $\partial\widetilde{X}$ will be denoted by $\mathscr{A}_{\partial\widetilde{X}}^{\text{mod}\,D}$. We define the sheaf $\mathscr{A}_{\mathfrak{I}^{\text{ét}}}^{\text{mod}\,D}$ exactly as in §5.a. The Riemann-Hilbert functor will then be defined as a functor from the category of germs along $D$ of meromorphic connections on $X$ with poles on $D$, that is, germs along $D$ of coherent $\mathscr{O}_X(*D)$-modules with a flat connection, to the category of Stokes-filtered local systems on $\partial\widetilde{X}$.

Let $\mathscr{M}$ be a meromorphic connection on $X$ with poles along $D$. We define $\mathrm{DR}^{\mathrm{mod}\,D}_{\mathcal{J}^{\text{ét}}}(\mathscr{M})$ as in §5.c. The sheaf $\mathscr{L}_{\leqslant} := \mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}_{\mathcal{J}^{\text{ét}}}(\mathscr{M})$ is naturally a subsheaf of $\mu^{-1}\mathscr{L}$, with $\mathscr{L} := \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{H}^0\,\mathrm{DR}(\mathscr{M}_{|X^*})$. At this point, we do not even claim that $\mathscr{L}_{\leqslant}$ is a pre-$\mathcal{J}$-filtration of $\mathscr{L}$. Nevertheless, we have defined a correspondence RH from the category of germs along $D$ of meromorphic connections on $X$ with poles along $D$ to the category of pairs $(\mathscr{L},\mathscr{L}_{\leqslant})$ consisting of a local system $\mathscr{L}$ on $\partial\widetilde{X}$ and a subsheaf $\mathscr{L}_{\leqslant}$ of $\mu^{-1}\mathscr{L}$. It is clear that this correspondence is functorial.

***Definition 10.3.*** — The Riemann-Hilbert functor RH is the functor defined above.

In order to obtain an equivalence, it is however necessary to have a goodness assumption, that we fix by the choice of a *good* $\mathcal{J}$-covering $\widetilde{\Sigma}$ of $\partial\widetilde{X}$, i.e., a closed subset $\widetilde{\Sigma} \subset \mathcal{J}^{\text{ét}}$ such that $\mu$ induces a finite covering $\mu : \widetilde{\Sigma} \to \partial\widetilde{X}$ which is good (see Definition 9.15 with only one stratum). This choice will be made once and for all in this lecture. We now describe the categories involved in the correspondence.

On the one hand, the category of germs along $D$ of good meromorphic connections with poles along $D$ and with associated $\mathcal{J}$-covering $\widetilde{\Sigma}(\mathscr{M})$ contained in $\widetilde{\Sigma}$, as defined in §10.a above.

On the other hand, the definition of the category of Stokes-filtered local systems on $\partial\widetilde{X}$ with associated stratified $\mathcal{J}$-covering contained in $\widetilde{\Sigma}$ has been given in Definitions 9.16 and 9.17.

***Lemma 10.4.*** — *If $\mathscr{M}$ has a good formal structure along D, then* $\mathrm{RH}(\mathscr{M}) = (\mathscr{L},\mathscr{L}_{\leqslant})$ *is a good Stokes-filtered local system on $\partial\widetilde{X}$, that we denote by $(\mathscr{L},\mathscr{L}_{\bullet})$.*

*Proof.* — The question is local on $\partial\widetilde{X}$, and we easily reduce to the case where $\mathscr{M}$ has a good formal decomposition. The Hukuhara-Turrittin theorem with a "good holomorphic parameter" is due to Sibuya [**80, 81**]. It implies that, near any $y \in \partial\widetilde{X}$,

$$\mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}} \otimes \mathscr{M} \simeq \mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}} \otimes \Big(\bigoplus_{\psi\in\Phi}(\mathscr{E}^{\psi}\otimes\mathscr{R}_{\psi})\Big), \tag{10.5}$$

where each $\mathscr{R}_\psi$ is a locally free $\mathscr{O}_{X,x}(*D)$-module ($x = \varpi(y)$) with a flat connection having a regular singularity along $D$. So, by Hukuhara-Turrittin-Sibuya, we can assume that $\mathscr{M} \simeq \bigoplus_{\psi\in\Phi}(\mathscr{E}^{\psi}\otimes\mathscr{R}_{\psi})$, where $\Phi$ is good. Arguing on each summand and twisting by $\mathscr{E}^{-\psi}$ reduces to the case where $\mathscr{M} = \mathscr{R}$ is regular. In such a case, we have to show that $\mathscr{L}_{\leqslant}$ is a pre-$\mathcal{J}$-filtration of $\mathscr{L}$, which moreover is the trivial graded $\mathcal{J}$-filtration. Since horizontal sections of $\mathscr{R}$ have moderate growth in any meromorphic basis of $\mathscr{R}$, we have, for any $y \in \partial\widetilde{X}$, $\mathscr{L}_{\leqslant\varphi,y} = \mathscr{L}_y$ if $\mathrm{Re}(\varphi) \leqslant 0$ in some neighbourhood of $y$, and $\mathscr{L}_{\leqslant\varphi,y} = 0$ otherwise. This defines the graded $\mathcal{I}$-filtration on $\mathscr{L}$ with $\Phi = \{0\}$, according to Example 2.11(1). $\square$

***Remark 10.6***. — Unlike the dimension-one case, the complex $\mathrm{DR}^{\mathrm{mod}\,D}_{\mathcal{I}\text{ét}}(\mathscr{M})$ has cohomology in degrees $\neq 0$ even if $\mathscr{M} = \mathscr{R}$ has regular singularities along $D$, as shown by Example 8.16.

The main result of this lecture is:

***Theorem 10.7***. — *In the previous setting, the Riemann-Hilbert functor induces an equivalence between the category of good meromorphic connections with poles along $D$ and associated $\mathcal{I}$-covering contained in $\widetilde{\Sigma}$, and the category of Stokes-filtered local systems on $\partial\widetilde{X}$ with associated $\mathcal{I}$-covering contained in $\widetilde{\Sigma}$.*

**10.c. Proof of the full faithfulness in Theorem 10.7.** — Let us start with a statement analogous to Theorem 5.3:

***Lemma 10.8***. — *Let $\mathscr{M}$ be a germ at $x_o \in D$ of meromorphic connection with poles along $D$. Assume that, after some the ramification $\rho_d$ of order $d$ around $D$, the pull-back connection satisfies* (10.5) *near any $y \in \varpi^{-1}(x_o)$, where every $\psi$ entering in the decomposition is purely monomial (see Definition* 9.8*). Then $\varpi_*\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{M}) = \mathscr{H}^0\,\mathrm{DR}(\mathscr{M})$.*

*Proof*. — The complex $\mathrm{DR}(\mathscr{M})$ is regarded as a complex on $D$ (i.e., we consider the sheaf restriction to $D$ of the usual de Rham complex, since $\mathscr{M}$ is a germ along $D$).

We first claim that $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{M})$ (see §8.c) has cohomology in degree 0 at most. This is a local statement on $\partial\widetilde{X}$. Assume first $d = 1$. By the decomposition (10.5), we are reduced to the case where $\mathscr{M} = \mathscr{E}^{\psi} \otimes \mathscr{R}$, and by an argument similar to that used in (2) of the proof of Theorem 5.3, we reduce to the case where $\mathscr{M} = \mathscr{E}^{\psi}$, where $\psi$ is purely monomial. The assertion is then a consequence of Proposition 8.15. If $d \geqslant 2$, as $\mathscr{M}$ is locally stably free (see [**52**]), we can apply the projection formula

$$\widetilde{\rho}_{d,*}\big(\mathscr{A}^{\mathrm{mod}\,D_d}_{\partial\widetilde{X}_d} \otimes \varpi_d^{-1}(\rho_d^{+}\mathscr{M})\big) = (\widetilde{\rho}_{d,*}\mathscr{A}^{\mathrm{mod}\,D_d}_{\partial\widetilde{X}_d}) \otimes \varpi^{-1}\mathscr{M},$$

and, as $\widetilde{\rho}_d$ is a covering or degree $d$, $\widetilde{\rho}_{d,*}\mathscr{A}^{\mathrm{mod}\,D_d}_{\partial\widetilde{X}_d} \simeq (\mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}})^d$ locally on $\partial\widetilde{X}$. This isomorphism is compatible with connections, hence, applying this to the moderate de Rham complex gives that, locally on $\partial\widetilde{X}$, $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{M})$ is a direct summand of $\boldsymbol{R}\widetilde{\rho}_{d,*}\,\mathrm{DR}^{\mathrm{mod}\,D_d}(\rho_d^{+}\mathscr{M})$. By the first part of the proof, and since $\widetilde{\rho}_d$ is finite, the latter term has cohomology in degree zero at most, hence the claim for general $d$.

As indicated in Remark 8.10, we have

$$\boldsymbol{R}\varpi_*\,\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{M}) = \mathrm{DR}(\mathscr{M}) \quad \text{in } D^b(X). \tag{10.9}$$

Recall that this uses

$$\boldsymbol{R}\varpi_*\Big(\mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}}\underset{\varpi^{-1}\mathscr{O}_X(*D)}{\otimes}\varpi^{-1}\mathscr{M}\Big)=\boldsymbol{R}\varpi_*\Big(\mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}}\overset{\boldsymbol{L}}{\underset{\varpi^{-1}\mathscr{O}_X(*D)}{\otimes}}\varpi^{-1}\mathscr{M}\Big)\tag{10.10}$$

$$=\big(\boldsymbol{R}\varpi_*\mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}}\big)\overset{\boldsymbol{L}}{\underset{\mathscr{O}_X(*D)}{\otimes}}\mathscr{M}\tag{10.11}$$

$$=\mathscr{M},\tag{10.12}$$

where (10.10) holds because $\mathscr{M}$ is locally stably free, (10.11) because of the projection formula, and (10.12) because of Proposition 8.7 and $\mathscr{M}$ is locally stably free. Extending this to $\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{M})$ as in the proof of Proposition 8.9 gives (10.9).

By the first part of the proof, the left-hand side of (10.9) is $\boldsymbol{R}\varpi_*\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}(\mathscr{M})$. Taking $\mathscr{H}^0$ on both sides gives the lemma. □

***Corollary 10.13***. — *The Riemann-Hilbert functor in Theorem* 10.7 *is fully faithful.*

*Proof.* — The proof is similar to that given in dimension one (§5.c), as soon as we show that Lemma 10.8 applies to $\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')$ and that the analogue of Lemma 5.12 (compatibility with $\mathscr{H}\!om$) holds. The point here is that the goodness property for $\mathscr{M},\mathscr{M}'$ does not imply the goodness property for $\mathscr{H}\!om(\mathscr{M},\mathscr{M}')$. But clearly, if Hukuhara-Turrittin-Sibuya's theorem applies to $\mathscr{M},\mathscr{M}'$, it applies to $\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')$. Moreover, since $\widetilde{\Sigma}(\mathscr{M}),\widetilde{\Sigma}(\mathscr{M}')\subset\widetilde{\Sigma}$ and $\widetilde{\Sigma}$ is good, the assumption of Lemma 10.8 holds for $\mathscr{H}\!om_{\mathscr{O}_X}(\mathscr{M},\mathscr{M}')$. The argument is similar in order to prove the compatibility with $\mathscr{H}\!om$. □

**10.d. Elementary and graded equivalences.** — Recall that $\widehat{D}$ denotes the formal completion of $X$ along $D$, and $\mathscr{O}_{\widehat{D}}:=\varprojlim_k\mathscr{O}_X/\mathscr{O}_X(-kD)$. Let $\widehat{\mathscr{M}}$ be a coherent $\mathscr{O}_{\widehat{D}}(*D)$-module with a flat connection(1). We assume that it is *good*, that is, (10.1) holds for $\widehat{\mathscr{M}}$ near each point $x_o\in D$, with a good index set $\Phi_d$.

***Proposition 10.14***

(1) *Locally at $x_o\in D$, any irreducible good coherent $\mathscr{O}_{\widehat{D}}(*D)$-module $\widehat{\mathscr{M}}$ with a flat connection takes the form $\rho_{d,+}(\mathscr{E}^{\varphi}\otimes\mathscr{R}_{\widehat{D}_d})$, for some $d\geqslant 1$, some purely monomial $\varphi\in\mathscr{O}_{X_d,x_o}(*D_d)/\mathscr{O}_{X_d,x_o}$, and some free rank-one $\mathscr{O}_{X_d,x_o}(*D_d)$-module $\mathscr{R}$ with a connection having regular singularities along $D$.*

(2) *Locally at $x_o\in D$, any good coherent $\mathscr{O}_{\widehat{D}}(*D)$-module $\widehat{\mathscr{M}}$ with a flat connection has a unique decomposition $\widehat{\mathscr{M}}=\bigoplus_\alpha\widehat{\mathscr{M}_\alpha}$, where*

- *each $\widehat{\mathscr{M}_\alpha}$ is isotypical, that is, takes the form $\rho_{d_\alpha,+}(\mathscr{E}^{\varphi_\alpha}\otimes\mathscr{R}_{\alpha|\widehat{D}_{d_\alpha}})$, for some $d_\alpha\geqslant 1$, some purely monomial $\varphi_\alpha\in\mathscr{O}_{X_{d_\alpha},x_o}(*D_{d_\alpha})/\mathscr{O}_{X_{d_\alpha},x_o}$, and some free $\mathscr{O}_{X_{d_\alpha},x_o}(*D_{d_\alpha})$-module $\mathscr{R}_\alpha$ with a connection having regular singularities along $D$,*

---

(1) Be careful that the notation $\widehat{\mathscr{M}}$ will have a different meaning in Lecture 11.

- *each* $\rho_{d_\alpha,+}\mathscr{E}^{\varphi_\alpha}$ *is irreducible,*
- *if* $\alpha \neq \beta$, $\rho_{d_\alpha,+}\mathscr{E}^{\varphi_\alpha} \not\simeq \rho_{d_\beta,+}\mathscr{E}^{\varphi_\beta}$.

(3) *Any good coherent* $\mathscr{O}_{\widehat{D}}(*D)$*-module* $\widehat{\mathscr{M}}$ *with a flat connection has a unique decomposition* $\widehat{\mathscr{M}} = \bigoplus_\alpha \widehat{\mathscr{M}_\alpha}$, *where each* $\widehat{\mathscr{M}_\alpha}$ *is locally isotypical.*

*Proof.* — The proof of (1) and (2) is similar to the analogous statements when $\dim X = 1$, see the unpublished preprint [**7**], see also [**77**, Prop. 3.1 & Cor. 3.3]. Then (3) follows by local uniqueness. ☐

***Definition 10.15* (Good elementary meromorphic connections)**

Let $\mathscr{M}$ be a coherent $\mathscr{O}_X(*D)$-module with a flat connection. We say that $\mathscr{M}$ is a *good elementary* connection if, locally near any $x_o$ on $D$ and after some ramification $\rho_d$ around $D$, $\rho_d^+\mathscr{M}$ has a decomposition as $\bigoplus_{\varphi\in\Phi_d}(\mathscr{E}^\varphi\otimes\mathscr{R}_\varphi)$ with a good set $\Phi_d$, and where each $\mathscr{R}_\varphi$ is a free $\mathscr{O}_{X_d,x_o}(*D_d)$-module of finite rank with connection having regular singularities along $D$. We then denote it by $\mathscr{M}^{\mathrm{el}}$.

***Proposition 10.16*.** — *The formalization functor* $\mathscr{O}_{\widehat{D}}\otimes_{\mathscr{O}_{X|D}}\bullet$ *is an equivalence between the category of elementary germs of meromorphic connection along* $D$ *with associated* $\mathfrak{I}$*-covering contained in* $\widetilde{\Sigma}$ *(full subcategory of that of germs along* $D$ *of meromorphic connections) and the category of formal connections along* $D$ *with associated* $\mathfrak{I}$*-covering contained in* $\widetilde{\Sigma}$ .

***Proposition 10.17*.** — *The* RH *functor induces an equivalence between the category of elementary germs of meromorphic connection along* $D$ *with associated* $\mathfrak{I}$*-covering contained in* $\widetilde{\Sigma}$ *and the category of graded* $\mathfrak{I}$*-filtered local systems on* $\partial\widetilde{X}$ *with associated* $\mathfrak{I}$*-covering contained in* $\widetilde{\Sigma}$.

*Proof of Propositions* 10.16 *and* 10.17. — We notice first that, for both propositions, it is enough to prove the equivalence for the corresponding categories of germs at $x_o \in D$ for any $x_o \in D$, to get the equivalence for the categories of germs along $D$. Indeed, if this is proved, then, given an object in the target category, one can present it as the result of gluing local objects for a suitable covering of $D$. By the local essential surjectivity, one can locally lift each of the local objects, and by the local full faithfulness, one can lift in a unique way the gluing isomorphisms, which satisfy the cocycle condition, also by the local full faithfulness. This gives the global essential surjectivity. The global full faithfulness is obtained in the similar way.

Similarly, it is enough to prove both propositions in the case where the covering $\widetilde{\Sigma}$ is trivial: if the equivalence is proved after a cyclic covering around $D$, then the essential surjectivity is obtained by using the full faithfulness after ramification to lift the Galois action, and the full faithfulness is proved similarly.

In each category, each object is decomposed, and the decomposition is unique. Moreover, each morphism is also decomposed, by the goodness property. One is then reduced to proving the equivalences of both propositions in the case when $\widetilde{\Sigma}$ is a covering of degree one and, by twisting, in the regular case, where it is standard. $\square$

**10.e. Proof of the essential surjectivity in Theorem 10.7.** — The important arguments have already been explained in [**76**, §II.6]. They are an adaptation to the case with a good parameter, of the arguments given in [**46, 47**]. We will not recall all details. Firstly, as already remarked in the proof of Propositions 10.16 and 10.17, one can reduce to the case of germs at a point of $D$ and it is enough to consider the case where $\widetilde{\Sigma}$ is a trivial covering. This is the setting considered below.

Let $(\mathscr{L},\mathscr{L}_\bullet)$ be a good Stokes-filtered local system on $\partial\widetilde{X}$ with associated $\mathcal{I}$-covering contained in $\widetilde{\Sigma}$. Our goal is construct a germ $\mathscr{M}_D$ of meromorphic connection along $D$ with $\mathrm{RH}(\mathscr{M}_D)\simeq(\mathscr{L},\mathscr{L}_\bullet)$.

*Step one.* — The good Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$ is determined in a unique way, up to isomorphism, by the graded Stokes filtration $\mathrm{gr}\,\mathscr{L}$ and an element of $H^1(\partial\widetilde{X},\mathscr{A}ut^{<0}(\mu_!\,\mathrm{gr}\,\mathscr{L}))$ (see Proposition 1.43). On the other hand, by Proposition 10.17, the graded Stokes filtration $\mathrm{gr}\,\mathscr{L}$ is isomorphic to $\mathrm{RH}(\mathscr{M}^{\mathrm{el}})$ for some germ along $D$ of good elementary meromorphic connection $\mathscr{M}^{\mathrm{el}}$.

***Lemma 10.18.*** — *Let $\mathscr{E}nd^{\mathrm{mod}\,D}(\mathscr{M}^{\mathrm{el}})$ be the sheaf of horizontal sections of $\mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}}\otimes\varpi^{-1}\mathscr{H}om_{\mathcal{O}_X}(\mathscr{M}^{\mathrm{el}},\mathscr{M}^{\mathrm{el}})$ and let $\mathscr{E}nd^{\mathrm{rd}\,D}(\mathscr{M}^{\mathrm{el}})$ be the subsheaf of sections with coefficients in $\mathscr{A}^{\mathrm{rd}\,D}_{\partial\widetilde{X}}$ of holomorphic functions on $X^*$ having rapid decay along $\partial\widetilde{X}$. Then $\mathscr{E}nd^{\mathrm{mod}\,D}(\mathscr{M}^{\mathrm{el}})=\mathscr{E}nd(\mu_!\,\mathrm{gr}\,\mathscr{L})_{\leqslant 0}$ and $\mathscr{E}nd^{\mathrm{rd}\,D}(\mathscr{M}^{\mathrm{el}})=\mathscr{E}nd(\mu_!\,\mathrm{gr}\,\mathscr{L})_{<0}$.*

*Proof.* — This is a special case of the compatibility of the Riemann-Hilbert functor with $\mathscr{H}om$, already used above, and similar to Lemma 5.12. $\square$

We denote by $\mathscr{A}ut^{\mathrm{rd}\,D}(\mathscr{M}^{\mathrm{el}})$ the sheaf $\mathrm{Id}+\mathscr{E}nd^{\mathrm{rd}\,D}(\mathscr{M}^{\mathrm{el}})$. Then $\mathscr{A}ut^{\mathrm{rd}\,D}(\mathscr{M}^{\mathrm{el}})=\mathscr{A}ut^{<0}(\mu_!\,\mathrm{gr}\,\mathscr{L})$.

*Step two.* — Consider the presheaf $\mathscr{H}_D$ on $D$ such that, for any open set $U$ of $D$, $\mathscr{H}_D(U)$ consists of isomorphism classes of pairs $(\mathscr{M},\widehat{\lambda})$, where $\mathscr{M}$ is a germ along $U$ of meromorphic connection on $X$ and $\widehat{\lambda}$ is an isomorphism $\mathscr{M}_{\widehat{D}}\xrightarrow{\sim}\widehat{\mathscr{M}^{\mathrm{el}}}$.

***Lemma 10.19.*** — *The presheaf $\mathscr{H}_D$ is a sheaf.*

*Proof.* — See [**76**, Lemma II.6.2]. $\square$

Similarly, let us fix a graded Stokes-filtered local system $\mathrm{gr}\,\mathscr{L}^o$ with associated covering contained in $\widetilde{\Sigma}$, and let $\mathcal{S}t_D$ be the presheaf on $D$ such that, for any open set $U$ of $D$, $\mathcal{S}t_D(U)$ consists of isomorphism classes of pairs $[(\mathscr{L},\mathscr{L}_\bullet),\widetilde{\lambda}]$, where $(\mathscr{L},\mathscr{L}_\bullet)$ is a Stokes-filtered local system on $U$ and $\widetilde{\lambda}$ is an isomorphism $\mathrm{gr}\,\mathscr{L}\simeq\mathrm{gr}\,\mathscr{L}^o_{|U}$ of graded Stokes-filtered local systems. In particular, the associated covering $\widetilde{\Sigma}(\mathscr{L},\mathscr{L}_\bullet)$ is contained in $\widetilde{\Sigma}$.

***Lemma 10.20.*** — *The presheaf* $\mathcal{S}t_D$ *is a sheaf.*

*Proof.* — The point is to prove that, given two pairs $[(\mathscr{L},\mathscr{L}_\bullet),\widetilde{\lambda}]$ and $[(\mathscr{L}',\mathscr{L}'_\bullet),\widetilde{\lambda}']$, there is at most one isomorphism between them. If there exists one isomorphism, we can assume that $(\mathscr{L}',\mathscr{L}'_\bullet)=(\mathscr{L},\mathscr{L}_\bullet)$ and we are reduced to proving that an automorphism $\lambda$ of $(\mathscr{L},\mathscr{L}_\bullet)$ is completely determined by $\operatorname{gr}\lambda$. Arguing as in the proof of Theorem 3.5, locally with respect to $D$, there exists a local trivialization of $(\mathscr{L},\mathscr{L}_\bullet)$ such that $\lambda$ is equal to $\operatorname{gr}\lambda$ in this trivializations. It is thus uniquely determined by $\operatorname{gr}\lambda$. □

***Remark 10.21.*** — Lemma 10.20 is not needed for the essential surjectivity, it is given by symmetry with Lemma 10.19. Note that the main argument in Lemma 10.19 is the faithful flatness of $\mathscr{O}_{\widehat{D}}$ over $\mathscr{O}_{X|D}$. The corresponding argument after applying RH comes from Theorem 3.5.

*Step three.* — Arguing as in [**46**], we have a natural morphism of sheaves

$$\mathscr{H}_D \longrightarrow \mathcal{S}t_D\,.$$

where the left-hand side is associated to $\mathscr{M}^{\mathrm{el}}$ and the right-hand side to $\mathrm{RH}(\mathscr{M}^{\mathrm{el}})$.

***Theorem 10.22.*** — *This morphism is an isomorphism.*

*Proof.* — See [**76**, Th. II.6.3]. □

***Remark 10.23.*** — The proof uses the *Malgrange-Sibuya theorem* on $\varpi^{-1}(x_o)\simeq S^1$, and a base change property for the Stokes sheaf $\mathcal{S}t_D$ is needed for that purpose (see [**76**, Prop. 6.9]). Note also that it is proved in loc. cit. that the Stokes sheaf $\mathcal{S}t_D$ is locally constant on $D$.

*Step four.* — We can now end the proof. Given a good Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$, we construct $\mathscr{M}^{\mathrm{el}}$ with $\mathrm{RH}(\mathscr{M}^{\mathrm{el}})\simeq\operatorname{gr}\mathscr{L}$, according to Proposition 10.17. We are then left with a class in $H^1(\partial\widetilde{X},\mathscr{A}ut^{<0}(\mu_!\operatorname{gr}\mathscr{L}))$. This defines a global section in $\Gamma(D,\mathcal{S}t_D)$ (in fact, Lemma 10.20 says that it *is* a global section of $\mathcal{S}t_D$ on $D$). By Theorem 10.22, this is also a class in $\Gamma(D,\mathscr{H}_D)$, that is, it defines a pair $(\mathscr{M},\widehat{\lambda})$. It is now clear from the construction that $\mathrm{RH}(\mathscr{M})\simeq(\mathscr{L},\mathscr{L}_\bullet)$. □

***Remarks 10.24***

(1) One can shorten the proof above in two ways: firstly, one can use directly a version of the Malgrange-Sibuya theorem with a parameter; secondly, one can avoid the introduction of the sheaves $\mathscr{H}_D,\mathrm{St}_D$, and use the full faithfulness of the Riemann-Hilbert functor to glue the locally defined meromorphic connections obtained by applying the local Riemann-Hilbert functor. This will be the strategy in the proof of Theorem 13.2. Nevertheless, it seemed interesting to emphasize these sheaves.

(2) One can deduce from the base change property mentioned in Remark 10.23 that, if we are given a holomorphic fibration $\pi : X \to D$ (such a fibration locally exists near any point of $D$) with $D$ (smooth and) simply connected, and for a fixed good $\mathcal{I}$-covering $\widetilde{\Sigma}$ of $\partial\widetilde{X} = \varpi^{-1}(D)$, the restriction functor from germs of meromorphic connections along $D$ with associated $\mathcal{I}$-covering contained in $\widetilde{\Sigma}$ to germs of meromorphic connections on $\pi^{-1}(x_o)$ with associated $\mathcal{I}$-covering contained in $\widetilde{\Sigma}_{|\varpi^{-1}(x_o)}$ is an equivalence of categories. A similar result holds for Stokes-filtered local systems.

Such a result has been generalized by T. Mochizuki to the case where $D$ has normal crossings (see [**64**]).

(3) The arguments of §10.d are useful to prove the analogue of Proposition 5.9, that is, the compatibility with the formal Riemann-Hilbert correspondence, which also holds in this case.

# LECTURE 11

# GOOD MEROMORPHIC CONNECTIONS

**Summary.** This Lecture is a prelude to the Riemann-Hilbert correspondence in higher dimensions. We explain the notion of a good formal structure for germs of meromorphic connections having poles along a divisor with normal crossings. While the notion of good formal decomposition—which is the good analogue in higher dimensions of the Turrittin-Levelt decomposition in dimension one—has been considered in [**76**] in the case of two variables, the much more efficient presentation given here is due to T. Mochizuki [**65**, Chap. 5], [**64**]. We also recall the many-variable version of the Hukuhara-Turrittin theorem, that we have already encountered in the case of a smooth divisor (proof of Lemma 10.4). The present version relies on the work of H. Majima [**42**]. In dimension two, it is proved in [**76**]. In higher dimensions, it is due to Mochizuki [**65**].

**11.a. Preliminary notation.** — All along this lecture, we will use the following notation, compatible with that of [**65**]:

- $\Delta$ is the open disc centered at 0 in $\mathbb{C}$ and of radius one, and $X = \Delta^n$, with coordinates $t_1, \dots, t_n$.
- $D$ is the divisor defined by $\prod_{i=1}^{\ell} t_i = 0$.
- For any $I \subset \{1, \dots, \ell\}$, $D_I = \bigcap_{i\in I}\{t_i = 0\}$, and $D(I^c) = \bigcup_{\substack{j\leqslant \ell\\ j\notin I}} D_j$. It will be convenient to set $L = \{1, \dots, \ell\}$, so $D_L$ is the stratum of lowest dimension of $D$.
- $\widehat{D}_I$ is $D_I$ endowed with the sheaf $\varprojlim_k \mathscr{O}_X/(t_{i\in I})^k$, that we denote $\mathscr{O}_{\widehat{D}_I}$ (it is sometimes denoted by $\mathscr{O}_{\widehat{X|D_I}}$). A germ $f \in \mathscr{O}_{\widehat{D}_I,0}$ is written as $\sum_{\nu\in\mathbb{N}^I} f_\nu t_I^\nu$ with $f_\nu \in \mathscr{O}(U \cap D_I)$ for some open neighbourhood $U$ of 0 independent of $\nu$.
- We will also denote by $\widehat{0}$ the origin endowed with the sheaf $\mathscr{O}_{\widehat{0}} := \mathbb{C}[\![t_1, \dots, t_n]\!]$.

***Example 11.1.*** — If $\ell = n = 2$, we have $D_1 = \{t_1 = 0\}$, $D_2 = \{t_2 = 0\}$, $D_{12} = \{0\}$, $D(1^c) = D_2$, $D(2^c) = D_1$, $D(\{1,2\}^c) = D(\varnothing) = \varnothing$. Given a germ $f \in \mathbb{C}\{t_1, t_2\}$, we can consider various formal expansions of $f$:

- $f = \sum_{i\in\mathbb{N}} f_i^{(1)}(t_2)t_1^i$ in $\mathscr{O}_{\widehat{D}_1}$, where all $f_i^{(1)}(t_2)$ are holomorphic in a common neighbourhood of $t_2 = 0$,

- $f=\sum_{j\in\mathbb{N}} f_j^{(2)}(t_1)t_2^j$ in $\mathscr{O}_{\widehat{D}_2}$, where all $f_j^{(2)}(t_1)$ are holomorphic in a common neighbourhood of $t_1=0$,
- $f=\sum_{(i,j)\in\mathbb{N}^2} f_{ij}t_1^i t_2^j$, with $f_{ij}\in\mathbb{C}$.

These expansions are of course related in a natural way. We will consider below the case of meromorphic functions with poles on $D$.

Any $f\in\mathscr{O}_{X,0}(*D)$ has a unique formal expansion $f=\sum_{\boldsymbol{\nu}\in\mathbb{Z}^\ell} f_{\boldsymbol{\nu}}t^{\boldsymbol{\nu}}$ with $f_{\boldsymbol{\nu}}\in\mathscr{O}_{D_L,0}$. When it exists, the minimum (for the natural partial order) of the set $\{\boldsymbol{\nu}\in\mathbb{Z}^\ell\mid f_{\boldsymbol{\nu}}\neq 0\}\cup\{0_\ell\}$ is denoted by $\operatorname{ord}^L(f)$. It belongs to $(-\mathbb{N})^\ell$ and only depends on the class $f_L$ of $f$ in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. With this definition, we have $\operatorname{ord}^L(f)=0\in(-\mathbb{N})^\ell$ iff $f_L=0\in\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. As in Definition 9.12, we then set $\boldsymbol{m}(f_L)=-\operatorname{ord}^L(f)\in\mathbb{N}^\ell$ for some (or any) lifting $f$ of $f_L$.

Similarly, for $I\subset L$ and $f\in\mathscr{O}_{X,0}(*D)$, the minimum (for the natural partial order) of the set $\{\boldsymbol{\nu}\in\mathbb{Z}^\ell\mid f_{\boldsymbol{\nu}}\neq 0\}\cup(0_I\times\mathbb{Z}^{I^c})$, *when it exists*, is denoted by $\operatorname{ord}^I(f)$. It belongs to $(-\mathbb{N})^\ell$, and only depends on the class $f_I$ of $f$ in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_X(*D(I^c))$. We have $f_I=0$ iff $\operatorname{ord}^I(f)\in\mathbb{Z}^{I^c}$. We then set $\boldsymbol{m}(f_I)=-\operatorname{ord}^I(f)\in\mathbb{N}^\ell$ for some (or any) lifting $f$ of $f_I$ in $\mathscr{O}_{X,0}(*D)$. We will also denote by $\boldsymbol{m}_I\in\mathbb{N}^I$ the $I$-component of $\boldsymbol{m}\in\mathbb{N}^\ell$. Then, when $\operatorname{ord}^I(f_I)$ exists, $f_I=0$ iff $\boldsymbol{m}_I=0$.

As a $\mathscr{O}_{D_L,0}$-module, $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ is isomorphic to

$$\bigoplus_{\varnothing\neq J\subset L}\Big(\mathscr{O}_{D_J,0}\otimes_{\mathbb{C}}(t^{-\mathbf{1}_J}\mathbb{C}[t_J^{-1}])\Big).$$

Under such an identification, we have for any $I\subset L$:

$$\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(I^c))=\bigoplus_{\substack{\varnothing\neq J\subset L\\ J\cap I\neq\varnothing}}\Big(\mathscr{O}_{D_J,0}\otimes_{\mathbb{C}}(t^{-\mathbf{1}_J}\mathbb{C}[t_J^{-1}])\Big).$$

***Lemma 11.2***. — *A germ $\widehat{f}\in\mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$ belongs to $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ iff for any $i\in L$, the class $\widehat{f}_i$ of $\widehat{f}$ in $\mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}(*D(i^c))$ belongs to $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(i^c))$.*

*Proof*. — We also have a decomposition of $\mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$ as a $\mathscr{O}_{\widehat{0}\cap D_L}$-module:

$$\bigoplus_{\varnothing\neq J\subset L}\Big(\mathscr{O}_{\widehat{0}\cap D_J}\otimes_{\mathbb{C}}(t^{-\mathbf{1}_J}\mathbb{C}[t_J^{-1}])\Big).$$

Then, a germ $\widehat{f}$ belongs to $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ iff, for any $J\neq\varnothing$, its $J$-component has coefficients in $\mathscr{O}_{D_J,0}$. The assumption of the lemma means that, for any $i\in L$ and for any $J\subset L$ with $J\neq\varnothing$ and $J\ni i$, the $J$-component of $\widehat{f}$ has coefficients in $\mathscr{O}_{D_J,0}$. This is clearly equivalent to the desired statement. ☐

## 11.b. Good formal decomposition

*Case of a smooth divisor*. — Let us revisit first the notion of good formal decomposition in the case where $D$ is smooth (i.e., $\#L=1$), as defined by (10.1). We will have in mind possible generalizations to the normal crossing case, where formal completion along various strata will be needed. Let $\mathscr{M}$ be a meromorphic connection on $X$ with poles along $D_1$. We wish to compare the following two properties:

(1) $\mathscr{M}$ has a *good formal decomposition* along $D_1$ near the origin, that is, there exists a *good set* $\Phi\subset\mathscr{O}_{X,0}(*D_1)/\mathscr{O}_{X,0}$ and a decomposition

$$\mathscr{M}_{\widehat{D}_1}:=\mathscr{O}_{\widehat{D}_1,0}\otimes_{\mathscr{O}_{X,0}}\mathscr{M}\simeq\bigoplus_{\varphi\in\Phi}(\mathscr{E}^{\varphi}\otimes\mathscr{R}_{\varphi|\widehat{D}_1}),$$

where $\mathscr{R}_{\varphi|\widehat{D}_1}=\mathscr{O}_{\widehat{D}_1,0}\otimes_{\mathscr{O}_{X,0}}\mathscr{R}_{\varphi}$ for some germ $\mathscr{R}_{\varphi}$ of meromorphic connection with regular singularity along $D_1$.

(2) The formal germ $\widehat{\mathscr{M}}=\mathscr{M}_{\widehat{0}}:=\mathscr{O}_{\widehat{0}}\otimes_{\mathscr{O}_{X,0}}\mathscr{M}$ as a good decomposition along $D_1$, that is, there exists a *good set* $\widehat{\Phi}\subset\mathscr{O}_{\widehat{0}}(*D_1)/\mathscr{O}_{\widehat{0}}$ and a decomposition

$$\widehat{\mathscr{M}}\simeq\bigoplus_{\varphi\in\Phi}(\mathscr{E}^{\varphi}\otimes\widehat{\mathscr{R}}_{\varphi})$$

where $\widehat{\mathscr{R}}_{\varphi}$ is a regular $\mathscr{O}_{\widehat{0}}(*D_1)$-connection.

***Problem 11.3.*** — *Let $\mathscr{M}$ be a meromorphic connection with poles along $D_1$. If $\mathscr{M}$ satisfies* (2)*, is it true that $\widehat{\Phi}\subset\mathscr{O}_{X,0}(*D_1)/\mathscr{O}_{X,0}$ (set then $\Phi=\widehat{\Phi}$) and $\mathscr{M}$ satisfies* (1)*?*

The problem reduces in fact, given a local basis $\widehat{e}$ of the $\mathscr{O}_{\widehat{0}}(*D_1)$-module $\widehat{\mathscr{M}}$ adapted to the decomposition given by (2), to finding a local basis $e$ of the $\mathscr{O}_{X,0}(*D_1)$-module $\mathscr{M}$ such that the base change $e=\widehat{e}\cdot\widehat{P}$ is given by a matrix in $\mathrm{GL}(\mathscr{O}_{\widehat{0}})$ (while there is by definition a base change with matrix in $\mathrm{GL}(\mathscr{O}_{\widehat{0}}(*D_1))$). Notice also that, according to [**52**, Prop. 1.2], it is enough to find a basis $e_{\widehat{D}_1}$ of $\mathscr{M}_{\widehat{D}_1}$ with a similar property with respect to $\widehat{e}$. The solution to the problem in such a case is given by Lemma 11.19 below, in the particular case where $I=L=\{1\}$.

The conclusion is that, even in the case of a smooth divisor, the two conditions may not be equivalent, and considerations of lattices make them equivalent.

*General case.* — Let $\Phi$ be a finite subset of $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. There exists an open neighbourhood $U$ of $0$ on which each element of $\Phi$ has a representative $\varphi\in\Gamma(U,\mathscr{O}_X(*D))$ (take $U$ such that $H^1(U,\mathscr{O}_X)=0$). Let $I$ be a subset of $L$. Then the restriction of $\varphi$ to $U\smallsetminus D(I^c)$ only depends on the class $\varphi_I$ of $\varphi$ in $\mathscr{O}_U(*D)/\mathscr{O}_U(*D(I^c))$. We denote by $\Phi_I$ the image of $\Phi$ in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(I^c))$.

***Definition 11.4* (Good decomposition and good formal decomposition)**

(1) Let $\widehat{\mathscr{M}}$ be a free $\mathscr{O}_{\widehat{0}}(*D)$-module equipped with a flat connection $\widehat{\nabla}:\widehat{\mathscr{M}}\to\Omega^1_{\widehat{0}}\otimes\widehat{\mathscr{M}}$. We say that $\widehat{\mathscr{M}}$ has a *good decomposition* if there exist a *good* finite set $\widehat{\Phi}\subset\mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$ (see Definition 9.12) and a decomposition

$$\widehat{\mathscr{M}}\simeq\bigoplus_{\widehat{\varphi}\in\widehat{\Phi}}(\mathscr{E}^{\widehat{\varphi}}\otimes\widehat{\mathscr{R}}_{\widehat{\varphi}}),\tag{11.4$\,*$}$$

where $\widehat{\mathscr{R}}_{\widehat{\varphi}}$ is a free $\mathscr{O}_{\widehat{0}}(*D)$-module equipped with a flat *regular* connection.

(2) Let $\mathscr{M}$ be a free $\mathscr{O}_{X,0}(*D)$-module equipped with a flat connection $\nabla : \mathscr{M} \to \Omega^1_{X,0} \otimes \mathscr{M}$. We say that $\mathscr{M}$ has a *punctual good formal decomposition* near the origin if, for any $x \in D$ in some neighbourhood (still denoted by) $X$ of the origin, the formal connection $\mathscr{O}_{\widehat{x}}(*D) \otimes_{\mathscr{O}_X} \mathscr{M}$ has a good decomposition.

(3) Let $\mathscr{M}$ be a free $\mathscr{O}_{X,0}(*D)$-module equipped with a flat connection $\nabla : \mathscr{M} \to \Omega^1_{X,0} \otimes \mathscr{M}$. We say that $\mathscr{M}$ has a *good formal decomposition* if there exists a *good* finite set $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ and for any $I \subset L$ and any $\varphi_I \in \Phi_I$ a free $\mathscr{O}_{\widehat{D}_I,0}(*D)$-module ${}^I\widehat{\mathscr{R}}_{\varphi_I}$ equipped with a flat connection such that, on some neighbourhood $U$ of 0 where all objects are defined, ${}^I\widehat{\mathscr{R}}_{\varphi_I|\widehat{D}_I^\circ}$ is *regular*, and there is a decomposition

$$\mathscr{M}_{U\cap\widehat{D}_I^\circ} := \mathscr{O}_{U\cap\widehat{D}_I^\circ} \otimes_{\mathscr{O}_{U\smallsetminus D(I^c)}} \mathscr{M}_{|U\smallsetminus D(I^c)} \simeq \bigoplus_{\varphi_I\in\Phi_I} (\mathscr{E}^{\varphi} \otimes {}^I\widehat{\mathscr{R}}_{\varphi_I})_{|U\cap\widehat{D}_I^\circ}, \tag{11.4 $**$}$$

where $D_I^\circ := D_I \smallsetminus D(I^c)$.

In the previous definition, for a germ $\mathscr{M}$ of $\mathscr{O}_{X,0}(*D)$-module with connection, we consider a representative in some neighbourhood of the origin, and a representative of the connection.

***Remarks 11.5***

(1) We note that the property in 11.4(3) is stronger than the notion introduced in [**76**], as the *same* data $\Phi$ and $\mathscr{R}_\varphi$ are used for any stratum $D_I^\circ$ whose closure contains the stratum $D_L$ going through the base point 0.

Note also that this property is *open*, that is, if $\mathscr{M}$ has a good formal decomposition then, for any $x \in D \cap U$ ($U$ small enough), $\mathscr{M}_x$ has a good decomposition with data $\Phi_x \subset \mathscr{O}_{X,x}(*D)/\mathscr{O}_{X,x}$ and $\mathscr{R}_{\varphi,x}$. That $\Phi_x$ is good has been checked in Remark 9.13(3), so $\mathscr{M}$ satisfies 11.4(2) near the origin.

(2) On the other hand, in 11.4(2), we do not insist on the relation between the various $\Phi_x$.

(3) These definitions are stable by a twist: for instance, $\widehat{\mathscr{M}}$ has a formal $\widehat{\Phi}$-decomposition iff for some $\widehat{\eta} \in \mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$, $\widehat{\mathscr{E}^{\widehat{\eta}}} \otimes \widehat{\mathscr{M}}$ has a $(\widehat{\Phi} + \widehat{\eta})$-decomposition.

(4) In [**65**], T. Mochizuki uses a goodness condition which is slightly stronger than ours (see Definition 9.12). For our purpose, the one we use is enough.

***Lemma 11.6.*** — *If $\widehat{\mathscr{M}}$ has a good decomposition, this decomposition is unique and $\widehat{\Phi}$ is unique.*

*Proof.* — It is enough to prove that, if $\widehat{\varphi} \neq \widehat{\psi} \in \mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$, then there is no nonzero morphism $\mathscr{E}^{\widehat{\varphi}} \otimes \widehat{\mathscr{R}}_{\widehat{\varphi}} \to \mathscr{E}^{\widehat{\psi}} \otimes \widehat{\mathscr{R}}_{\widehat{\psi}}$. This amounts to showing that $\mathscr{E}^{\widehat{\varphi}-\widehat{\psi}} \otimes \widehat{\mathscr{R}}$ has no horizontal section, when $\widehat{\mathscr{R}}$ is regular. We can even assume that $\widehat{\mathscr{R}}$ has rank one, since a general $\widehat{\mathscr{R}}$ is an extension of rank-one regular meromorphic connections. So, we are looking for horizontal sections of $(\mathscr{O}_{\widehat{0}}(*D), d + d\widehat{\eta} + \omega)$ with $\widehat{\eta} \in \mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$, $\widehat{\eta} \neq 0$, and $\omega = \sum_{i=1}^{\ell} \omega_i dt_i/t_i$, $\omega_i \in \mathbb{C}$.

Assume that $f \in \mathscr{O}_{\widehat{0}}(*D)$ is a nonzero horizontal section. Then for any $i \in L$, it satisfies $z_i\partial_{z_i}(f)+z_i\partial_{z_i}(\eta)\cdot f+\omega_i f=0$. Let us consider its Newton polyhedron, which is the convex hull of the union of octants $\boldsymbol{\nu}+\mathbb{R}_+^\ell$ for which $f_{\boldsymbol{\nu}} \neq 0$ (see Notation in §11.a). By assumption, there exists $i \in L$ for which the Newton polyhedron of $z_i\partial_{z_i}(\eta)$ is not contained in $\mathbb{R}_+^\ell$. Then, on the one hand, the Newton polyhedron of $z_i\partial_{z_i}(\eta)\cdot f$ is *equal* to the Minkowski sum of that of $f$ and that of $z_i\partial_{z_i}(\eta)$, hence is not contained in that of $f$. On the other hand, the Newton polyhedron of $z_i\partial_{z_i}(f)+\omega_i f$ is contained in that of $f$. Therefore, the sum of the two corresponding terms cannot be zero, a contradiction. ☐

With the previous definition of a (punctual) good formal decomposition, it is natural to set the analogue of Problem 11.3, that is, to ask wether, given $\mathscr{M}$, the existence of a punctual good formal decomposition of $\mathscr{M}$ (near $\widehat{0}$) comes from a good formal decomposition of $\mathscr{M}$. This is a tautology if $\dim X=1$.

***Theorem 11.7* (T. Mochizuki [65, 64]).** — *Let $\mathscr{M}$ be a free $\mathscr{O}_{X,0}(*D)$-module equipped with a flat connection. If $\mathscr{M}$ has a punctual good formal decomposition near the origin, then $\mathscr{M}$ has a good formal decomposition near* $0$*, so in particular the sets $\widehat{\Phi}_x \subset \mathscr{O}_{\widehat{x}}(*D)/\mathscr{O}_{\widehat{x}}$ ($x \in D$ near $0$) are induced by a single subset $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$.*

***Remark 11.8*.** — In dimension two, T. Mochizuki already proved in [**65**] that the theorem holds with the a priori weaker condition of the existence of a formal good decomposition of $\mathscr{M}_{\widehat{0}}$. However, due to known results on the generic existence of a good decomposition (see e.g. [**52**]), the weaker condition is in fact equivalent to the stronger.

The difficulty for solving Problem 11.3 (under Condition 11.4(2) in dimension $\geqslant 3$) comes from the control of the formal coefficients of the polar parts of elements of $\widehat{\Phi}$, like $a(x_2)/x_1^k$. The original idea in [**65**] consists in working with lattices (that is, locally free $\mathscr{O}_X$-modules) instead of $\mathscr{O}_X(*D)$-modules. The notion of a good lattice will be essential (see Definition 11.12 below). One of the main achievements in [**64**] is to prove the existence of good lattices under the assumption of punctual formal decomposition of $\mathscr{M}$. This was already done in dimension two in [**65**].

The proof of Theorem 11.7 is done in two steps. In [**64**], T. Mochizuki proves the existence of a good lattice. The second step is given by Theorem 11.16 below, which follows [**65**]. We will only explain the second step.

***Definition 11.9* (Good meromorphic connection).** — Let $\mathscr{M}$ be a meromorphic connection with poles along a divisor $D$ with normal crossings. We say that $\mathscr{M}$ is a *good meromorphic connection* if, near any $x_o \in X$, there exists a ramification $\rho_{\boldsymbol{d}}: X_{\boldsymbol{d}} \to X$ around the components of $D$ going through $x_o$, such that $\rho_{\boldsymbol{d}}^+\mathscr{M}$ has a good formal decomposition near $x_o$.

***Remark 11.10*** **(Existence after blowing-up).** — The basic conjecture [**73**, Conj. I.2.5.1] asserted that, given any meromorphic bundle with connection on a complex surface, there exists a sequence of point blowing-ups such that the pull-back connection by this proper modification is good along the divisor of its poles. Fortunately, this conjecture has now been settled, in the algebraic setting by T. Mochizuki [**60**] and in general by K. Kedlaya [**37**]. The natural extension of this conjecture in higher dimension is also settled in the algebraic case by T. Mochizuki [**65**] (see also the survey [**61**]) and in the local analytic case by K. Kedlaya [**38**].

***Remark 11.11*** **(The stratified $\mathcal{I}$-covering attached to a good meromorphic connection)**

Let $\mathscr{M}$ be a germ at 0 of good meromorphic connection with poles on $D$ at most. Assume first that $\mathscr{M}$ has a good formal decomposition indexed by a good finite set $\Phi\subset\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. If $U$ is small neighbourhood of 0 in $D$ on which each $\varphi\in\Phi$ is defined, the subset $\widetilde{\Sigma}_{|\varpi^{-1}(U)}\bigcup_{\varphi\in\Phi}\varphi(\varpi^{-1}(U))\subset\mathcal{I}^{\text{ét}}_{\mathbf{1}}$ defines a stratified $\mathcal{I}_{\mathbf{1}}$-covering of $\varpi^{-1}(U)$ relative to the stratification induced by the $(Y_I)_{I\subset L}$ (see §9.b for the notation). If $\mathscr{M}$ is only assumed to be good (Definition 11.9), i.e., has a good formal decomposition after a ramification of order $\boldsymbol{d}$, one defines similarly a subset $\widetilde{\Sigma}_{|\varpi^{-1}(U)}$ of $\mathcal{I}^{\text{ét}}_{\boldsymbol{d}}$, hence of $\mathcal{I}^{\text{ét}}$, which is a stratified $\mathcal{I}$-covering with respect to the stratification $(Y_I)_{I\subset L}$.

By the uniqueness of the decomposition, this set is intrinsically attached to $\mathscr{M}$, and therefore is globally defined along $D$ when $\mathscr{M}$ is so.

**11.c. Good lattices.** — It will be implicit below that the poles of the meromorphic objects are contained in $D$.

***Definition 11.12*** **(Good decomposition and good lattice, T. Mochizuki [65])**

(1) Let $\widehat{E}$ be a free $\mathscr{O}_{\widehat{0}}$-module equipped with a flat meromorphic connection $\widehat{\nabla}:\widehat{E}\to\Omega^1_{\widehat{0}}(*D)\otimes\widehat{E}$. We say that $\widehat{E}$ has a *good decomposition* if there exist a *good* finite set $\widehat{\Phi}\subset\mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$ and a decomposition

$$(\widehat{E},\widehat{\nabla})\simeq\bigoplus_{\widehat{\varphi}\in\widehat{\Phi}}(\widehat{E}^{\widehat{\varphi}}\otimes\widehat{R}_{\widehat{\varphi}},\widehat{\nabla}),$$

where $\widehat{R}_{\widehat{\varphi}}$ is a free $\mathscr{O}_{\widehat{0}}$-module equipped with a flat meromorphic connection $\widehat{\nabla}$ such that $\widehat{\nabla}$ is logarithmic and $(\widehat{E}^{\widehat{\varphi}},\widehat{\nabla})=(\mathscr{O}_{\widehat{0}},d+d\widehat{\varphi})$.

(2) Let $E$ be a free $\mathscr{O}_{X,0}$-module equipped with a flat meromorphic connection $\nabla:E\to\Omega^1_{X,0}(*D)\otimes E$. We say that $(E,\nabla)$ is a (non-ramified) *good lattice* if $(E,\nabla)_{\widehat{0}}:=\mathscr{O}_{\widehat{0}}\otimes_{\mathscr{O}_{X,0}}(E,\nabla)$ has a good decomposition indexed by some good finite set $\widehat{\Phi}$.

In order to better understand the notion of a good lattice over $\mathscr{O}_{X,0}$, we need supplementary notions, using the notation of §11.a.

***Definition 11.13*** **($I$-goodness).** — Let $I$ be a subset of $L$. We say that a finite subset $\Phi_I$ of $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(I^c))$ is *good* if $\#\Phi_I = 1$ or for any $\varphi_I \neq \psi_I$ in $\Phi_I$, the order $\operatorname{ord}^{\boldsymbol{I}}(\varphi_I - \psi_I)$ exists (so belongs to $(-\mathbb{N})^\ell \smallsetminus \mathbb{Z}^{I^c}$) and the corresponding coefficient of $\varphi - \psi$, for some (or any) lifting of $\varphi_I - \psi_I$, does not vanish at 0 (in other words, $\varphi_I - \psi_I$ is *purely $I$-monomial*). Setting $\boldsymbol{m}(\varphi_I - \psi_I) = -\operatorname{ord}^{\boldsymbol{I}}(\varphi_I - \psi_I)$, for any *fixed* $\psi_I \in \Phi_I$, the set $\{\boldsymbol{m}(\varphi_I - \psi_I) \mid \varphi_I \in \Phi_I,\, \varphi_I \neq \psi_I\} \subset \mathbb{N}^\ell \smallsetminus \mathbb{Z}^{I^c}$ is totally ordered and its maximum, which does not depend on the choice of $\psi_I \in \Phi_I$, is denoted by $\boldsymbol{m}(\Phi_I)$. If $\#\Phi_I = 1$, we set $\boldsymbol{m}(\Phi_I) = 0$.

***Definition 11.14*** **($\Phi$ and $\Phi_I$-decompositions, T. Mochizuki [65])**

(1) Let $I \subset L$. Let $({}^I\widehat{E}, {}^I\widehat{\nabla})$ be a free $\mathscr{O}_{\widehat{D}_I,0}$-module with flat meromorphic connection and let $\Phi_I$ be a finite subset of $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(I^c))$. We say that $({}^I\widehat{E}, {}^I\widehat{\nabla})$ has a $\Phi_I$-decomposition along $D$ if there is a decomposition

$$({}^I\widehat{E}, {}^I\widehat{\nabla}) \simeq \bigoplus_{\varphi_I \in \Phi_I} (\widehat{E}^{\widetilde{\varphi}_I} \otimes {}^I\widehat{R}_{\varphi_I}, {}^I\widehat{\nabla}),$$

with $(\widehat{E}^{\widetilde{\varphi}_I}, {}^I\widehat{\nabla}) = (\mathscr{O}_{\widehat{D}_I,0}, d - d\widetilde{\varphi}_I)$ for some (or any) lifting $\widetilde{\varphi}_I$ of $\varphi_I$ in $\mathscr{O}_{\widehat{D}_I,0}(*D)$, and $({}^I\widehat{R}_{\varphi_I}, {}^I\widehat{\nabla})$ is $I$-logarithmic, that is,

$${}^I\widehat{\nabla}({}^I\widehat{R}_{\varphi_I}) \subset {}^I\widehat{R}_{\varphi_I} \otimes \Big(\mathscr{O}_{\widehat{D}_I,0}(*D(I^c)) \cdot \Omega^1_{\widehat{D}_I,0}(\log D)\Big),$$

i.e., is logarithmic only partially with respect to $D(I)$, but can have poles of arbitrary order along $D(I^c)$. If $\Phi_I$ is good, the $\Phi_I$-decomposition is said to be good.

(2) Let $(E, \nabla)$ be a free $\mathscr{O}_{X,0}$-module with flat meromorphic connection. If there exists $\Phi \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ such that, denoting by $\Phi_I$ the image of $\Phi$ in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(I^c))$, $(E, \nabla)_{|\widehat{D}_I}$ has a $\Phi_I$-decomposition for any $I \subset L$, we say that $(E, \nabla)$ has a formal $\Phi$-decomposition. If $\Phi$ is good (so that any $\Phi_I$ is good), the formal $\Phi$-decomposition is said to be good.

***Remark 11.15.*** — We have stability by twist: Fix any $\eta_I \in \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}(*D(I^c))$. Then for any lifting $\eta$ of $\eta_I$, $({}^I\widehat{E}, {}^I\widehat{\nabla} + d\eta)$ has a $(\Phi_I + \eta_I)$-decomposition iff $(E, \nabla)$ has a $\Phi_I$-decomposition.

If $({}^I\widehat{E}, {}^I\widehat{\nabla})$ has a $\Phi_I$-decomposition at 0, it has such a decomposition on $D_I \cap U$ on some neighbourhood $U$ of 0. If the $\Phi_I$-decomposition is good at 0, it is good in some neighbourhood of 0. The same property holds for a formal $\Phi$-decomposition.

***Theorem 11.16*** **(T. Mochizuki [65]).** — *Let $(E, \nabla)$ be a free $\mathscr{O}_{X,0}$-module with flat meromorphic connection. If $(E, \nabla)$ is a (non-ramified) good lattice (see Definition* 11.12(2)*) with formal exponential factors $\widehat{\Phi}$, then $\widehat{\Phi} \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ and $(E, \nabla)$ has a good formal $\Phi$-decomposition at* 0 *(see Definition* 11.14(2)*).*

***Remark 11.17.*** — One can find more properties of good lattices in [**65**], in particular good Deligne-Malgrange lattices, which are essential for proving the local and global existence of (possibly ramified) good lattices, extending in this way the result of Malgrange [**52, 54**], who shows the existence of a lattice which is generically good.

### 11.d. Proof of Theorem 11.16.

— We first generalize to the present setting the classical decomposition with respect to the eigenvalues of the principal part of the connection matrix.

Let $U$ be some open neighbourhood of 0 in $X$ and let $I \subset L$. Set $O_I = \mathscr{O}(D_I \cap U)[\![z_I]\!]$.

***Lemma 11.18* (Decomposition Lemma).** — *Let $E$ be a free $O_I$-module with a flat connection $\nabla : E \to E \otimes \Omega^1_{O_I}(*D)$. Assume that there exists $\boldsymbol{m} \in \mathbb{N}^\ell - \{0\}$, $i \in I$ and a $O_I$-basis of $E$ such that*

(1) *$m_i > 0$,*

(2) *the matrix of $\nabla$ in the given basis can be written as $z^{-\boldsymbol{m}}(\Omega + z_i A)$, where the entries of $\Omega, A$ are in $\Omega^1_{O_I}(\log D)$ and $\Omega$ is block-diagonal* $\operatorname{diag}(\Omega_1, \Omega_2)$,

(3) *the components $\Omega_1^{(i)}, \Omega_2^{(i)}$ of $\Omega_1, \Omega_2$ on $dz_i/z_i$ are such that $\Omega_1^{(i)}(0), \Omega_2^{(i)}(0)$ have no common eigenvalues.*

*Then, after possibly shrinking $U$, there exists a base change with matrix $G = \operatorname{Id} + \left(\begin{smallmatrix} 0 & z_iX \\ z_iY & 0 \end{smallmatrix}\right)$, where $X, Y$ have entries in $O_I$, such that the matrix of $\nabla$ in the new basis is $z^{-\boldsymbol{m}}\Omega'$, where $\Omega'$ has entries in $\Omega^1_{O_I}(\log D)$, is block-diagonal as $\Omega$ is and $\Omega'(0) = \Omega(0)$.*

*Moreover, such a decomposition $(E, \nabla) = (E_1, \nabla) \oplus (E_2, \nabla)$ is unique.*

*Sketch of proof.* — We can assume that $\Omega$ has entries in $O_{I'}$ with $I' = I - \{i\}$ (so that $O_I = O_{I'}[\![z_i]\!]$). Let us start with the component of $z^{-\boldsymbol{m}}(\Omega + z_i A)$ on $dz_i/z_i$. In order to find $G$ such that the transformed matrix of $z_i \nabla_{\partial_{z_i}}$ is block-diagonal, we have to solve

$$\Omega_2^{(i)} Y - Y \Omega_1^{(i)} = A_{21}^{(i)} + z_i A_{22}^{(i)} Y + z_i Y A_{11}^{(i)} + (z_i \partial_{z_i} + 1) Y + z_i^2 Y A_{12}^{(i)} Y$$

for $Y$, and a similar equation fo $X$. The assumption implies that the determinant of the endomorphism $Z \mapsto (\Omega_2^{(i)}(0) Z - Z \Omega_1^{(i)}(0))$ is not zero. Choose $U$ such that the determinant of $Z \mapsto (\Omega_2^{(i)} Z - Z \Omega_1^{(i)})$ does not vanish on $U$. We find a solution of this equation by expanding $Y$ with respect to $z_i$.

Using the integrability condition, a similar argument shows that the matrix $z^{-\boldsymbol{m}}\Omega'$ is block-diagonal.

For the uniqueness, we are reduced to prove that there is no nonzero morphism between $(E_1, \nabla_1)$ and $(E_2', \nabla_2')$ (with obvious notation). The proof is similar. ☐

*Proof of Theorem* 11.16. — We start with:

***Lemma 11.19.*** — *Let $I \subset L$ and let $({}^I\widehat{E}, {}^I\widehat{\nabla})$ be a free $\mathscr{O}_{\widehat{D}_I,0}$-module with flat meromorphic connection. Assume that there exists a good finite subset $\widehat{\Phi} \subset \mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}$*

*such that*

$$({}^I\widehat{E}, {}^I\widehat{\nabla})_{\widehat{0}} = \bigoplus_{\widehat{\varphi}\in\widehat{\Phi}} (\widehat{E}^{\widehat{\varphi}} \otimes \widehat{R}_{\widehat{\varphi}}, \widehat{\nabla}) \tag{11.19 $*$}$$

*with*

$$\widehat{\nabla}\widehat{R}_{\widehat{\varphi}} \subset \widehat{R}_{\widehat{\varphi}} \otimes \Omega^1_{\widehat{0}}(\log D) \tag{11.19 $**$}$$

*for any* $\widehat{\varphi} \in \widehat{\Phi}$*. Then* $\widehat{\Phi}_I \subset \mathscr{O}_{\widehat{D}_I,0}(*D)/\mathscr{O}_{\widehat{D}_I,0}(D(I^c))$ *(set* $\Phi_I = \widehat{\Phi}_I$*) and* $({}^I\widehat{E}, {}^I\widehat{\nabla})$ *has a* $\Phi_I$*-decomposition.*

*Proof.* — Set $\boldsymbol{m} = \boldsymbol{m}(\widehat{\Phi})$. The proof is by induction on $\#\widehat{\Phi}_I$, where $\widehat{\Phi}_I$ denotes the image of $\widehat{\Phi}$ in $\mathscr{O}_{\widehat{0}}(*D)/\mathscr{O}_{\widehat{0}}(D(I^c))$. Assume first $\#\widehat{\Phi}_I = 1$, that is, $\widehat{\Phi}_I = \{\widehat{\varphi}_I\}$. Then for any lifting $\widehat{\varphi}$ we have

$$({}^I\widehat{\nabla} - d\widehat{\varphi}){}^I\widehat{E}_{\widehat{0}} \subset {}^I\widehat{E}_{\widehat{0}} \otimes \Big(\mathscr{O}_{\widehat{0}}(*D(I^c))\Omega^1_{\widehat{0}}(\log D)\Big).$$

Writing this in some $\mathscr{O}_{\widehat{D}_I,0}$-basis of ${}^I\widehat{E}_{\widehat{0}}$ implies that $\widehat{\varphi}_I = \varphi_I \in \mathscr{O}_{\widehat{D}_I,0}(*D)/\mathscr{O}_{\widehat{D}_I,0}(D(I^c))$, hence the first statement. After a twist by $\mathscr{E}^{-\widetilde{\varphi}_I}$ for some lifting $\widetilde{\varphi}_I$ of $\varphi_I$, we can then assume that $\widehat{\Phi}_I = 0$, and we choose $\widehat{\varphi} = 0$. Then

$${}^I\widehat{\nabla}{}^I\widehat{E}_{\widehat{0}} \subset {}^I\widehat{E}_{\widehat{0}} \otimes \Big(\mathscr{O}_{\widehat{0}}(*D(I^c))\Omega^1_{\widehat{0}}(\log D)\Big),$$

and this implies

$${}^I\widehat{\nabla}{}^I\widehat{E} \subset {}^I\widehat{E} \otimes \Big(\mathscr{O}_{\widehat{D}_I,0}(*D(I^c))\Omega^1_{\widehat{D}_I,0}(\log D)\Big),$$

that is, the $\Phi_I$-decomposition with $\Phi_I = 0$.

Assume now that $\#\widehat{\Phi}_I \geqslant 2$, so $\boldsymbol{m}_I \neq 0$. Let $i \in I$ be such that $m_i > 0$ and set $I' = I - \{i\}$. Assume first that there exists $\widehat{\varphi} \in \widehat{\Phi}$ with $\operatorname{ord}^L(\widehat{\varphi}) = -\boldsymbol{m}$. By the $\widehat{\Phi}$-decomposition of $({}^I\widehat{E}, {}^I\widehat{\nabla})_{\widehat{0}}$ we can find a $\mathscr{O}_{\widehat{D}_I,0}$-basis of ${}^I\widehat{E}$ in which the matrix of ${}^I\widehat{\nabla}$ is $z^{-\boldsymbol{m}}(\Omega + z_i A)$ with $\Omega$ block-diagonal with respect to the values $(z^{\boldsymbol{m}}\widehat{\Phi})(0) \subset \mathbb{C}$ and $\Omega$ with entries in $\Omega^1_{\widehat{D}_{I'},0}(\log D)$ and $A$ with entries in $\Omega^1_{\widehat{D}_I,0}(\log D)$. Note that $\#(z^{\boldsymbol{m}}\widehat{\Phi})(0) \geqslant 2$ since $\boldsymbol{m} \neq 0$. Applying the Decomposition Lemma 11.18 (existence part), we find a flat decomposition of $({}^I\widehat{E}, {}^I\widehat{\nabla}) = \bigoplus_c ({}^I\widehat{E}_c, {}^I\widehat{\nabla}_c)$ indexed by $c \in (z^{\boldsymbol{m}}\widehat{\Phi})(0)$, whose matrix is $z^{-\boldsymbol{m}}(\Omega + z_i B)$ where $B$ is now block-diagonal as $\Omega$. Let us set $\widehat{\Phi}_c = \{\widehat{\varphi} \in \widehat{\Phi} \mid (z^{-\boldsymbol{m}}\widehat{\varphi})(0) = c\}$. Then $\widehat{\Phi} = \sqcup_c \widehat{\Phi}_c$ and, as $\boldsymbol{m}_I \neq 0$, $\widehat{\Phi}_I = \sqcup_c \widehat{\Phi}_{c,I}$. Thus, for every $c \in (z^{\boldsymbol{m}}\widehat{\Phi})(0)$, $\#\widehat{\Phi}_{c,I} < \#\widehat{\Phi}_I$. By the uniqueness in the Decomposition Lemma 11.18, we have

$$({}^I\widehat{E}_c, {}^I\widehat{\nabla}_c)_{\widehat{0}} = \bigoplus_{\widehat{\varphi}\in\widehat{\Phi}_c} (\widehat{E}^{\widehat{\varphi}} \otimes {}^I\widehat{R}_{\widehat{\varphi}}, \widehat{\nabla}).$$

We conclude by induction on $\#\widehat{\Phi}_I$.

Assume now that there is no $\widehat{\varphi} \in \widehat{\Phi}$ such that $\operatorname{ord}^L(\widehat{\varphi}) = -\boldsymbol{m}$. Then there exists $f \in \mathscr{O}_{\widehat{0}}(*D)/z^{-\boldsymbol{m}}\mathscr{O}_{\widehat{0}}$ such that $\operatorname{ord}^L(\widehat{\varphi} - f) \geqslant -\boldsymbol{m}$ for any $\widehat{\varphi} \in \widehat{\Phi}$ (and equality for some $\widehat{\varphi} \in \widehat{\Phi}$). It is thus enough to prove that $f \in \mathscr{O}_{\widehat{D}_I,0}(*D)/z^{-\boldsymbol{m}}\mathscr{O}_{\widehat{D}_I,0}$ and (see Lemma 11.2) it is equivalent to show that, for any $i \in L$, the class $f^{(i)}$ of $f$ in $\mathscr{O}_{\widehat{0}}(*D)/z_i^{-m_i}\mathscr{O}_{\widehat{0}}(*D(i^c))$ belongs to $\mathscr{O}_{\widehat{D}_I,0}(*D)/z_i^{-m_i}\mathscr{O}_{\widehat{D}_I,0}(*D(i^c))$.

Let us fix $i \in L$ and set $f^{(i)} = \sum_{\nu=m_i+1}^{\nu_i} f_\nu^{(i)} z_i^{-\nu}$. Let us prove by induction on $k \in \{0, \dots, \nu_i - m_i - 1\}$ that $f_{\nu_i-k}^{(i)} \in \mathscr{O}_{\widehat{D}_I \cap D_i}(*D)/z_i^{-m_i}\mathscr{O}_{\widehat{D}_I \cap D_i}(*D(i^c))$. Because of (11.19 $**$), for any $\widehat{\varphi} \in \widehat{\Phi}$ we have

$$\Big(z_i^{\nu_i+1}\widehat{\nabla}_{\partial_{z_i}} - \sum_{\mu=0}^{k}(\nu_i - \mu) f_{\nu_i-\mu}^{(i)} z_i^{\mu}\Big)\widehat{E}_{\widehat{\varphi}}(*D(i^c)) \subset z_i^{k+1} \cdot \widehat{E}_{\widehat{\varphi}}(*D(i^c)),$$

hence also

$$\Big(z_i^{-\nu_i+1}\,{}^I\widehat{\nabla}_{\partial_{z_i}} - \sum_{\mu=0}^{k}(\nu_i - \mu) f_{\nu_i-\mu}^{(i)} z_i^{\mu}\Big)\,{}^I\widehat{E}_{\widehat{0}}(*D(i^c)) \subset z_i^{k+1} \cdot {}^I\widehat{E}_{\widehat{0}}(*D(i^c)).$$

This implies that $z_i^{\nu_i}\,{}^I\widehat{\nabla}(z_i\partial_i)\,{}^I\widehat{E}(*D(i^c)) \subset {}^I\widehat{E}(*D(i^c))$, and by induction that the $\mathscr{O}_{\widehat{D}_I \cap D_i}(*D(i^c))$-linear endomorphism that $\big(z_i^{-\nu_i+1}\,{}^I\widehat{\nabla}_{\partial_{z_i}} - \sum_{\mu=0}^{k-1}(\nu_i - \mu) f_{\nu_i-\mu}^{(i)} z_i^{\mu}\big)$ induces on $z_i^k \cdot {}^I\widehat{E}(*D(i^c))/z_i^{k+1} \cdot {}^I\widehat{E}(*D(i^c))$ acts as $(\nu_i - k) f_{\nu_i-k}^{(i)}\,\mathrm{Id}$. It follows that $f_{\nu_i-k}^{(i)} \in \mathscr{O}_{\widehat{D}_I \cap D_i}(*D)/z_i^{-m_i}\mathscr{O}_{\widehat{D}_I \cap D_i}(*D(i^c))$. ☐

*End of the proof of Theorem* 11.16. — The assumption of the theorem is that (11.19 $*$) and (11.19 $**$) hold for $I = L$. Therefore, the assumptions of Lemma 11.19 are satisfied by $({}^I\widehat{E}, {}^I\widehat{\nabla}) := (E, \nabla)_{\widehat{D}_I}$ for any $I \subset L$, so $\widehat{\Phi}_I \subset \mathscr{O}_{\widehat{D}_I,0}(*D)/\mathscr{O}_{\widehat{D}_I,0}(D(I^c))$ for any such $I$, and in particular for any $I = \{i\}$. Lemma 11.2 then implies that $\widehat{\Phi} \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. The second part of the theorem follows then from Lemma 11.19 applied to any $I \subset L$. ☐

**11.e. The Hukuhara-Turrittin-Majima theorem.** — We keep the setting of §11.a. For short, we will denote by $\widetilde{X}$ the real blow-up space $\widetilde{X}(D_{i\in L})$, by $\varpi : \widetilde{X} \to X$ the real blowing up map and by $\mathscr{A}_{\widetilde{X}}$ the sheaf on $\widetilde{X}$ consisting of $C^\infty$ functions on $\widetilde{X}$ which are annihilated by $\overline{t}_i\overline{\partial}_{t_i}$ ($i \in L$) and $\overline{\partial}_{t_i}$ ($i \notin L$). We refer to [**42**], [**72**] and [**73**, Chap. 2] for the main properties of this sheaf. In particular, we have a Taylor map $T_L : \mathscr{A}_{\widetilde{X}} \to \mathscr{A}_{\widehat{D}_L} = \varpi^{-1}\mathscr{O}_{\widehat{D}_L}$. Notice also that $\mathscr{A}_{\widetilde{X}}$ is a subsheaf of the sheaf $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$ introduced in §8.b.

***Theorem 11.20*** **(Hukuhara-Turrittin-Majima).** — *Let $\mathscr{M}$ be a meromorphic connection with poles along $D$. Assume that $\mathscr{M}$ has a good formal decomposition (see Definition* 11.4(3)*) with set of exponential factors $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. Then, for any $\theta_o \in \varpi^{-1}(0)$, the decomposition* (11.4 $*$) *can be lifted as a decomposition*

$$\mathscr{A}_{\widetilde{X},\theta_o} \otimes_{\varpi^{-1}\mathscr{O}_{X,0}} \varpi^{-1}\mathscr{M}_0 \simeq \mathscr{A}_{\widetilde{X},\theta_o} \otimes_{\varpi^{-1}\mathscr{O}_{X,0}} \Big(\bigoplus_{\varphi\in\Phi}(\mathscr{E}^{\varphi} \otimes \mathscr{R}_{\varphi})_0\Big),$$

*where each $\mathscr{R}_\varphi$ is a meromorphic connection with poles along $D$, and regular singularity along $D$.*

We have implicitly used Theorem 11.7 to ensure that $\widehat{\Phi} \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$.

***Remark 11.21.*** — This theorem has already been used in Lemma 10.4 when $D$ is smooth, referring to Sibuya [**80, 81**] for its proof. In order to handle the case with normal crossings, the asymptotic theory developed by H. Majima [**42**] is needed. Notice also that previous approaches to this asymptotic theory can be found in [**22**].

When $\dim X = 2$, this theorem is proved in [**73**] (see Th. 2.1.1 in loc. cit.). However, the proof of loc. cit. does not seem to extend in arbitrary dimension. Here, we give an alternative proof, essentially due to T. Mochizuki [**65**], by applying another aspect of the asymptotic theory of Majima, through [**42**, §III.2, Th. 2.1] (in some cases) or [**72**, Th. A.12]. The new idea in this proof, compared to that of [**73**] in dimension two, is the use of the existence of a good lattice. See also [**26**, Appendix] for a similar proof.

***Corollary 11.22.*** — *Under the assumptions of Theorem* 11.20*, if moreover each* $\varphi \in \Phi$ *is purely monomial (see Definition* 9.8*), that is, if* $\Phi \cup \{0\}$ *is also good, then* $\mathrm{DR}^{\mathrm{mod}\, D} \mathscr{M}$ *is a sheaf.*

*Proof.* — According to Theorem 11.20, it is enough to prove the result for $\mathscr{M} = \mathscr{E}^{\varphi} \otimes \mathscr{R}_{\varphi}$, where $\varphi$ is purely monomial, since the statement is local on $\partial\widetilde{X}$. The proof is then similar to that indicated for Proposition 8.15. ☐

*Proof of Theorem* 11.20. — We will argue in a way similar to that of the proof of [**73**, Th. II.2.1.4]. For the sake of the induction, we will need to consider $\mathscr{A}(*D)$-connections and good $\mathscr{A}$-lattices. Let us fix $\theta_o \in \varpi^{-1}(0)$. By a $\mathscr{A}(*D)$-connection $\mathscr{M}^{\mathscr{A}}$, we mean a free $\mathscr{A}_{\widetilde{X},\theta_o}$-module of finite rank equipped with a flat connection

$$\nabla : \mathscr{M}^{\mathscr{A}} \longrightarrow \varpi^{-1}\Omega^1_{X,0}(*D) \otimes_{\varpi^{-1}\mathscr{O}_{X,0}} \mathscr{M}^{\mathscr{A}}.$$

By a $\mathscr{A}$-lattice $E^{\mathscr{A}}$ of $\mathscr{M}^{\mathscr{A}}$, we mean a free $\mathscr{A}_{\widetilde{X},\theta_o}$-submodule of $\mathscr{M}^{\mathscr{A}}$ such that $\mathscr{M}^{\mathscr{A}} = \mathscr{A}_{\widetilde{X},\theta_o}(*D) \otimes_{\mathscr{A}_{\widetilde{X},\theta_o}} E^{\mathscr{A}}$. The lattice is said to be good if the associated formal lattice (via the Taylor map $T_L$ at $\theta_o$) $(\widehat{E}, \widehat{\nabla})$ is a good lattice of $\widehat{\mathscr{M}}$ and if the corresponding set $\widehat{\Phi}$ is contained in $\mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$. In particular, if $(E, \nabla)$ is a good lattice in the sense of Definition 11.12(2), then $E^{\mathscr{A}} := \mathscr{A}_{\widetilde{X},\theta_o} \otimes_{\mathscr{O}_{X,0}} E$ is a good $\mathscr{A}$-lattice.

The $\mathscr{A}$-lattice $(E^{\mathscr{A}}, \nabla)$ is said to be logarithmic if $(\widehat{E}, \widehat{\nabla})$ is so, and is a Deligne logarithmic $\mathscr{A}$-lattice if moreover, for any $i \in L$, the eigenvalues of the residue of $\widehat{\nabla}_{t_i\partial_{t_i}}$ do not differ by a nonzero integer (we will fix that their real part belong to $[0, 1)$ for instance).

***Remark 11.23.*** — Arguing as in [**73**, Prop. II.2.1.10], by using the regular case of Majima's existence theorems in asymptotic analysis, we find that any Deligne $\mathscr{A}$-lattice $(E^{\mathscr{A}}, \nabla)$ is of the form $\mathscr{A}_{\widetilde{X},\theta_o} \otimes_{\varpi^{-1}\mathscr{O}_{X,0}} (E, \nabla)$ for some Deligne lattice $(E, \nabla)$.

More generally, a good $\mathscr{A}$-lattice is a good Deligne $\mathscr{A}$-lattice if each component $\widehat{R}_\varphi$ in the decomposition of 11.12(1) is a Deligne logarithmic lattice.

***Lemma 11.24.*** — *Assume that $\mathscr{M}^{\mathscr{A}}$ has a good $\mathscr{A}$-lattice. Then it has a good Deligne $\mathscr{A}$-lattice.*

*Sketch of proof.* — Because the Taylor map $T_L$ is onto (Borel-Ritt), any formal base change on $\widehat{\mathscr{M}}$ can be lifted as a $\mathscr{A}_{\widetilde{X},\theta_o}(*D)$-base change on $\mathscr{M}^{\mathscr{A}}$. The result follows. $\square$

We now argue by induction on $\#\Phi$. If $\#\Phi = 1$, we can twist by $\mathscr{E}^{-\varphi}$ to reduce to the logarithmic case, which is solved by the remark above. If $\#\Phi \geqslant 2$, since we know that $\mathscr{M}$ has a good lattice (see [**64**] and Remark 11.17) we can assume that $\mathscr{M}^{\mathscr{A}}$ has a good Deligne $\mathscr{A}$-lattice $E^{\mathscr{A}}$ by the lemma above (or directly by using a Deligne-Malgrange lattice $E$ given by [**64**]), and we fix a basis of this $\mathscr{A}$-lattice lifting a basis of $\widehat{E}$ compatible with the $\Phi$-decomposition. Moreover, by twisting with some $\mathscr{E}^{-\varphi}$, we can assume that $0 \in \Phi$. We then have $\boldsymbol{m}(\Phi) \neq 0$ (see Remark 9.13(5)). Let $\varphi_o \in \Phi$ be such that $\boldsymbol{m}(\varphi_o) = \boldsymbol{m}(\Phi)$ and let $\Phi_o = \{\varphi \in \Phi \mid \boldsymbol{m}(\varphi - \varphi_o) < \boldsymbol{m}(\Phi)\}$. Then $\Phi_o \subsetneq \Phi$, since $0 \notin \Phi_o$. We denote by $(\widehat{E}_1, \widehat{\nabla}_1) \oplus (\widehat{E}_2, \widehat{\nabla}_2)$ the corresponding decomposition of $(\widehat{E}, \widehat{\nabla})$.

Assume first that all the components $\boldsymbol{m}(\Phi)_i$ $(i \in L)$ are $> 0$. Because of the surjectivity of the Taylor map $T_L$, any $\mathscr{O}_{X,0}$-basis of a Deligne formal lattice $\widehat{E}$ lifts as a $\mathscr{A}_{\widetilde{X},\theta_o}$-basis of the lifted Deligne $\mathscr{A}$-lattice $E^{\mathscr{A}}$. Therefore, the decomposition $\widehat{E} = \widehat{E}_1 \oplus \widehat{E}_2$ lifts as $E^{\mathscr{A}} = E_1^{\mathscr{A}} \oplus E_2^{\mathscr{A}}$. There exists then a $\mathscr{A}_{\widetilde{X},\theta_o}$-basis of $E^{\mathscr{A}}$ in which the matrix of $\nabla$ takes the form

$$\Omega = \begin{pmatrix} \Omega_{11} & \Omega_{12} \\ \Omega_{21} & \Omega_{22} \end{pmatrix} \tag{11.25}$$

where $\widehat{\Omega_{12}} = 0$, $\widehat{\Omega_{21}} = 0$, and $\widehat{\Omega_{11}}$ (resp. $\widehat{\Omega_{22}}$) consists of diagonal blocks $d\varphi \operatorname{Id} + \sum_{i\in L} \Lambda_{\varphi,i} dt_i/t_i$, with $\varphi \in \Phi_o$ (resp. in $\Phi \smallsetminus \Phi_o$) and each $\Lambda_{\varphi,i}$ are constant and its eigenvalues have a real part in $[0,1)$. We look for a base change

$$P = \begin{pmatrix} \operatorname{Id} & P_{12} \\ P_{21} & \operatorname{Id} \end{pmatrix} \tag{11.26}$$

where $P$ has entries in $\mathscr{A}_{\widetilde{X},\theta_o}$ and $\widehat{P_{12}} = 0$, $\widehat{P_{21}} = 0$ which splits $\Omega$, that is, such that

$$dP_{12} = -\Omega_{12} + (\Omega_{11}P_{12} - P_{12}\Omega_{22}) + P_{12}\Omega_{21}P_{12} \tag{11.27}$$

$$dP_{21} = -\Omega_{21} + (\Omega_{22}P_{21} - P_{21}\Omega_{11}) + P_{21}\Omega_{12}P_{21}. \tag{11.28}$$

The eigenvalues of the endomorphism $P_{12} \mapsto \Omega_{11}P_{12} - P_{12}\Omega_{22}$ are the $d(\varphi - \psi)$, $\varphi \in \Phi_o$ and $\psi \in \Phi \smallsetminus \Phi_o$. By assumption, for any $i \in L$, $t_i\partial_{t_i}(\varphi - \psi) = t^{-\boldsymbol{m}(\Phi)} \cdot \text{unit}$, and we can apply [**72**, Th. A.12] to get a solution of (11.27). We argue similarly for (11.28). This partial splitting of $E^{\mathscr{A}}$ extends as a partial splitting of $\mathscr{M}^{\mathscr{A}}$, and each summand satisfies the inductive assumption, with $\Phi_o$ and $\Phi\setminus\Phi_o$ as sets of exponential factors. This concludes the proof in this first case.

If some $\boldsymbol{m}(\Phi)_i$ is zero, then, in order to apply [**72**, Th. A.12] we need to ensure that, for any such $i$, $\Omega_{12|t_i=0}=0$ and $\Omega_{21|t_i=0}=0$. We also need the assumption on the eigenvalues of $\Lambda_{\varphi,i}$ for such values of $i$.

For any $I\subset L$, we have a natural map $\varpi_I:\widetilde{X}(D_{i\in L})_{|D_I}\to\widetilde{D}_I(D_{j\in L\setminus I})$, and $\mathscr{A}_I:=\mathscr{A}_{\widetilde{X}}\big/\sum_{i\in I}t_i\mathscr{A}_{\widetilde{X}}=\varpi_I^{-1}\mathscr{A}_{\widetilde{D}_I}$. We regard $\mathscr{A}_{I,\theta_o}$ as a subring of $\mathscr{A}_{\widetilde{X},\theta_o}$.

Let $L'=\{i\in L\mid \boldsymbol{m}(\Phi)_i=0\}$ and $L''=L\setminus L'$. For any $I$ subset $L'$, the residual connection $({}^I\!E^{\mathscr{A}},{}^I\nabla)$ is well defined: we set ${}^I\!E^{\mathscr{A}}=\mathscr{A}_{I,\theta_o}\otimes_{\mathscr{A}_{\widetilde{X},\theta_o}}E^{\mathscr{A}}$, and for any $j\notin I$, $\nabla_{t_j\partial_{t_j}}$ sends $E^{\mathscr{A}}$ to $E^{\mathscr{A}}(*D(L''))$, hence can be restricted to $t_i=0$ ($i\in I\subset L'$). We will prove the following property by decreasing induction on $k$:

(H$_k$) For any $I\subset L'$ with $\#I\geqslant k$ there exists a basis $\boldsymbol{e}_I^{\mathscr{A}}$ of ${}^I\!E^{\mathscr{A}}$, which restricts to $\boldsymbol{e}_J^{\mathscr{A}}$ when tensoring with $\mathscr{A}_{J,\theta_o}$ for any $J$ such that $I\subset J\subset L'$, in which the matrix of ${}^I\nabla$ is block-diagonal, i.e.,

$$({}^I\!E^{\mathscr{A}},{}^I\nabla)=({}^I\!E_1^{\mathscr{A}},{}^I\nabla_1)\oplus({}^I\!E_2^{\mathscr{A}},{}^I\nabla_2).$$

Let us set $\ell'=\#L'$. The first part of the proof shows that (H$_{\ell'}$) holds. On the other hand, the theorem will be proved if (H$_0$) holds. Assume that (H$_k$) holds and let $I\subset L'$ with $\#I=k-1$. Firstly, by a simple Mayer-Vietoris argument, the various bases $\boldsymbol{e}_J^{\mathscr{A}}$ for $J\supset I$, $J\subset L'$ and $\#J=k$ are compatible with the various restrictions $\mathscr{A}_{K,\theta_o}\otimes$, with $L'\subset K\supset J$, and therefore there exist a basis $\boldsymbol{e}_I^{\mathscr{A}}$ compatible with the various restrictions $\mathscr{A}_{J,\theta_o}\otimes$. Indeed, note first that if we are given a family $(f_J)$ of elements of $\mathscr{A}_{J,\theta_o}$ which are compatible with restrictions to $t_k=0$, $k\in L'\setminus I$, there exists $f_I\in\mathscr{A}_{I,\theta_o}$ which restricts to $f_J$ at $t_j=0$ with $J=I\cup\{j\}$: it suffices to set

$$f_I=\sum_{\#J=\#I+1}f_J-\sum_{\#J=\#I+2}f_J+\cdots.$$

Starting with any basis of ${}^I\!E$, we denote by $P_J$ the base change from this basis restricted to $t_j=0$, $j\in J$, to the basis $\boldsymbol{e}_J^{\mathscr{A}}$. The family of matrices $P_J$ is compatible with restrictions to $t_j=0$, and therefore lifts as a matrix $P_I$, which remains invertible, and defines the desired basis $\boldsymbol{e}_I^{\mathscr{A}}$.

The matrix ${}^I\Omega$ of ${}^I\nabla$ decomposes as in (11.25), where ${}^I\Omega_{12}$ and ${}^I\Omega_{21}$ restrict identically to 0 when $t_j=0$, for $j\in L'\setminus I$. According to [**72**, Th. A.12], we can now solve (11.27) and (11.28) in such a way that $P_{12}$ and $P_{21}$ restrict identically to 0 when $t_j=0$, for $j\in L'\setminus I$. Therefore, the new decomposition $({}^I\!E^{\mathscr{A}},{}^I\nabla)=({}^I\!E_1^{\mathscr{A}},{}^I\nabla_1)\oplus({}^I\!E_2^{\mathscr{A}},{}^I\nabla_2)$ restricts to the given one when $t_j=0$, $j\in L'\setminus I$, and so (H$_{k-1}$) holds. $\square$

# LECTURE 12

# LEAKY PIPES

**Summary.** In this lecture, we experiment the compatibility of the Riemann-Hilbert correspondence with direct image on a simple but not trivial example. Although we did not already defined this correspondence in full generality, we have enough material to make such a computation when the target space has dimension one. Compared with the computations in Lecture 7, we go one step further. We refer to the literature for the basic notions of holonomic $\mathscr{D}$-module and regularity (see e.g. [**10, 34, 56, 57**]).

**12.a. The setting.** — Let $\Delta$ be an open disc with coordinate $t$ and let $\mathbb{A}^1$ be the affine line with coordinate $x$. We will denote by $y$ the coordinate $1/x$ at $\infty$ on $\mathbb{A}^1$. Let $M$ be a holonomic $\mathscr{D}_\Delta[x]\langle\partial_x\rangle$-module with regular singularities (included along $x=\infty$). We denote by $S\subset\mathbb{A}^1\times\Delta$ the union of the irreducible components of its singular support distinct from $\mathbb{P}^1\times\{0\}$: away from $S\cup(\mathbb{P}^1\times\{0\})$, $M$ is a holomorphic bundle with flat connection. We will assume that $\Delta$ is small enough so that any irreducible component of the closure $\overline{S}$ of $S$ in $\mathbb{P}^1\times\Delta$ cuts $\mathbb{P}^1\times\{0\}$ and is smooth away from $\mathbb{P}^1\times\{0\}$ (see Figure 1). We denote by $p:\mathbb{P}^1\times\Delta\to\Delta$ the projection. By working in the analytic category with respect to $\mathbb{P}^1$, we also regard $M$ as a holonomic $\mathscr{D}_{\mathbb{P}^1\times\Delta}$-module with regular singularities, and we have $\mathscr{O}_{\mathbb{P}^1\times\Delta}(*\infty)\otimes_{\mathscr{O}_{\mathbb{P}^1\times\Delta}}M=M$, where $\infty$ stands for the divisor $\{\infty\}\times\Delta$ in $\mathbb{P}^1\times\Delta$.

Let us set $\mathscr{E}^x=(\mathscr{O}_\Delta[x],d+dx)$. The $\mathscr{O}_\Delta[x]\langle\partial_x\rangle$-module $\mathscr{E}^x\otimes_{\mathscr{O}_\Delta[x]}M$ has an irregular singularity along $x=\infty$. The direct image $p_+(\mathscr{E}^x\otimes_{\mathscr{O}_\Delta[x]}M)$ is a complex which satisfies $\mathscr{H}^\ell p_+(\mathscr{E}^x\otimes M)=0$ if $\ell\neq-1,0$. Moreover, $\mathscr{H}^{-1}p_+(\mathscr{E}^x\otimes M)$ is supported at the origin of $\Delta$: Indeed, if $\Delta$ is small enough, $\mathscr{H}^{-1}p_+(\mathscr{E}^x\otimes M)$ is a vector bundle on $\Delta^*$, whose fibre at $t=t_o\neq0$ can be computed as $\operatorname{Ker}[\nabla_{\partial_x}:(\mathscr{E}^x\otimes M_{t_o})\to(\mathscr{E}^x\otimes M_{t_o})]$, where $M_{t_o}=M/(t-t_o)M$. Note that $M_{t_o}$ is a regular holonomic $\mathbb{C}[x]\langle\partial_x\rangle$-module, and that the kernel is also $\operatorname{Ker}[(\nabla_{\partial_x}+\mathrm{Id}):M_{t_o}\to M_{t_o}]$. It is well-known that this kernel is 0 for a regular holonomic $M_{t_o}$ (this can be checked directly on cyclic $\mathbb{C}[x]\langle\partial_x\rangle$-modules and the general case follows by also considering



FIGURE 1

modules supported on points). Therefore, the only interesting module is

$$N := \mathscr{H}^0 p_+(\mathscr{E}^x \otimes M). \tag{12.1}$$

Information on the Levelt-Turrittin decomposition of $\mathscr{H}^\ell p_+(\mathscr{E}^x \otimes M)$ has been given in [**69, 70, 71**].

***Problem 12.2.*** — *To compute the Stokes filtration of $N$ at the origin of $\Delta$ in terms of the Stokes filtration of the analytic de Rham complex* $\mathrm{DR}^{\mathrm{an}} M$.

As noticed in [**27**], this problem can be translated into a problem of computing sectorial asymptotic expansions of integrals of the form

$$I(t) = \int_{\gamma_t} f(x,t) e^x dx,$$

where $f(x,t)$ is a multivalued solution on $(\mathbb{A}^1 \times \Delta) \smallsetminus S$ of a regular holonomic system of differential equation and $\gamma_t$ is a suitable family of cycles of the fibre $\mathbb{A}^1 \times \{t\}$ parametrized by $t$.

Let $\varpi : \widetilde{\Delta} \to \Delta$ be the real blowing up of the origin and set $S^1 = \varpi^{-1}(0)$. On $\widetilde{\Delta}$, we consider the sheaf $\mathscr{A}_{\widetilde{\Delta}}^{\mathrm{mod}\,0}$, and similarly, on $\mathbb{P}^1 \times \widetilde{\Delta}$, we consider the sheaf $\mathscr{A}_{\mathbb{P}^1 \times \widetilde{\Delta}}^{\mathrm{mod}\,0}$, where 0 now denotes the divisor $\mathbb{P}^1 \times \{0\}$. We denote by $\widetilde{p} : \mathbb{P}^1 \times \widetilde{\Delta} \to \widetilde{\Delta}$ the projection.

We will also consider the moderate de Rham complex $\mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^x \otimes M)$:

$$0 \longrightarrow \mathscr{A}_{\mathbb{P}^1 \times \widetilde{\Delta}}^{\mathrm{mod}\,0} \otimes_{\varpi^{-1}\mathscr{O}_{\mathbb{P}^1 \times \Delta}} M \xrightarrow{\nabla + dx} \mathscr{A}_{\mathbb{P}^1 \times \widetilde{\Delta}}^{\mathrm{mod}\,0} \otimes_{\varpi^{-1}\mathscr{O}_{\mathbb{P}^1 \times \Delta}} (\Omega^1_{\mathbb{P}^1 \times \Delta} \otimes M)$$
$$\xrightarrow{\nabla + dx} \mathscr{A}_{\mathbb{P}^1 \times \widetilde{\Delta}}^{\mathrm{mod}\,0} \otimes_{\varpi^{-1}\mathscr{O}_{\mathbb{P}^1 \times \Delta}} (\Omega^2_{\mathbb{P}^1 \times \Delta} \otimes M) \longrightarrow 0.$$

This is a complex on $\mathbb{P}^1 \times \widetilde{\Delta}$.

**12.b. Direct images and Riemann-Hilbert.** — It is known that the complex $\mathrm{DR}^{\mathrm{mod}\,0} N$ has cohomology in degree 0 at most (see Theorem 5.3). On the other hand, the complex $\mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^x \otimes M)$ only depends on the localized module $\mathscr{O}_{\mathbb{P}^1\times\Delta}[1/t] \otimes_{\mathscr{O}_{\mathbb{P}^1\times\Delta}} M$.

**Lemma 12.3.** — *There is a functorial morphism*

$$\mathscr{H}^0(\mathrm{DR}^{\mathrm{mod}\,0} N) \longrightarrow \mathscr{H}^1 \boldsymbol{R}\widetilde{p}_* \,\mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^x \otimes M), \tag{$12.3_{\leqslant 0}$}$$

*which is injective.*

*Proof.* — This is completely similar to Lemma 7.4. ☐

**Theorem 12.4.** — *The morphism of Lemma* 12.3 *is an isomorphism.*

The proof will be similar to that of Theorem 7.6, in the sense that it will involve a better topological understanding of the right-hand side in terms of $\mathrm{DR}^{\mathrm{an}} M$, that is, a solution to Problem 12.2, but the geometric situation is a little more complicated and uses more complex blowing-ups.

*Proof.* — We can assume that $M = \mathscr{O}_\Delta[1/t] \otimes_{\mathscr{O}_\Delta} M$, as the computation of $\mathrm{DR}^{\mathrm{mod}\,0}$ only uses the localized module (on $\mathbb{P}^1 \times \Delta$ or on $\Delta$). From the injectivity in Lemma 12.3, and as the theorem clearly holds away from $|t| = 0$, it is enough to check that the germs at $\theta \in S^1 = \partial\widetilde{\Delta}$ of both terms of ($12.3_{\leqslant 0}$) have the same dimension. It is then enough to prove the theorem after a ramification with respect to $t$ (coordinate of $\Delta$), so that we are reduced to assuming that, in the neighbourhood of $\mathbb{P}^1 \times \{0\}$, the irreducible components of the singular support $\overline{S}$ of $M$ are smooth and transverse to $\mathbb{P}^1 \times \{0\}$.

We can also localize $M$ along its singular support $\overline{S}$. The kernel and cokernel of the localization morphism are supported on $\overline{S}$, and the desired assertion is easy to check for these modules. We can therefore assume that $M$ is a meromorphic bundle along $\overline{S}$ with a flat connection having regular singularities. In particular, $M$ is a locally free $\mathscr{O}_{\mathbb{P}^1\times\Delta}(*\overline{S})$-module of finite rank (see [**76**, Prop. I.1.2.1]).

Let $e : X \to \mathbb{P}^1 \times \Delta$ be a sequence of point blowing up over $(\infty, 0)$, with exceptional divisor $E := e^{-1}(0, \infty)$, such that the strict transform of $\overline{S}$ intersects the pull-back of $(\mathbb{P}^1 \times \{0\}) \cup (\{\infty\} \times \Delta)$ only at smooth points of this pull-back. We can choose for $e$ a sequence of $n$ blowing-ups of the successive intersection points of the exceptional divisor with the strict transform of $\{\infty\} \times \Delta$. We set $D = e^{-1}(\mathbb{P}^1 \times \{0\})$, $D' = e^{-1}[(\mathbb{P}^1 \times \{0\}) \cup (\{\infty\} \times \Delta)]$. This is illustrated on Figure 2.

**Lemma 12.5.** — *The pull-back connection $e^+(\mathscr{E}^x \otimes M)$ is good except possibly at the intersection points of the strict transform of $\overline{S}$ with $D'$.*

*Proof.* — Indeed, $e^+M$ has regular singularities along its polar locus, and $e^+\mathscr{E}^x = \mathscr{E}^{1/y\circ e}$ is purely monomial (see Definition 9.8). At the intersection points of $\overline{S}$ with $D'$, the polar locus $\overline{S} \cup D'$ is not assumed to be a normal crossing divisor, which explains



FIGURE 2. The divisor $D$ is given by the thick lines. The vertical thick line is the strict transform of $\{t = 0\}$, the horizontal line is the strict transform of $\{x = \infty\}$ and the other thin lines are the strict transforms of the $\overline{S}_i$. Each dot is the center of a chart with coordinates $(u_k, v_k)$ $(k = 1, \dots, n)$ with $t \circ e = u_k v_k$ and $y \circ e = u_k^{k-1} v_k^k = v_k \cdot (t \circ e)^{k-1}$. The chart centered at the last dot has coordinates $(u'_n, v'_n)$ and $t \circ e = u'_n$, $y \circ e = u'^n_n v'_n = v'_n \cdot (t \circ e)^n$.

the restriction in the lemma. Of course, blowing-up these points sufficiently enough would lead to a good meromorphic connection, but we try to avoid these supplementary blowing-ups. ☐

Let $\widetilde{X}(D')$ be the real blowing up of the irreducible components of $D'$ (this notation is chosen to shorten the notation introduced in §8.a, which should be $\widetilde{X}(0, \infty, E_{i\in\{1,\dots,n\}})$, where $0$ (resp. $\infty$) denotes the strict transform of $\mathbb{P}^1 \times \{0\}$ (resp. $\{\infty\} \times \Delta$); this does not correspond to the real blow-up of the divisor $D'$). We denote by $\mathscr{A}^{\mathrm{mod}\,D'}_{\widetilde{X}(D')}$ the corresponding sheaf of functions (see §8.b). The morphism $e$ lifts to a morphism $\widetilde{e} : \widetilde{X}(D') \to \mathbb{P}^1 \times \widetilde{\Delta}$ and we have

$$\mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^x \otimes M) \simeq \boldsymbol{R}\widetilde{e}_*\, \mathrm{DR}^{\mathrm{mod}\,D'}_{\widetilde{X}(D')}[e^+(\mathscr{E}^x \otimes M)].$$

Indeed, this follows from Proposition 8.9 and from the isomorphism $e_+ e^+(\mathscr{E}^x \otimes M) = \mathscr{E}^x \otimes M$, which is a consequence of our assumption that $M$ is localized along $\mathbb{P}^1 \times \{0\}$.

Let us set $\mathscr{F}_{\leqslant 0} := \mathrm{DR}^{\mathrm{mod}\,D'}_{\widetilde{X}(D')}[e^+(\mathscr{E}^x \otimes M)]_{|\partial\widetilde{X}(D')}$. The proof of the theorem reduces to proving that, for any $\theta \in S^1 = \partial\widetilde{\Delta}$,

$$\dim \mathscr{H}^0(\mathrm{DR}^{\mathrm{mod}\,0} N)_\theta = \dim \boldsymbol{H}^1\big((\widetilde{p} \circ \widetilde{e})^{-1}(\theta), \mathscr{F}_{\leqslant 0}\big). \tag{12.6}$$

Indeed, if we denote by $\mathscr{F}_{\leqslant 0,\theta}$ the sheaf-theoretic restriction of $\mathscr{F}_{\leqslant 0}$ to $(\widetilde{p} \circ \widetilde{e})^{-1}(\theta)$, then, as $\widetilde{p} \circ \widetilde{e}$ is proper, we get

$$[\boldsymbol{R}^1 \widetilde{p}_*\, \mathrm{DR}^{\mathrm{mod}\,0}(\mathscr{E}^x \otimes M)]_\theta \simeq \boldsymbol{H}^1(\widetilde{X}(D')_\theta, \mathscr{F}_{\leqslant 0,\theta}).$$

Let us describe the inverse image by $\widetilde{p} \circ \widetilde{e} : \widetilde{X}(D') \to \widetilde{\Delta}$ of $\theta \in \partial\widetilde{\Delta} = S^1$. At a crossing point of index $k$ $(k = 1, \dots, n)$, $\widetilde{X}(D')$ is the product $(S^1 \times \mathbb{R}_+)^2$, with coordinates $(\alpha_k, |u_k|, \beta_k, |v_k|)$, and $\arg(t \circ e) = \theta$ is written $\alpha_k + \beta_k = \theta$. At the crossing point with coordinates $(u'_n, v'_n)$, we have coordinates $(\alpha'_n, |u'_n|, \beta'_n, |v'_n|)$ and $\arg(t \circ e) = \theta$ is written $\alpha'_n = \theta$. More globally, $\partial\widetilde{X}(D')_\theta := (\widetilde{p} \circ \widetilde{e})^{-1}(\theta)$ looks like a leaky pipe (see Figure 3, which has to be seen as lying above Figure 2), where the punctures (small black dots on the pipe for the visible ones, small circles for the ones which are behind) correspond to the intersection with the strict transforms of the $\overline{S}_i$.



FIGURE 3

*First step: hypercohomology of* $\mathscr{F}_{\leqslant 0,\theta}$

**Lemma 12.7.** — *Away from the punctures, the complex* $\mathscr{F}_{\leqslant 0,\theta}$ *has cohomology in degree* 0 *at most.*

*Proof.* — This is Corollary 11.22. ☐



FIGURE 4

Considering the growth of the functions $e^{1/u_k^{k-1}v_k^k}$ or $e^{1/u'_n v'^n_n}$, one obtains that, away from the punctures, the sheaf $\mathscr{F}_{\leqslant 0,\theta}$ is locally constant on a semi-open leaky half-pipe as in Figure 4, which is topologically like in Figure 5. Moreover, $\mathscr{F}_{\leqslant 0,\theta}$ is extended by 0 at the dashed boundary.



FIGURE 5

Let us denote by $\overline{S}_i$ $(i\in I)$ the components of $\overline{S}$ which contain the point $(0,\infty)$ and by $\overline{S}_j$ $(j\in J)$ the components which do not. Denoting as above by $y$ the coordinate on $\mathbb{P}^1$ at $\infty$ given by $y=1/x$, the local equation of each component $\overline{S}_i$ near $(0,\infty)$ takes the form $\mu_i(t)y=t^{q_i}$, with $\mu_i$ holomorphic and $\mu_i(0)\neq 0$. Let us set $\varphi_i(t)=\mu_i(t)/t^{q_i} \bmod \mathbb{C}\{t\}$. The punctures in the non-vertical part of Figure 4 or in the right part of Figure 5 correspond to the components $\overline{S}_i$ $(i\in I)$ for which $\varphi_i <_\theta 0$. We denote by $I_\theta\subset I$ the corresponding subset of $I$.

On the other hand, the punctures on the vertical part of Figure 4 or on the left part of Figure 5 correspond to the components $\overline{S}_j$ $(j\in J)$.

***Lemma 12.8.*** — *Near the punctures, $\mathscr{F}_{\leqslant 0,\theta}[1]$ is a perverse sheaf. It is zero if the puncture does not belong to the half-pipe of Figure 4, and the dimension of its vanishing cycle space at the puncture is the number of curves $\overline{S}_i$ $(i\in I_\theta$ or $i\in J)$ going through this puncture.*

*Proof.* — One checks that, blowing up the puncture and then taking the real blow-up of the components of the new normal crossing divisor, and then restricting to $\arg t=\theta$, amounts to change an neighbourhood of the puncture in Figure 5 (say) with a disc where the the puncture has been replaced by as many punctures as distinct tangent line of the curves $\overline{S}_i$ going through the original puncture, and the new sheaf $\widetilde{\mathscr{F}}_{\leqslant 0,\theta}$ remains locally constant away from the new punctures. By an easy induction, one reduces to the case where only one $\overline{S}_i$ goes through each puncture, and then the result is easy. ☐

***Corollary 12.9.*** — *$\boldsymbol{H}^k(\widetilde{X}(D')_\theta,\mathscr{F}_{\leqslant 0,\theta})$ is zero if $k\neq 1$ and $\dim \boldsymbol{H}^1(\widetilde{X}(D'),\mathscr{F}_{\leqslant 0,\theta}) = \operatorname{rk} M\cdot \#(J\cup I_\theta)$.*

*Proof.* — According to Lemma 12.8, this follows from Lemma 7.13. ☐

*End of the proof.* — According to [**70**], the possible exponential factors of $N$ (defined by (12.1)) are the $\varphi_i$ with $i\in I$. Denoting by $\phi_i \operatorname{DR} M$ the local system on $S_i$ of vanishing cycles of $\operatorname{DR} M$ along the function $f_i(t,y)=\mu_i(t)y-t^{q_i}$, we have

$$\dim \mathscr{H}^0(\mathrm{DR}^{\mathrm{rd}\,0} N)_\theta = \sum_{i\,|\,\varphi_i<_\theta 0} \operatorname{rk} \phi_i \operatorname{DR} M.$$

As $M$ is assumed to be a meromorphic bundle, we have $\operatorname{rk}\phi_i\operatorname{DR} M=\operatorname{rk} M$, so the previous formula reads

$$\dim \mathscr{H}^0(\mathrm{DR}^{\mathrm{rd}\,0} N)_\theta = \operatorname{rk} M\cdot \# I_\theta.$$

We now use

$$\dim \mathscr{H}^0(\mathrm{DR}^{\mathrm{mod}\,0} N)_\theta = \dim \mathscr{H}^0(\mathrm{DR}^{\mathrm{rd}\,0} N)_\theta + \dim \psi_t N,$$

where $\psi_t N$ denote the moderate nearby cycles of $N$ (computed with the Kashiwara-Malgrange $V$-filtration, see [**49**]). Arguing as in [**70**], we compute them as the direct image of $\psi_t(\mathscr{E}^x \otimes M)$ by $\mathbb{P}^1 \times \{0\} \to \{0\}$. By the same argument, this is computed as the direct image of $\psi_{t\circ e} e^+(\mathscr{E}^x \otimes M)$. Arguing as in [**73**, Lemme III.4.5.10(2)], this is supported on the vertical part of $D$. Still using the argument of [**73**, Lemme III.4.5.10], this is finally the direct image of $\psi_t(M) \otimes \mathscr{E}^x$, whose dimension is that of the rank of the Fourier transform of $\psi_t(M)$, regarded as a $\mathbb{C}[x]\langle\partial_x\rangle$-module, that is, the dimension of the vanishing cycles of $\psi_t(M)$ (see [**49**] for such a formula, compare also with Proposition 7.11).

Let us compute this dimension at a point $x_o \neq \infty$. Since $M$ is assumed to be a meromorphic connection, one checks that $\dim \phi_{x-x_o}\psi_t(M)$ is the number of components of $\overline{S}$ going through $x_o$. The sum $\sum_{x_o} \dim \phi_{x-x_o}\psi_t(M)$ is then equal to $\#J$, by definition of $J$.

Summarizing, we get

$$\dim \psi_t N = \operatorname{rk} M \cdot \#J.$$

This concludes the proof of $(12.3_{\leqslant 0})$. □

# LECTURE 13

# THE RIEMANN-HILBERT CORRESPONDENCE FOR GOOD MEROMORPHIC CONNECTIONS

**Summary.** This lecture is similar to Lecture 10, but we now assume that $D$ is a divisor with normal crossings. The new point is that we have to deal with non-Hausdorff étale spaces, and we need to use the level structure to prove the local essential surjectivity of the Riemann-Hilbert functor. We also apply the fundamental results of K. Kedlaya [**37, 38**] and T. Mochizuki [**60, 65, 64**] to show that the Hermitian dual of a holonomic $\mathscr{D}$-module is holonomic, generalizing the main result of M. Kashiwara [**33**] to possibly irregular holonomic $\mathscr{D}$-modules and the result of Lecture 6 to higer dimension.

**13.a. The Riemann-Hilbert functor.** — We consider the following setting:

- $X$ is a complex manifold and $D$ is a divisor with normal crossings in $X$, with smooth components $D_j$ ($j\in J$), and $X^*:=X\smallsetminus D$,
- $j:X^*\hookrightarrow X$ and $\widetilde{\jmath}:X^*\hookrightarrow \widetilde{X}(D_{j\in J})$ denote the open inclusions, and $i:D\hookrightarrow X$ and $\widetilde{\imath}:\partial\widetilde{X}(D_{j\in J})\hookrightarrow \widetilde{X}(D_{j\in J})$ denote the closed inclusions,
- the ordered sheaf $\mathfrak{I}$ on $\partial\widetilde{X}(D_{j\in J})$ is as in Definitions 9.4 and 9.5.

In this lecture, we consider germs along $D$ of meromorphic connections with poles on $D$ at most, which are locally good in the sense of Definition 11.9 (in particular, we assume the local existence of a good lattice, but we do not care of the global existence of such a lattice, which could be proved by using a good Deligne-Malgrange lattice, see [**65**]).

The arguments of §10.b apply exactly in the same way here, so Definition 10.3 will be used below. Similarly to Lemma 10.4 we also have:

***Lemma 13.1.*** — *If $\mathscr{M}$ is a good meromorphic connection with poles along $D$ (see Definition 11.9) with associated stratified $\mathfrak{I}$-covering $\widetilde{\Sigma}$, then $\mathrm{RH}(\mathscr{M})=(\mathscr{L},\mathscr{L}_{\leqslant})$ is a good Stokes-filtered local system on $\partial\widetilde{X}$ (see Definition 9.17) with associated stratified $\mathfrak{I}$-covering $\widetilde{\Sigma}$, that we denote by $(\mathscr{L},\mathscr{L}_{\bullet})$.*

*Proof*. — Same proof as for Lemma 10.4, if we use the Hukuhara-Turrittin-Majima theorem 11.20 instead of the Hukuhara-Turrittin-Sibuya theorem. □

***Theorem 13.2***. — *Let $\widetilde{\Sigma}$ be a good stratified $\mathfrak{I}$-covering with respect to the (pull-back to $\partial\widetilde{X}(D)$ of the) natural stratification of $D$. The Riemann-Hilbert functor induces an equivalence between the category of germs of good meromorphic connections along $D$ with stratified $\mathfrak{I}$-covering (see Remark* 11.11*) contained in $\widetilde{\Sigma}$ and the category of good Stokes-filtered $\mathbb{C}$-local systems on $\partial\widetilde{X}$ with stratified $\mathfrak{I}$-covering (see Definition* 9.17*) contained in $\widetilde{\Sigma}$.*

**13.b. Local theory.** — In this section, the setting and the notation are as in Lecture 11. In particular, $X = \Delta^\ell \times \Delta^{n-\ell}$ and we may shrink $X$ when necessary. We also set $D = \{t_1 \cdots t_\ell = 0\}$, and $\varpi : \widetilde{X} := \widetilde{X}(D) \to X$ denotes the real blow-up of the components of $D$ (see §8.a), so that in particular $\varpi^{-1}(0) \simeq (S^1)^\ell$. Let $\Phi \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ be a good finite set of exponential factors (see Definition 9.12). We will prove the theorem for germs at 0 of good meromorphic connections and germs along $\varpi^{-1}(0)$ of good Stokes-filtered local systems.

As in the cases treated before (§5.c and 10.e), we will use the corresponding variant of the Hukuhara-Turrittin theorem, which is the Hukuhara-Turrittin-Majima theorem 11.20. This gives the analogue of Lemma 10.8. It then remains to show the analogue of Lemma 5.12 (compatibility with $\mathscr{H}\!om$) to conclude the proof of the full faithfulness of the Riemann-Hilbert functor. The proof is done similarly, and the goodness condition for $\Phi\cup\Phi'$ makes easy to show the property that (taking notation of the proof of Lemma 5.12) $e^{\varphi-\varphi'}u_{\varphi,\varphi'}$ has moderate growth in some neighbourhood of $\theta_o \in \varpi^{-1}(0)$ if and only if $\varphi \leqslant_{\theta_o} \varphi'$.

For the essential surjectivity, let us consider a Stokes-filtered local system $(\mathscr{L}, \mathscr{L}_\bullet)$ on $\partial\widetilde{X}$ whose associated $\mathfrak{I}$-covering over $\varpi^{-1}(0_\ell \times \Delta^{n-\ell})$ is trivial and contained in $\Phi \times \Delta^{n-\ell}$ (in particular, we consider the non-ramified case).

***Proposition 13.3***. — *Under these assumptions, there exists a germ at $0 \in X$ of good meromorphic connection $\mathscr{M}$ (in the sense of Definition* 11.9*) such that, for any local section $\varphi$ of $\mathfrak{I}$, $\mathrm{DR}(e^\varphi \mathscr{A}^{\mathrm{mod}\,D}_{\partial\widetilde{X}} \otimes \varpi^{-1}\mathscr{M}) \simeq \mathscr{L}_{\leqslant\varphi}$ in a way compatible with the filtration.*

As in the cases treated before (§5.c and 10.e), the main point is the *Malgrange-Sibuya theorem*. A detailed proof of this theorem will be given in §13.c.

***Theorem 13.4* (Malgrange-Sibuya in dimension $\geqslant 2$).** — *The image of*

$$H^1\big((S^1)^\ell, \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{A}_{\partial\widetilde{X}})\big) \longrightarrow H^1\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})\big)$$

*is the identity.*

If one uses the multi-dimensional analogue of the theorem of Borel-Ritt due to Majima (see [**73**, Prop. II.1.1.16]), which gives the existence of an exact sequence

$$H^0\big((S^1)^\ell, \mathrm{GL}_d(\varpi^{-1}\mathscr{O}_{\widehat{D}})\big) = \mathrm{GL}_d(\mathscr{O}_{\widehat{D},0}) \longrightarrow H^1\big((S^1)^\ell, \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{A}_{\partial\widetilde{X}})\big) \\ \longrightarrow H^1\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})\big),$$

it follows that for each class $\gamma \in H^1\big((S^1)^\ell, \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{A}_{\partial\widetilde{X}})\big)$, there exist $\widehat{g} \in \mathrm{GL}_d(\mathscr{O}_{\widehat{D},0})$, a covering $(U_i)$ of $(S^1)^\ell$, and liftings $g_i \in \Gamma\big(U_i, \mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})\big)$ of $\widehat{g}$ on $U_i$, such that $\gamma$ is represented by the cocycle $(g_i g_j^{-1})$.

In order to apply it in the normal crossing case, we need to be sure that all differences $\varphi - \psi$, where $\varphi, \psi \in \Phi$, have poles along $D$ exactly. This does not hold in general, but can be achieved (up to changing $D$) at each step of the level structure considered in §9.16.

*Proof of Proposition* 13.3. — The proof of Proposition 13.3 will proceed by induction on the pairs $(\ell, \boldsymbol{m}(\Phi))$ (see Remark 9.13(5) for the definition of $\boldsymbol{m}(\Phi)$) through the level structure, where $\ell$ denotes the codimension of the stratum of $D$ to which belongs the origin.

For each $\ell \geqslant 1$, the case $\boldsymbol{m} = 0$ corresponds (up to a twist, see Remark 9.13(5)), to the case where the Stokes filtration is trivial, and in such a case the regular meromorphic connection associated with the local system $\mathscr{L}$ fulfills the conditions of Proposition 13.3.

We fix $\ell$ and $\boldsymbol{m} = \boldsymbol{m}(\Phi)$ and we assume that Proposition 13.3 holds for any Stokes-filtered local system $(\mathscr{L}', \mathscr{L}'_\bullet)$ with associated $\mathcal{I}$-covering over $\varpi^{-1}(0_\ell \times \Delta^{n-\ell})$ contained in $\Phi' \times \Delta^{n-\ell}$, with a good $\Phi' \subset \mathscr{O}_{X,0}(*D)/\mathscr{O}_{X,0}$ satisfying $\boldsymbol{m}(\Phi') < \boldsymbol{m}$ (with respect to the partial order of $\mathbb{N}^\ell$), and also for any pair $(\ell', \boldsymbol{m}')$ with $\ell' < \ell$, and we will prove it for the pair $(\ell, \boldsymbol{m})$. We will therefore assume that $\boldsymbol{m} > 0$ (in $\mathbb{N}^\ell$). We also fix some element $\varphi_o \in \Phi$ and we denote by $\boldsymbol{\ell}_{\varphi_o} = \boldsymbol{\ell} =\in \mathbb{N}^\ell$ the submaximum of $\Phi - \varphi_o$. By twisting we may assume for simplicity that $\varphi_o = 0$.

Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be a good Stokes-filtered local system on $(S^1)^\ell$ with set of exponential factors contained in $\Phi$. According to Proposition 9.23 (applied with $\boldsymbol{\ell} = \boldsymbol{\ell}_{\varphi_o}$) and to Remark 9.13(6), this Stokes-filtered local system induces a Stokes-filtered local system $(\mathscr{L}, \mathscr{L}_{[\bullet]_{\boldsymbol{\ell}}})$ of level $\geqslant \boldsymbol{\ell}$, such that each $\big(\mathrm{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L}, (\mathrm{gr}_{[\varphi]_{\boldsymbol{\ell}}}\mathscr{L})_\bullet\big)$ is a Stokes-filtered local system to which we can apply the inductive assumption.

If $[\varphi]_{\boldsymbol{\ell}} \in \Phi(\boldsymbol{\ell}) = \mathrm{image}(\Phi \to \mathcal{P}_\ell(\boldsymbol{\ell}))$, it takes the form $ct^{-\boldsymbol{m}} \bmod (t^{-\boldsymbol{\ell}}\mathbb{C}[t])$ for some $c \in \mathbb{C}$.

By induction, we get germs of meromorphic connections $\mathscr{M}_c$ ($c \in \mathbb{C}$) corresponding to $\big(\mathrm{gr}_{[ct^{-\boldsymbol{m}}]_{\boldsymbol{\ell}}}\mathscr{L}, (\mathrm{gr}_{[ct^{-\boldsymbol{m}}]_{\boldsymbol{\ell}}}\mathscr{L})_\bullet\big)$. Each $\mathscr{M}_c$ is good (by the inductive assumption) and its set of exponential factors is contained in the set of $\varphi \in \Phi$ of the form $\varphi \equiv ct^{-\boldsymbol{m}} \bmod (t^{-\boldsymbol{\ell}}\mathbb{C}[t])$.

We will now construct $\mathscr{M}$ from $\mathscr{M}_{\boldsymbol{\ell}} := \bigoplus_{c\in\mathbb{C}}\mathscr{M}_c$, by considering the Stokes-filtered local system $(\mathscr{L},\mathscr{L}_{[\bullet]_{\boldsymbol{\ell}}})$ of level $\geqslant \boldsymbol{\ell}$, whose corresponding graded object is the Stokes-filtered local system $(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L},(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet)$ graded at the level $\geqslant \boldsymbol{\ell}$.

**Lemma 13.5.** — *We have a natural isomorphism*

$$\mathscr{E}nd_{\mathscr{D}}^{\mathrm{mod}\,D}(\mathscr{M}_{\boldsymbol{\ell}}) \simeq \mathscr{E}nd(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})_{\leqslant 0}.$$

*Proof.* — Since by definition $\mathscr{H}^0\,\mathrm{DR}_{\mathrm{j\acute{e}t}}^{\mathrm{mod}\,D}(\mathscr{M}_{\boldsymbol{\ell}}) \simeq (\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L},(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet)$, the assertion is proved as in Lemma 10.18, using the compatibility of the Riemann-Hilbert functor with $\mathscr{H}om$ mentioned above. ☐

*The case where $m_i > 0$ for all $i = 1,\dots,\ell$*

**Lemma 13.6.** — *In this case, the isomorphism of Lemma* 13.5 *induces an inclusion* $\mathscr{E}nd(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})_{<_{[\bullet]_{\boldsymbol{\ell}}}0} \subset \mathscr{E}nd_{\mathscr{D}}^{\mathrm{rd}\,D}(\mathscr{M}_{\boldsymbol{\ell}})$.

*Proof.* — Recall that the left-hand side consists of those endomorphisms $\lambda$ which satisfy $\mathrm{gr}_{[\varphi]_{\boldsymbol{\ell}}}\lambda = 0$ for each $\varphi$. Note that the condition $\varphi <_{[\bullet]_{\boldsymbol{\ell}}} \psi$ near $\theta \in (S^1)^\ell$ implies that the $\boldsymbol{m}$-dominant part of $\varphi$ and $\psi$ are distinct, and so, by our assumption on $\boldsymbol{m}$, $\varphi-\psi$ has a pole along each component of $D$ and $\varphi < \psi$ near $\theta$, so $e^{\varphi-\psi}$ has rapid decay. This applies to the $(\varphi,\psi)$ components of $\lambda$, expressed as in Lemma 5.12. ☐

Let us finish the proof of Proposition 13.3 in this case. According to Remark 9.24, given the Stokes-filtered local system $(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L},(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet)$ graded at the level $\geqslant \boldsymbol{\ell}$, the Stokes-filtered local system $(\mathscr{L},\mathscr{L}_\bullet)$ determines (and is determined by) a class $\gamma$ in the pointed set $H^1\big((S^1)^\ell,\mathscr{A}ut^{<_{[\bullet]_{\boldsymbol{\ell}}}0}(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})\big)$, hence a class $\gamma$ in $H^1\big((S^1)^\ell,\mathscr{A}ut_{\mathscr{D}}^{\mathrm{rd}\,D}(\mathscr{M}_{\boldsymbol{\ell}})\big)$, after Lemma 13.6.

The germ $\mathscr{M}_{\boldsymbol{\ell}}$ is a free $\mathscr{O}_{X,0}(*D)$-module with connection $\nabla_{\boldsymbol{\ell}}$. With respect to some basis of $\mathscr{M}_{\boldsymbol{\ell}}$, the class $\gamma$ becomes a class in $H^1\big((S^1)^\ell,\mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{A}_{\partial\widetilde{X}})\big)$. By the Malgrange-Sibuya theorem 13.4, this class is a coboundary of $\mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})$ on $(S^1)^\ell$. Let $(U_i)$ be a covering of $(S^1)^\ell$ on which this coboundary is defined by sections $g_i \in \Gamma\big(U_i,\mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})\big)$ with $\widehat{g}_i = \widehat{g} \in \mathrm{GL}_d(\mathscr{O}_{\widehat{D},0})$ for all $i$. On $U_i$ we twist the connection on $\mathscr{M}_{\boldsymbol{\ell}}$ by setting $\nabla_i = g_i^{-1}\nabla_{\boldsymbol{\ell}} g_i$. Since $g_i g_j^{-1} = \gamma_{ij}$ is $\nabla_{\boldsymbol{\ell}}$-flat on $U_i\cap U_j$, the $\nabla_i$ glue together to define a new connection $\nabla$ on the free $\mathscr{O}_{X,0}(*D)$-module $\mathscr{M}_{\boldsymbol{\ell}}$, that we now denote by $(\mathscr{M},\nabla)$. By construction, $\mathscr{H}^0\,\mathrm{DR}_{\mathrm{j\acute{e}t}}^{\mathrm{mod}\,D}(\mathscr{M})$ is the Stokes-filtered local system determined by $(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L},(\mathrm{gr}_{\boldsymbol{\ell}}\mathscr{L})_\bullet)$ and the class $\gamma$, hence is isomorphic to $(\mathscr{L},\mathscr{L}_\bullet)$.

*The case where $m_i = 0$ for some $i = 1,\dots,\ell$.* — In this case, $(\mathscr{L},\mathscr{L}_\bullet)$ is partially regular along $D$ and we shall use the equivalence of Proposition 9.36. We set $L = L'\cup L''$ with $m_i = 0$ if and only if $i \in L''$, and we have $\Phi \subset \mathscr{O}_{X,0}(*D(L'))/\mathscr{O}_{X,0}$. By Proposition 9.36, giving $(\mathscr{L},\mathscr{L}_\bullet)$ is equivalent to giving a Stokes-filtered local system $(\mathscr{L}',\mathscr{L}'_\bullet)$ on $\partial\widetilde{X}'$ together with commuting automorphisms $T_k$, $k \in L''$. By induction on $\ell$, there exists a free $\mathscr{O}_{X,0}(*D(L'))$-module $\mathscr{M}'$ with flat connection $\nabla'$ whose associated Stokes-filtered local system is $(\mathscr{L}',\mathscr{L}'_\bullet)$. Moreover, there exist commuting endomorphisms $C_k$ of $(\mathscr{L}',\mathscr{L}'_\bullet)$ such that $\exp(-2\pi C_k) = T_k$ (represent the local system $\mathscr{L}'$ as a vector space $V'$ equipped with automorphisms $T'_j$, $j \in L'$; then $T_k$ are automorphisms of $V'$ which commute with the $T'_j$, and any choice $C_k$ of a logarithm of $-(2\pi i)^{-1}T_k$ also commutes with $T'_j$ and defines an endomorphism of $\mathscr{L}'$; similarly, this endomorphism is filtered with respect to the Stokes filtration $\mathscr{L}'_\bullet$). By the full faithfulness of the Riemann-Hilbert correspondence, the commuting endomorphisms $C_k$ of $(\mathscr{L}',\mathscr{L}'_\bullet)$ define commuting endomorphisms $C_k$ of $(\mathscr{M}',\nabla')$. The free $\mathscr{O}_{X,0}(*D)$-module $\mathscr{M}'(*D'')$ can be equipped with the flat connection $\nabla := \nabla' + \sum_{k\in L''} C_k\, dt_k/t_k$. Then one checks that the Stokes-filtered local system associated to $(\mathscr{M},\nabla)$ on $\partial\widetilde{X}$ is isomorphic to $(\mathscr{L},\mathscr{L}_\bullet)$. ☐

*Proof of Theorem* 13.2 *for germs.* — It now remains to treat the local reconstruction (Proposition 13.3) in the ramified case. The data of a possibly ramified $(\mathscr{L},\mathscr{L}_\bullet)$ is equivalent to that of a non-ramified one on a covering $(S^1)^\ell_{\boldsymbol{d}}$ which is stable with respect to the Galois action of the covering. The same property holds for germs of meromorphic connections. By the full faithfulness of the Riemann-Hilbert functor, the Galois action on a Stokes-filtered local system is lifted in a unique way as a Galois action on the reconstructed connection after ramification, giving rise to a meromorphic connection before ramification, whose associated Stokes-filtered local system is isomorphic to $(\mathscr{L},\mathscr{L}_\bullet)$. ☐

*Proof of Theorem* 13.2 *in the global setting.* — Due to the local full faithfulness of the Riemann-Hilbert functor, one gets at the same time the global full faithfulness and the global essential surjectivity by lifting in a unique way the local gluing morphisms, which remain therefore gluing morphisms (i.e., the cocycle condition remains satisfied after lifting). ☐

**13.c. Proof of the Malgrange-Sibuya theorem 13.4.** — Since the proof given in [**73**, §II.1.2] contains a small mistake[(1)], we give a detailed proof here.

We denote by $\mathscr{E}_{\widetilde{X}}$ the sheaf of $C^\infty$ functions on $\widetilde{X}$ and by $\mathscr{E}_{|\partial\widetilde{X}}$ its sheaf-theoretic restriction to $\partial\widetilde{X}$. We can then define the subsheaf $\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}}$ of $C^\infty$ functions with rapid decay and the quotient sheaf is that of $C^\infty$ formal functions along $\partial\widetilde{X}$. We note that a local section of $\mathrm{M}_d(\mathscr{E}_{|\partial\widetilde{X}})$ (matrices of size $d$ with entries in $\mathscr{E}_{|\partial\widetilde{X}}$) is a section of $\mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}})$ if and only if its image in $\mathrm{M}_d(\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}})$ belongs to $\mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}})$: indeed, given a matrix in $\mathrm{M}_d(\mathscr{E}_{|\partial\widetilde{X}})$, the image in $\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}}$ of its determinant is the determinant of its image in $\mathrm{M}_d(\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}})$; use now that the values of a local section of $\mathscr{E}_{|\partial\widetilde{X}}$ at $\partial\widetilde{X}$ are also the values of its image in $\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}}$.

---

(1) whose correction is available at `http://www.math.polytechnique.fr/~sabbah/sabbah_ast_263_err.pdf`

***Lemma 13.7.*** — *We have* $H^1\big((S^1)^\ell, \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{E}_{|\partial\widetilde{X}})\big) = \mathrm{Id}$*, where we have set as above* $\mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{E}_{|\partial\widetilde{X}}) := \mathrm{Id} + \mathrm{M}_d^{\mathrm{rd}\,D}(\mathscr{E}_{|\partial\widetilde{X}})$.

*Proof.* — By definition, we have an exact sequence of groups

$$\mathrm{Id} \longrightarrow \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{E}_{|\partial\widetilde{X}}) \longrightarrow \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}}) \longrightarrow \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}}) \longrightarrow \mathrm{Id}\,.$$

We first show that $H^0\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}})\big) \to H^0\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}})\big)$ is onto. Locally, a section of the right-hand term can be lifted. Using a partition of unity, we lift it globally as a section of $\mathrm{M}_d(\mathscr{E}_{|\partial\widetilde{X}})$, and by the remark above, it is a section of $\mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}})$.

It remains thus to show that $H^1\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}})\big) \to H^1\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}}/\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}})\big)$ is injective, and since $\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}} \subset t_1\cdots t_\ell \mathscr{E}_{|\partial\widetilde{X}}$, it is enough to show a similar assertion of the restriction map $H^1\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{E}_{|\partial\widetilde{X}})\big) \to H^1\big((S^1)^\ell, \mathrm{GL}_d(\mathscr{E}_{\partial\widetilde{X}})\big)$, where $\mathscr{E}_{\partial\widetilde{X}} = \mathscr{E}_{|\partial\widetilde{X}}/t_1\cdots t_d\mathscr{E}_{|\partial\widetilde{X}}$ is the sheaf of $C^\infty$ functions on $\partial\widetilde{X}$.

Using the interpretation of an element of $H^1$ as giving an isomorphism class of vector bundle, we are reduced to showing that, given a $C^\infty$ vector bundle in the neighbourhood of $(S^1)^\ell$ whose restriction to $(S^1)^\ell$ is trivializable, it is trivializable in some (possibly smaller) neighbourhood of $(S^1)^\ell$. For that purpose, it is enough to prove that any global section of the restriction can be lifted to a global section of the original bundle in some neighbourhood of $(S^1)^\ell$, because a lift of a basis of global sections will remain a basis of sections in some neighbourhood of $(S^1)^\ell$. Now, such a lifting property for a global section can be done locally on $(S^1)^\ell$ and glued with a partition of unity. □

Let $\alpha$ be a class in $H^1\big((S^1)^\ell, \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{A}_{\partial\widetilde{X}})\big)$ represented by a cocycle $(\alpha_{ij})$ on some covering $\mathscr{U} = (U_i)$ of $(S^1)^\ell$. According to the previous lemma, $H^1\big(\mathscr{U}, \mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{E}_{|\partial\widetilde{X}})\big) = 0$ for any covering $\mathscr{U}$ of $(S^1)^\ell$, and therefore $\alpha_{ij} = \beta_i^{-1}\beta_j$, where $\beta_i$ is a section over $U_i$ of $\mathrm{GL}_d^{\mathrm{rd}\,D}(\mathscr{E}_{|\partial\widetilde{X}})$.

The operator $\overline{\partial}$ is well defined on $\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}}$. We set

$$\gamma_i = \overline{\partial}\beta_i \cdot \beta_i^{-1}.$$

Then $\gamma_i = \gamma_j$ on $U_i \cap U_j$ and the $\gamma_i$ glue together as a matrix $\gamma$ of 1-forms with entries in $\mathscr{E}^{\mathrm{rd}\,D}_{X|D} = \varpi_*\mathscr{E}^{\mathrm{rd}\,D}_{|\partial\widetilde{X}}$, and of type $(0,1)$. Moreover, the $\gamma_i$ (hence $\gamma$) satisfy

$$\overline{\partial}\gamma_i + \gamma_i \wedge \gamma_i = 0$$

because this equality is already satisfied away from $\partial\widetilde{X}$. For $\gamma$, this equality is read on $X$.

***Lemma 13.8.*** — *There exists a neighbourhood of* $0 \in X$ *on which the equation* $\overline{\partial}\varphi = -\varphi\cdot\gamma$ *has a solution* $\varphi$ *which is a section of* $\mathrm{GL}_d(\mathscr{E}_X)$.

*Proof.* — This is Theorem 1 in [**43**, Chap. X]. ☐

Then for each $i$ one has $\overline{\partial}(\varphi\beta_i) = 0$, so that $\varphi\beta_i$ is a section on $U_i$ of $\mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})$ and $\alpha_{ij} = (\varphi\beta_i)^{-1}\cdot(\varphi\beta_j)$, in other words, the image of $\alpha$ in $H^1\big(\mathscr{U}, \mathrm{GL}_d(\mathscr{A}_{\partial\widetilde{X}})\big)$ is the identity. ☐

**13.d. Application to Hermitian duality of holonomic $\mathscr{D}$-modules.** — The Riemann-Hilbert correspondence for good meromorphic connections, as stated in Theorem 13.2, together with the fundamental results of K. Kedlaya and T. Mochizuki, allows one to give a complete answer to a question asked by M. Kashiwara in [**33**], namely, to prove that the Hermitian dual $C_X\mathscr{M}$ of a holonomic $\mathscr{D}_X$-module is still holonomic. This application has also been considered in [**64**].

Recall the notation of Lecture 6, but now in arbitrary dimension. We now denote by $\mathscr{D}_X$ the sheaf of holomorphic linear differential operators on a complex manifold $X$ and by $\mathfrak{Db}_X$ the sheaf of distributions on the underlying $C^\infty$ manifold, which is a left $\mathscr{D}_X \otimes_{\mathbb{C}} \mathscr{D}_{\overline{X}}$-module. The Hermitian dual $C_X\mathscr{M}$ of $\mathscr{M}$ is the $\mathscr{D}_X$-module $\mathscr{H}om_{\mathscr{D}_{\overline{X}}}(\overline{\mathscr{M}}, \mathfrak{Db}_X)$.

***Theorem 13.9.*** — *If* $\mathscr{M}$ *is holonomic, then so is* $C_X\mathscr{M}$, *and* $\mathscr{E}xt^k_{\mathscr{D}_{\overline{X}}}(\overline{\mathscr{M}}, \mathfrak{Db}_X) = 0$ *for* $k > 0$.

*Sketch of the proof.* — It is done in many steps, and is very similar to that in dimension one (see Theorem 6.4), except for the goodness property, which is now essential:

(1) One first reduces (see [**33**] see also [**73**]) to the case where $\mathscr{M}$ is a meromorphic bundle with connection along a divisor $D$, and to proving that

- $C_X^{\mathrm{mod}\,D}\mathscr{M} := \mathscr{H}om_{\mathscr{D}_{\overline{X}}}(\mathscr{M}, \mathfrak{Db}_X(*D))$ is a meromorphic bundle with connection,
- $\mathscr{E}xt^k_{\mathscr{D}_{\overline{X}}}(\mathscr{M}, \mathfrak{Db}_X(*D)) = 0$ for $k > 0$.

(2) The problem is local on $X$ and one can apply the resolution of singularities in the neighbourhood of a point of $D$ to assume that $D$ has normal crossings. The problem remains local, and one can apply the result of K. Kedlaya [**38**] (when $\dim X = 2$, one refers to [**37**]; in the algebraic setting and $\dim X = 2$, one can use [**60**], and [**65**] for $\dim X \geqslant 3$, see Remark 11.10) to reduce to the case where $\mathscr{M}$ is a good meromorphic bundle with connection. This reduction is of course essential. We can also assume that it has no ramification.

(3) As in dimension one, one reduces to proving a similar result on the real blow-up space $\widetilde{X}(D)$, by replacing $\mathscr{M}$ with $\widetilde{\mathscr{M}} = \mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D} \otimes_{\varpi^{-1}\mathscr{O}_X} \varpi^{-1}\mathscr{M}$ and $\mathfrak{Db}_X^{\mathrm{mod}\,D}$ with $\mathfrak{Db}_{\widetilde{X}}^{\mathrm{mod}\,D}$. Now, Theorem 11.20 asserts that $\widetilde{\mathscr{M}}$ is of good Hukuhara-Turrittin type (a definition analogous to Definition 6.1, supplemented of the goodness assumption).

Recall that an important point here is the existence of a good lattice proved in [**64**], see Remark 11.17.

(4) One now proves as in [**73**, Prop. II.3.2.6] the vanishing of the $\mathscr{E}xt^k$ for $k > 0$, and that $C_{\widetilde{X}}^{\mathrm{mod}\,D}(\widetilde{\mathscr{M}})$ is a locally free $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$-module with flat connection of Hukuhara-Turrittin type.

(5) We can now repeat the arguments of Proposition 6.2 and Corollary 6.3, by using Theorem 13.2 instead of Theorem 5.7, to prove that $C_{\widetilde{X}}^{\mathrm{mod}\,D}(\widetilde{\mathscr{M}}) = \mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D} \otimes_{\varpi^{-1}\mathscr{O}_X} \varpi^{-1}\mathscr{N}$ for some meromorphic bundle with connection $\mathscr{N}$, and thus $\mathscr{N} = C_X^{\mathrm{mod}\,D}\mathscr{M}$. $\square$

# LECTURE 14

# NEARBY CYCLES OF STOKES-FILTERED LOCAL SYSTEMS

**Summary.** In this lecture, we define a nearby cycle functor for a good Stokes-filtered local system on $(X, D)$, relative to a holomorphic function whose zero set is contained in the normal crossing divisor $D$. We then show that the Riemann-Hilbert correspondence of Lecture 13 is compatible with taking nearby cycles, in a suitable sense.

**14.a. Preliminaries.** — Let $g: X' \to X$ be a holomorphic map between complex manifolds $X'$ and $X$. We assume that $X$ and $X'$ are equipped with normal crossing divisors $D$ and $D'$ with smooth components $D_{j\in J}$ and $D'_{j'\in J'}$, and that

(1) $D' = g^{-1}(D)$,
(2) $g: X' \smallsetminus D' \to X \smallsetminus D$ is smooth.

Let $\varpi: \widetilde{X}(D_{j\in J}) \to X$ (resp. $\varpi': \widetilde{X}'(D'_{j'\in J'}) \to X'$) be the real blowing up of the components $D_{j\in J}$ in $X$ (resp. $D'_{j'\in J'}$ in $X'$). There exists a lifting $\widetilde{g}: \widetilde{X}' \to \widetilde{X}$ of $g$ (see §8.a) such that the following diagram commutes:

$$\begin{tikzcd}
\widetilde{X}' \arrow[r, "\varpi'"] \arrow[d, "\widetilde{g}"'] & X' \arrow[d, "g"] \\
\widetilde{X} \arrow[r, "\varpi"] & X
\end{tikzcd}$$

Notice that $\partial\widetilde{X}' = \widetilde{g}^{-1}(\partial\widetilde{X})$.

***Remark 14.1*** **(Direct images of local systems).** — We set $X^* = X \smallsetminus D$, $X'^* = X' \smallsetminus D'$, and we denote by $\widetilde{\jmath}$ (resp. $\widetilde{\jmath}'$) the inclusion $X^* \hookrightarrow \widetilde{X}$ (resp. $X'^* \hookrightarrow \widetilde{X}'$), and by $\widetilde{\imath}$ (resp. $\widetilde{\imath}'$) the inclusion $\partial\widetilde{X} \hookrightarrow \widetilde{X}$ (resp. $\partial\widetilde{X}' \hookrightarrow \widetilde{X}'$). We assume here that $g$ is proper.

Let $\mathscr{F}'^*$ be a local system on $X'^*$. Then, since $g: X'^* \to X^*$ is smooth and proper, each $\boldsymbol{R}^k g_*\mathscr{F}'^*$ $(k \geqslant 0)$ is a local system on $X^*$ and, after Corollary 8.3, $\mathscr{L}' := \widetilde{\imath}'^{-1}\widetilde{\jmath}'_*\mathscr{F}'^*$ and $\mathscr{L}^k := \widetilde{\imath}^{-1}\widetilde{\jmath}_*\boldsymbol{R}^k g_*\mathscr{F}'^*$ $(k \geqslant 0)$ are local systems on $\partial\widetilde{X}'$ and $\partial\widetilde{X}$ respectively. We claim that

$$\forall\, k \geqslant 0, \quad \boldsymbol{R}^k\widetilde{g}_*\mathscr{L}' = \mathscr{L}^k. \tag{14.1 $*$}$$

Indeed, we have

$$\begin{aligned}\boldsymbol{R}\widetilde{g}_*\mathscr{L}' &= \boldsymbol{R}\widetilde{g}_*\widetilde{\imath}'^{-1}\widetilde{\jmath}'_*\mathscr{F}'^* \\ &= \widetilde{\imath}^{-1}\boldsymbol{R}\widetilde{g}_*\widetilde{\jmath}'_*\mathscr{F}'^* \quad (\widetilde{g}\text{ proper}) \\ &= \widetilde{\imath}^{-1}\boldsymbol{R}\widetilde{g}_*\boldsymbol{R}\widetilde{\jmath}'_*\mathscr{F}'^* \quad (\text{Corollary 8.3}) \\ &= \widetilde{\imath}^{-1}\boldsymbol{R}\widetilde{\jmath}_*\boldsymbol{R}g_*\mathscr{F}'^*.\end{aligned}$$

Using Corollary 8.3 once more, we get (14.1 $*$) by taking the $k$-th cohomology of both terms.

In particular, let us consider the case where, in (2) above, $g$ is an isomorphism, that is, $g : X' \to X$ is a proper modification. We then identify $X \smallsetminus D$ with $X' \smallsetminus D'$, so that $\widetilde{\jmath} = \widetilde{g} \circ \widetilde{\jmath}'$, and $\mathscr{F}^* := g_*\mathscr{F}'^*$ with $\mathscr{F}'^*$. Then (14.1 $*$) reads

$$\boldsymbol{R}\widetilde{g}_*\mathscr{L}' = \mathscr{L}, \tag{14.1 $**$}$$

where $\mathscr{L} = \mathscr{L}^0 = \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$.

***Definition 14.2* (Stokes-filtered local system on $(X, D)$).** — Let $(X, D)$ and $\widetilde{X}$ be as above and $\mathfrak{I}_{\widetilde{X}}$ as in Definition 9.4, and let $\mathscr{F}_\leqslant$ be a pre-$\mathfrak{I}_{\widetilde{X}}$-filtration, i.e., an object of $\mathrm{Mod}(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}},\leqslant})$. We say that $\mathscr{F}_\leqslant$ is a *Stokes-filtered local system on* $(X, D)$ if the following holds:

(1) $\mathscr{F}^* := \widetilde{\jmath}^{-1}\mathscr{F}_\leqslant$ is a local system of finite dimensional $\boldsymbol{k}$-vector spaces on $X^*$,

(2) $\mathscr{F}_\leqslant$ has no subsheaf supported on $\partial\widetilde{X}$, that is, the natural morphism $\widetilde{\imath}^{-1}\mathscr{F}_\leqslant \to \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$ is injective,

(3) this inclusion is a Stokes filtration $(\mathscr{L}, \mathscr{L}_\bullet)$ of $\mathscr{L} := \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$.

We say that $\mathscr{F}_\leqslant$ is *good* if the Stokes-filtered local system $(\mathscr{L}, \mathscr{L}_\bullet)$ is good (see Definition 9.17).

This definition is similar to Definition 4.3, but we do not consider constructible objects $\mathscr{F}^*$, only local systems. We also denote by the same letters $\widetilde{\imath}, \widetilde{\jmath}$ the natural inclusions either in $\widetilde{X}$ or in $\mathfrak{I}^{\text{ét}}$, hoping that this abuse induces no confusion. We will also denote by $(\mathscr{F}, \mathscr{F}_\bullet)$ such a Stokes-filtered local system, in order to emphasize the local system $\mathscr{F} := \widetilde{\imath}^{-1}\widetilde{\jmath}_*\mathscr{F}^*$ on $\widetilde{X}$.

**14.b. Adjunction.** — Let $(\mathscr{F}, \mathscr{F}_\bullet)$ be a Stokes-filtered local system on $(X, D)$ and let $g : (X', D') \to (X, D)$ be a holomorphic map (see notation of §9.e). The diagram of morphisms

$$\begin{tikzcd} \mathfrak{I}_{\widetilde{X}'} & \widetilde{g}^{-1}\mathfrak{I}_{\widetilde{X}} \arrow[l, "g^*"'] \arrow[r, "\widetilde{g}"] & \mathfrak{I}_{\widetilde{X}} \end{tikzcd}$$

allows us to consider the pull-back functor $g^+$ as in Lemma 1.49. It is easy to check that goodness is preserved by pull-back. Below, we will make this property more precise. We will also use the direct image functor $g_+ : D^b(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}}_{\widetilde{X}'},\leqslant}) \to D^b(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}}_{\widetilde{X}},\leqslant})$ of Definition 1.23.

**Proposition 14.3.** — *Assume that* $g : (X', D') \to (X, D)$ *satisfies* 14.a(1) *and* (2) *and is proper, and let* $(\mathscr{F}, \mathscr{F}_\bullet)$ *be a good Stokes-filtered local system on* $(X, D)$*. Then* $g_+ g^+ \mathscr{F}_\leqslant \simeq \boldsymbol{R}\widetilde{g}_* \boldsymbol{k}_{\widetilde{X}'} \otimes \mathscr{F}_\leqslant$ *(where the tensor product is taken in the sense of Remark* 1.18*) in* $D^b(\boldsymbol{k}_{\mathfrak{I}^{\text{ét}}, \leqslant})$*. In particular, each* $\mathscr{H}^k(g_+ g^+ \mathscr{F}_\leqslant)$ *is a good Stokes-filtered local system, isomorphic to* $\boldsymbol{R}^k \widetilde{g}_* \boldsymbol{k}_{\widetilde{X}'} \otimes \mathscr{F}_\leqslant$ *(where the tensor product is taken in the sense of Remark* 1.12*). If moreover* $g$ *is a proper modification, then* $g_+ g^+ \mathscr{F}_\leqslant \simeq \mathscr{F}_\leqslant$*.*

The last assertions are a straightforward consequence of Remark 14.1.

*Proof.* — Let us consider the diagram (1.20) with $q_g = g^*$. The point is to prove that the natural adjunction morphism $\widetilde{g}^{-1}\mathscr{F}_\leqslant \to q_g^{-1}(q_g \widetilde{g} \mathscr{F})_\leqslant$ (notation of Definition 1.34, where we denote by $\widetilde{g}$ both maps $\widetilde{X}' \to \widetilde{X}$ and $\widetilde{X}' \times_{\widetilde{X}} \mathfrak{I}^{\text{ét}} \to \mathfrak{I}^{\text{ét}}$) is an isomorphism. This is clear on $X'^*$, so it is enough to check this on $\partial\widetilde{X}'$, and since the question is local, we can assume that $\mathscr{F}_\leqslant$ is graded. Let $\widetilde{\Sigma} \subset \mathfrak{I}^{\text{ét}}$ be the stratified $\mathfrak{I}$-covering attached to $\mathscr{F}_\leqslant$, let $y' \in \partial\widetilde{X}'$ and set $y = \widetilde{g}(y') \in \partial\widetilde{X}$. The subset $\widetilde{\Sigma}_y \subset \mathfrak{I}^{\text{ét}}_y$ is good, that is, is identified with a good subset $\Phi_{\boldsymbol{d}}$ of $\mathscr{O}_{X_{\boldsymbol{d}},0}(*D)/\mathscr{O}_{X_{\boldsymbol{d}},0}$ after a local ramification of the germ $(X, 0)$ around the germ $(D, 0)$. According to the definition of $(q_g \widetilde{g}^{-1}\mathscr{F})_\leqslant$, we are thus reduced to proving that, for any two germs $\varphi_y, \psi_y \in \widetilde{\Sigma}_y$, we have $g^*(\varphi_y) \leqslant_{y'} g^*(\psi_y)$ only if $\varphi_y \leqslant_y \psi_y$. Since we do not assume that $\widetilde{g}$ is open, we cannot use the general argument of Proposition 9.30, but we can use Lemma 9.34, since $\varphi_y - \psi_y$ is purely monomial when considered in $\mathscr{O}_{X_{\boldsymbol{d}},0}(*D)/\mathscr{O}_{X_{\boldsymbol{d}},0}$, by the goodness assumption.

It is then enough to remark that $\varphi_y \leqslant_y \psi_y$ if and only if, for any $y_{\boldsymbol{d}} \in \partial\widetilde{X}_{\boldsymbol{d}}$ above $y$, we have $\varphi_y \leqslant \psi_y$ at $y_{\boldsymbol{d}}$, when $\varphi, \psi$ are considered as elements of $\Phi_{\boldsymbol{d}}$. ☐

**14.c. Nearby cycles along a function (the good case).** — We denote by $S$ a neighbourhood of the origin in $\mathbb{C}$ and we consider a holomorphic function $f : X \to S$ such that the divisor $D = f^{-1}(0)$ has normal crossings with smooth irreducible components $D_{j\in J}$.

Let $\varpi : \widetilde{X} = \widetilde{X}(D_{j\in J}) \to X$ be the real blow-up of $X$ along the components of $D$ and let $\widetilde{S}$ be the real blow-up of $S$ at the origin. There is a lifting $\widetilde{f} : \widetilde{X} \to \widetilde{S}$ of $f$. We have a diagram of sheaves of ordered abelian groups

$$\begin{tikzcd} \mathfrak{I}_{\widetilde{X}} & \widetilde{f}^{-1}\mathfrak{I}_{\widetilde{S}} \arrow[l, "q_f = f^*"'] \arrow[r, "\widetilde{f}"] & \mathfrak{I}_{\widetilde{S}}. \end{tikzcd}$$

Because $S$ is one-dimensional, the morphism $f^*$ is injective (see Proposition 9.30). Let us also notice that any local section $\varphi$ of $\mathfrak{I}_{\partial\widetilde{S}}$ determines a finite covering $\widetilde{\Sigma}_\varphi \subset \mathfrak{I}^{\text{ét}}_{\partial\widetilde{S}}$ of $\partial\widetilde{S}$. Indeed, this is clear if $\varphi$ is non-ramified, that is, a local section of $\mathfrak{I}_{\partial\widetilde{S},1}$. If it is ramified of order $d$, then one argues by using a ramified covering $\rho_d$ of $(S, 0)$. If $\widetilde{\Sigma}_\varphi$ is such a covering, then its pull-back $\partial\widetilde{X} \times_{\partial\widetilde{S}} \widetilde{\Sigma}_\varphi$ is a finite covering of $\partial\widetilde{X}$, and its image by the inclusion $f^*$ is a finite $\mathfrak{I}_{\partial\widetilde{X}}$-covering of $\partial\widetilde{X}$: this is $\widetilde{\Sigma}_{f^*\varphi}$.

Notice that $D \times \partial\widetilde{S}$ is the boundary of the real blow-up space of $X$ along $f^{-1}(0)$ (see Lemma 8.1) and we have a natural map $(\varpi, \widetilde{f}) : \partial\widetilde{X} \to D \times \partial\widetilde{S}$. For $x$ belonging to a stratum of $D$ of codimension $\ell$ in $X$, and for $\theta \in \partial\widetilde{S} \simeq S^1$, the fibre $(\varpi, \widetilde{f})^{-1}(x, \theta)$ is a union of a finite number of copies of $(S^1)^{\ell-1}$, since $f$ is locally monomial. The natural map $\widetilde{\Sigma}_{f^*\varphi} \to D \times \partial\widetilde{S}$ has a similar property.

***Definition 14.4***. — Let $\widetilde{\Sigma}_\varphi \subset \mathcal{I}^{\text{ét}}_{\partial\widetilde{S}}$ be the covering associated to a local section $\varphi$ of $\mathcal{I}_{\partial\widetilde{S}}$, and let $\widetilde{\Sigma} \subset \mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$ be a good $\mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$-stratified covering of $\partial\widetilde{X}$. We say that $\varphi$ is good with respect to $\widetilde{\Sigma}$ if $\widetilde{\Sigma} \cup \widetilde{\Sigma}_{f^*\varphi}$ (which is a stratified covering of $\partial\widetilde{X}$) is *good*.

Let $(\mathscr{L}, \mathscr{L}_\bullet)$ be a $\mathcal{I}_{\partial\widetilde{X}}$-filtered local system on $\partial\widetilde{X}$. Since $\mathcal{I}_{\partial\widetilde{X}}$ is not Hausdorff over the crossing points of $D$, the graded sheaf $\operatorname{gr}\mathscr{L}$ has to be taken on each stratum of $D$. On the other hand, $\widetilde{f}^{-1}\mathcal{I}_{\partial\widetilde{S}}$ is Hausdorff, since $\mathcal{I}_{\partial\widetilde{S}}$ is so (see Remark 1.1(4)) and, since $q_f^{-1}\mathscr{L}_\leqslant$ defines a pre-$\widetilde{f}^{-1}\mathcal{I}_{\partial\widetilde{S}}$-filtration of $\mathscr{L}$, it is meaningful to consider $\operatorname{gr}\mathscr{L}$ as a subsheaf on $(\widetilde{f}^{-1}\mathcal{I}_{\partial\widetilde{S}})^{\text{ét}} = \partial\widetilde{X} \times_{\partial\widetilde{S}} \mathcal{I}^{\text{ét}}_{\partial\widetilde{S}}$. We will denote this sheaf as $\operatorname{gr}^f\mathscr{L}$ in order to avoid any confusion. For any local section $\varphi$ of $\mathcal{I}_{\partial\widetilde{S}}$, we denote by $\operatorname{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L}$ the restriction of $\operatorname{gr}^f\mathscr{L}$ to $\partial\widetilde{X} \times_{\partial\widetilde{S}} \widetilde{\Sigma}_\varphi$ and we still denote by $\varpi$ the projection $\partial\widetilde{X} \times_{\partial\widetilde{S}} \widetilde{\Sigma}_\varphi \to D$.

***Proposition 14.5***. — *Assume that $(\mathscr{L}, \mathscr{L}_\bullet)$ is a $\mathcal{I}_{\partial\widetilde{X}}$-filtered local system. Then, for any local section $\varphi$ of $\mathcal{I}_{\partial\widetilde{S}}$ we have, over each stratum of $D$, a surjective morphism*

$$\operatorname{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L} \longrightarrow q_f^{-1}\operatorname{gr}_{f^*\varphi}\mathscr{L}.$$

*Assume moreover that $\varphi$ is good with respect to $\widetilde{\Sigma}$. Then the restriction of $\operatorname{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L}$ over each stratum of $D$ is a local system, and $\operatorname{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L}$ is constructible on $\partial\widetilde{X} \times_{\partial\widetilde{S}} \widetilde{\Sigma}_\varphi$ with respect to the pull-back stratification of $D$.*

*Proof*. — By definition, we have $(q_f^{-1}\mathscr{L}_\bullet)_{\leqslant\widetilde{f}^{-1}\varphi} = \mathscr{L}_{\leqslant f^*\varphi}$. Let us fix $y \in \partial\widetilde{X}$ and let us set $\theta = \widetilde{f}(y) \in \partial\widetilde{S}$. The point is to show that $(q_f^{-1}\mathscr{L}_\bullet)_{<\widetilde{f}^{-1}\varphi,y} \subset \mathscr{L}_{<f^*\varphi,y}$. Let $\widetilde{\Sigma} \subset \mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$ be the stratified $\mathcal{I}$-covering attached to $(\mathscr{L}, \mathscr{L}_\bullet)$. On the one hand,

$$(q_f^{-1}\mathscr{L}_\bullet)_{<\widetilde{f}^{-1}\varphi,y} = \sum_{\psi<_\theta\varphi} \mathscr{L}_{\leqslant f^*\psi,y} = \sum_{\psi<_\theta\varphi} \bigoplus_{\substack{\eta\in\widetilde{\Sigma}_y\\ \eta\leqslant_y f^*\psi}} \operatorname{gr}_\eta\mathscr{L}_y = \bigoplus_{\substack{\eta\in\widetilde{\Sigma}_y\\ \exists\,\psi<_\theta\varphi,\,\eta\leqslant_y f^*\psi}} \operatorname{gr}_\eta\mathscr{L}_y. \tag{14.5 $*$}$$

On the other hand,

$$\mathscr{L}_{<f^*\varphi,y} = \bigoplus_{\substack{\eta\in\widetilde{\Sigma}_y\\ \eta<_y f^*\varphi}} \operatorname{gr}_\eta\mathscr{L}_y. \tag{14.5 $**$}$$

Since $f^*$ is compatible with the order, we have $\psi <_\theta \varphi \Rightarrow f^*\psi \leqslant_y f^*\varphi$ and, according to the second part of Proposition 9.30 since $\widetilde{f} : \widetilde{X} \to \widetilde{S}$ is open, we moreover have $f^*\psi \neq f^*\varphi$, that is, $f^*\psi <_y f^*\varphi$. This gives the inclusion $(14.5\,*) \subset (14.5\,**)$.

For the second part of the proposition, we will need a lemma.

***Lemma 14.6.*** — *Let $\eta \in \mathcal{I}_{\partial\widetilde{X},y}$ be such that the associated stratified covering $\widetilde{\Sigma}_\eta \cup \{0\} \subset \mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$ of $\partial\widetilde{X}$ in some neighbourhood of $y$ is good (i.e., after some finite ramification around $D$ near $x = \varpi(y)$, $\rho^*_{\boldsymbol{d}}\eta$ is purely monomial). Assume that $\eta <_y 0$. Then, setting $\theta = \widetilde{f}(y) \in \partial\widetilde{S}$, the property*

$$\exists\, \psi <_\theta 0 \in \mathcal{I}_{\partial\widetilde{S},\theta}, \quad \eta \leqslant_y f^*\psi \tag{14.6$\,*$}$$

*holds if and only if $\eta$ has poles along all the local components of $D$ at $x = \varpi(y)$.*

*Proof.* — Assume first that $\eta$ has poles along all the local components of $D$. We will prove that, if $\psi \in \mathcal{I}_{\partial\widetilde{S},\theta}$ has a pole of order $1/d$ with $d$ big enough (i.e., $\rho_d^{-1}\psi$ has a pole of order one), and if $\psi <_\theta 0$ (such a $\psi$ clearly exists), then $\eta \leqslant_y f^*\psi$. As in Proposition 9.30, one can reduce the statement to the case where $\eta$ is non-ramified, and so $\eta$ is purely monomial. In local coordinates adapted to $D$, we have $D = \{x_1 \cdots x_\ell = 0\}$, $f(x_1, \dots, x_n) = x^{\boldsymbol{k}}$ with $\boldsymbol{k} \in (\mathbb{N}^*)^\ell$, $\eta = u(x_1, \dots, x_n)/x^{\boldsymbol{m}}$ for some $\boldsymbol{m} \in \mathbb{N}^\ell$ and $u$ is a unit, and our assumption means that $\boldsymbol{m} \in (\mathbb{N}^*)^\ell$. It is then enough to choose $d$ such that $\boldsymbol{k} < d\boldsymbol{m}$ with respect to the natural partial ordering of $\mathbb{N}^\ell$. In such a case, $\eta - f^*\psi$ remains purely monomial with the same leading term as $\eta$.

Let us now assume that some $m_j$ is zero, with $j \in \{1, \dots, \ell\}$. Let $\psi$ be such that $\psi <_\theta 0$. Both $\eta$ and $f^*\psi$ are purely monomial, with leading monomial $x^{-\boldsymbol{m}}$ and $x^{-r\boldsymbol{k}}$ respectively, if $\psi$ has leading monomial $t^{-r}$, $r \in \mathbb{Q}^*_+$. By our assumption on $\boldsymbol{m}$, there exists a local modification $\varepsilon : (X', D') \to (X, D)$ near $x \in D$ such that the leading term of $\varepsilon^*(\eta - f^*\psi)$ is $-f^*\psi$. If we had $\eta - f^*\psi \leqslant_y 0$ and $\psi <_\theta 0$, we would also have $-\varepsilon^* f^*\psi \leqslant_{y'} 0$ for any $y' \in \widetilde{\varepsilon}^{-1}(y)$ and $\varepsilon^* f^*\psi <_{y'} 0$, a contradiction. ☐

According to (14.5 $*$) and (14.5 $**$) and to the previous lemma, the kernel of $\mathrm{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L}_y \to q_f^{-1}\mathrm{gr}_{f^*\varphi}\mathscr{L}_y$ is equal to the sum of $\mathrm{gr}_\eta\mathscr{L}_y$, where $\eta \in \widetilde{\Sigma}_y$ is such that $\eta - f^*\varphi$ has no pole along some irreducible component of $D$ going through $x = \varpi(y)$. This condition does not depend on $y$, but only on the stratum of $D$ which $x$ belongs to. This shows that this kernel is a local system over each stratum of $D$. On the other hand, $\mathrm{gr}_{f^*\varphi}\mathscr{L}$ is also a local system over each stratum of $D$. Therefore, so is $\mathrm{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L}$. ☐

***Definition 14.7* (Nearby cycles, the good case).** — Let $(\mathscr{F}, \mathscr{F}_\bullet)$ be a *good* Stokes-filtered local system on $(X, D)$ (see Definition 14.2) with associated stratified $\mathcal{I}$-covering $\widetilde{\Sigma} \subset \mathcal{I}^{\text{ét}}_{\partial\widetilde{X}}$, and let $\varphi$ be a local section of $\mathcal{I}_{\partial\widetilde{S}}$ defining a finite covering $\widetilde{\Sigma}_\varphi \subset \mathcal{I}^{\text{ét}}_{\partial\widetilde{S}}$ of $\partial\widetilde{S} \simeq S^1$. Let us assume that $\varphi$ is *good* with respect to $\widetilde{\Sigma}$. We then set

$$\begin{aligned}\widetilde{\psi}^\varphi_f(\mathscr{F}, \mathscr{F}_\bullet) &= \mathrm{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L},\\ \psi^\varphi_f(\mathscr{F}, \mathscr{F}_\bullet) &= \boldsymbol{R}(\varpi, \widetilde{f})_*\, \mathrm{gr}^f_{\widetilde{f}^{-1}\varphi}\mathscr{L}.\end{aligned}$$

We have seen that $\widetilde{\psi}_f^\varphi(\mathscr{F},\mathscr{F}_\bullet)$ is locally constant with respect to the pull-back stratification of $D$. Since the map $(\varpi,\widetilde{f})$ is a topological fibration when restricted above each stratum of $D$ with fibre homeomorphic to a finite number of copies of $(S^1)^{\ell-1}$ when the stratum has codimension $\ell$, it follows that the cohomology sheaves of $\psi_f^\varphi(\mathscr{F},\mathscr{F}_\bullet)$ are locally constant on $D_I\times\partial\widetilde{S}$ for each stratum $D_I$ of $D$. According to Lemma 9.37, they can be regarded as $\mathbb{C}$-constructible sheaves on $D$ (constructible with respect to the natural stratification) equipped with an automorphism (the monodromy around $f=0$). We will denote by $(\psi_f^\varphi(\mathscr{F},\mathscr{F}_\bullet),T)$ the corresponding object of $D^b_c(D)$ equipped with its automorphism $T$ (we implicitly extend the equivalence of Lemma 9.37 to the derived category).

More precisely, let us denote by $D^b_{D\text{-c}}(D\times\partial\widetilde{S})$ the full subcategory of $D^b(D\times\partial\widetilde{S})$ whose objects are constructible with respect to the natural stratification $(D_I\times\partial\widetilde{S})_I$. For each $\lambda\in\mathbb{C}^*$, we denote by $L_{\lambda^{-1},\infty}$ the local system on $\partial\widetilde{S}$ whose fibre is the polynomial ring $\mathbb{C}[x]$ and the monodromy is the automorphism $\lambda^{-1}\cdot\mathrm{U}_\infty$, where $\mathrm{U}_\infty=\exp(2\pi i\mathrm{N}_\infty)$ and $\mathrm{N}_\infty:\mathbb{C}[x]\to\mathbb{C}[x]$ is defined by $\mathrm{N}_\infty(x^k)=x^{k-1}$ if $k\geqslant 1$ and $\mathrm{N}_\infty(1)=0$. We denote similarly the pull-back of $L_{\lambda^{-1},\infty}$ to $D\times\partial\widetilde{S}$, and by $p:D\times\partial\widetilde{S}\to D$ the projection. We define the functor $\psi_\lambda$ from $D^b_{D\text{-c}}(D\times\partial\widetilde{S})$ to $D^b(D)$ by

$$\psi_\lambda(\mathscr{G})=\boldsymbol{R}p_*(L_{\lambda^{-1},\infty}\otimes_{\mathbb{C}}\mathscr{G}).$$

This functor takes values in the derived category of bounded constructible complexes on $D$ (constructible with respect to the natural stratification). Moreover, $\psi_\lambda(\mathscr{G})$ is equipped functorially with an automorphism $T=\lambda\cdot(\boldsymbol{R}p_*(\mathrm{U}_\infty\otimes\mathrm{Id}))$.

***Lemma 14.8.*** — *The functor $\bigoplus_{\lambda\in\mathbb{C}^*}\psi_\lambda$ induces an equivalence between $D^b_{D\text{-}c}(D\times\partial\widetilde{S})$ and the category $(D^b_c(D),T)$ whose objects are pairs of an object of $D^b_c(D)$ (constructibility with respect to the natural stratification is understood) and an automorphism $T$ of this object.* ☐

This lemma is a natural extension to derived categories of Lemma 9.37, if one uses the finite determination functor of [**13**, Lemm 1.5]. In particular, for a given object $\mathscr{G}$ of $D^b_{D\text{-c}}(D\times\partial\widetilde{S})$, all $\psi_\lambda\mathscr{G}$ but a finite number are isomorphic to zero locally on $D$. In the following, we will have to consider $(\psi^\varphi_{f,\lambda}(\mathscr{F},\mathscr{F}_\bullet),T)$ for each $\lambda\in\mathbb{C}^*$.

The following proposition will be essential to define nearby cycles when the goodness condition on $\varphi$ is not fulfilled.

***Proposition 14.9.*** — *Let us keep the previous assumptions and let $\varepsilon:(X',D')\to(X,D)$ be a proper modification, where $D'=\varepsilon^{-1}(D)$ is a divisor with normal crossings and $\varepsilon:X'\smallsetminus D'\to X\smallsetminus D$ is an isomorphism. Let us set $f'=f\circ\varepsilon:(X',D')\to(S,0)$. We have*

$$\widetilde{\psi}_f^\varphi(\mathscr{F},\mathscr{F}_\bullet)=\boldsymbol{R}\widetilde{\varepsilon}_*\widetilde{\psi}^\varphi_{f'}\varepsilon^+(\mathscr{F},\mathscr{F}_\bullet),\tag{14.9 $*$}$$

$$\psi_f^\varphi(\mathscr{F},\mathscr{F}_\bullet)=\boldsymbol{R}(\varepsilon,\mathrm{Id}_{\partial\widetilde{S}})_*\psi^\varphi_{f'}\varepsilon^+(\mathscr{F},\mathscr{F}_\bullet).\tag{14.9 $**$}$$

*A preliminary result.* — Let us denote by $D_{j\in J}$ (resp. $D'_{j'\in J'}$) the irreducible components of $D$ (resp. $D'$). By assumption, they are smooth, as well as their various intersections. Since $X$ and $X'$ are smooth, the modification $\varepsilon$ is an isomorphism in codimension two on $X$. For any $j\in J$, there exists thus a unique $j'\in J'$ such that $\varepsilon: D'_{j'}\to D_j$ is dominant, and it is then a proper modification of $D_j$. We will denote $j$ instead of $j'$ and we will regard $J$ as a subset of $J'$. We will call $D'_j$ the *strict transform* of $D_j$. Recall that, for a subset $I\subset J$, $D(I)$ denotes $\bigcup_{j\in I} D_j$ and similarly $D'(I)=\bigcup_{j\in I} D'_j$, and $D_I=\bigcap_{j\in I} D_j$, $D'_I=\bigcap_{j\in I} D'_j$. Recall also that we denote by $\varpi:\widetilde X\to X$ (resp. $\varpi':\widetilde X'\to X'$) the real blowing-up of the components of $D$ (resp. $D'$) and by $\widetilde\varepsilon:\widetilde X'\to\widetilde X$ the lifting of $\varepsilon$.

**Lemma 14.10.** — *Let us fix a subset $I$ of $J$ and let $\mathscr{F}'_I$ denote the constant sheaf on $\partial\widetilde X'\smallsetminus\varpi'^{-1}(D'(I))$ extended by $0$ on $\varpi'^{-1}(D'(I))$. Then $\boldsymbol{R}\widetilde\varepsilon_*\mathscr{F}'_I$ is zero on $\varpi^{-1}(D(I))$.*

*Proof.* — It is equivalent to proving that $\boldsymbol{R}\varepsilon_*\mathbb{C}_{\widetilde X'_{|D'(I)}}=\mathbb{C}_{\widetilde X_{|D(I)}}$ since, according to (14.1 $**$), we have $\boldsymbol{R}\widetilde\varepsilon_*\mathbb{C}_{\partial\widetilde X'}=\mathbb{C}_{\partial\widetilde X}$. Assume first that $I=\{j_o\}$. By definition (see Lemma 8.2), we have a cartesian square

$$\begin{tikzcd}
\widetilde X_{|D_{j_o}}=\widetilde X(D_{j\in J})_{|D_{j_o}} \arrow[r] \arrow[d] & \widetilde X(D_{j_o})_{|D_{j_o}}=\partial\widetilde X(D_{j_o}) \arrow[d] \\
\widetilde X(D_{j\in J\smallsetminus\{j_o\}})_{|D_{j_o}} \arrow[r] & D_{j_o}
\end{tikzcd}$$

and $\widetilde X(D_{j\in J\smallsetminus\{j_o\}})_{|D_{j_o}}$ is nothing but the real blow-up space of $D_{j_o}$ along the components $D_{j_o}\cap D_{j\in J\smallsetminus\{j_o\}}$, that we will denote by $\widetilde D_{j_o}$ for short. We denote similarly by $\widetilde D'_{j_o}$ the real blow-up space of $D'_{j_o}$ along the components $D'_{j_o}\cap D'_{j'\in J'\smallsetminus\{j_o\}}$, that we identify to $\widetilde X'(D'_{j'\in J'\smallsetminus\{j_o\}})_{|D'_{j_o}}$. Then the map $\widetilde\varepsilon:\widetilde X'_{|D'_{j_o}}=\widetilde X'(D'_{j'\in J'})_{|D'_{j_o}}\to\widetilde X_{|D_{j_o}}$ factorizes through

$$\widetilde{\varepsilon_{|D'_{j_o}}}\times\mathrm{Id}:\widetilde D'_{j_o}\times_{D_{j_o}}\partial\widetilde X(D_{j_o})\longrightarrow\widetilde D_{j_o}\times_{D_{j_o}}\partial\widetilde X(D_{j_o}).$$

We will show that the map $\widetilde X'_{|D'_{j_o}}\to\widetilde D'_{j_o}\times_{D_{j_o}}\partial\widetilde X(D_{j_o})$ is an isomorphism. Assume that this is proved. Then we have an equality of maps

$$\begin{tikzcd}
\widetilde X'_{|D'_{j_o}} \arrow[r, equal] \arrow[d, "\widetilde\varepsilon"'] & \widetilde D'_{j_o}\times_{D_{j_o}}\partial\widetilde X(D_{j_o}) \arrow[d, "\widetilde{\varepsilon_{|D'_{j_o}}}\times\mathrm{Id}"] \\
\widetilde X_{|D_{j_o}} \arrow[r, equal] & \widetilde D_{j_o}\times_{D_{j_o}}\partial\widetilde X(D_{j_o})
\end{tikzcd}\tag{14.11}$$

We know, by (14.1 $**$), that $\boldsymbol{R}\widetilde\varepsilon_*\mathbb{C}_{\widetilde D'_{j_o}}=\mathbb{C}_{\widetilde D_{j_o}}$. It follows that $\boldsymbol{R}\widetilde\varepsilon_*\mathbb{C}_{\widetilde X'_{D'_{j_o}}}=\mathbb{C}_{\widetilde X_{D_{j_o}}}$.

To prove the assertion is a local question on $D'_{j_o}$. We can use the setting already introduced in §9.e, and replace $f$ there with $\varepsilon$ here. We will assume that $D_{j_o}$ is locally defined by $x_1=0$ and $D'_{j_o}$ by $x'_1=0$. The set $L$ parametrizes the components of $D$ going through the chosen origin in $X$ and $L'$ those components passing through the chosen origin 0 in $X'$. We check the assertion on $\varpi'^{-1}(0)$. Then $\varepsilon$ also has the local form (9.28). Moreover, in the neighbourhood of a generic point of $D'_{j_o}$, $\varepsilon$ is an isomorphism, which implies that the component $k_{1,1}$ in (9.28) is equal to one. The formula for $\theta_1$ obtained as the first line in (9.29) shows then that $\theta'_1$ can be uniquely recovered from $\theta_1$ and $\theta'_2,\dots,\theta'_{\ell'},x'_{\ell'+1},\dots,x'_n$, which is the desired assertion.

By induction on $\#I$ we obtain that $\boldsymbol{R}\widetilde{\varepsilon}_*\mathbb{C}_{\widetilde{X}'_{|D'_I}}=\mathbb{C}_{\widetilde{X}_{|D_I}}$: indeed, if $I=\{j_o\}\cup I_1$, we can apply the inductive assumption to $\widetilde{D}'_{j_o|I_1}\to\widetilde{D}_{j_o|I_1}$; we notice that, for $k\in I_1$, $D'_{j_o}\cap D'_k$ is the strict transform of the divisor $D_{j_o}\cap D_k$ of $D_{j_o}$ by the proper modification $D'_{j_o}\to D_{j_o}$ induced by $\varepsilon$; therefore, $D'_{j_o|I_1}=D'_I$; it suffices then to use (14.11) restricted to $D'_I$ and $D_I$.

The Mayer-Vietoris sequence gives a quasi-isomorphism

$$\mathbb{C}_{\widetilde{X}'_{|D'(I)}}\xrightarrow{\sim}\Big\{\bigoplus_{\substack{K\subset I\\ \#K=1}}\mathbb{C}_{\widetilde{X}'_{|D'_K}}\longrightarrow\bigoplus_{\substack{K\subset I\\ \#K=2}}\mathbb{C}_{\widetilde{X}'_{|D'_K}}\longrightarrow\cdots\Big\}.$$

According to the previous result (applied to $K$ instead of $I$) for each $\mathbb{C}_{\widetilde{X}'_{|D'_K}}$ with $K\subset I$, we obtain that $\boldsymbol{R}\varepsilon_*\mathbb{C}_{\widetilde{X}'_{|D'(I)}}$ is quasi-isomorphic to the Mayer-Vietoris complex

$$\Big\{\bigoplus_{\substack{K\subset I\\ \#K=1}}\mathbb{C}_{\widetilde{X}_{|D_K}}\longrightarrow\bigoplus_{\substack{K\subset I\\ \#K=2}}\mathbb{C}_{\widetilde{X}_{|D_K}}\longrightarrow\cdots\Big\},$$

that is, to $\mathbb{C}_{\widetilde{X}_{|D(I)}}$. $\square$

*Proof of Proposition* 14.9. — We will consider the following diagram:

$$\begin{tikzcd}
& \widetilde{f}'^{-1}\mathfrak{I}_{\widetilde{S}} \arrow[rr, "\widetilde{\varepsilon}"] \arrow[dl, "q_{f'}"'] \arrow[dr, "\widetilde{\varepsilon}^{-1}q_f"] & & \widetilde{f}^{-1}\mathfrak{I}_{\widetilde{S}} \arrow[r, "\widetilde{f}"] \arrow[d, "q_f"] & \mathfrak{I}_{\widetilde{S}} \arrow[dd] \\
\mathfrak{I}_{\widetilde{X}'} \arrow[dr] & & \widetilde{\varepsilon}^{-1}\mathfrak{I}_{\widetilde{X}} \arrow[ll, "q_\varepsilon"'] \arrow[r] \arrow[dl] & \mathfrak{I}_{\widetilde{X}} \arrow[d] & \\
& \widetilde{X}' \arrow[rr, "\widetilde{\varepsilon}"] & & \widetilde{X} \arrow[r, "\widetilde{f}"] & \widetilde{S}
\end{tikzcd}$$

where we recall that, for a map $g$, $q_g$ is a notation for $g^*$.

On the one hand, we have by definition $(q_{f'}^{-1}\varepsilon^+\mathscr{L})_{\leqslant\widetilde{f}'^{-1}\varphi}=(\varepsilon^+\mathscr{L})_{\leqslant f'^*\varphi}$. Recall also (see Definition 1.34) that $(\varepsilon^+\mathscr{L})_{\leqslant}=\mathrm{Tr}_{\leqslant q_\varepsilon}(\widetilde{\varepsilon}^{-1}\mathscr{L}_{\leqslant})$. Therefore,

$$(q_{f'}^{-1}\varepsilon^+\mathscr{L})_{\leqslant\widetilde{f}'^{-1}\varphi}=\sum_{\eta,\,\varepsilon^*\eta\leqslant f'^*\varphi}\widetilde{\varepsilon}^{-1}\mathscr{L}_{\leqslant\eta}.$$

On the other hand, $\widetilde{\varepsilon}^{-1}(q_f^{-1}\mathscr{L})_{\leqslant\widetilde{f}^{-1}\varphi}=\sum_{\eta\leqslant f^*\varphi}\widetilde{\varepsilon}^{-1}\mathscr{L}_{\leqslant\eta}$. We thus have a natural inclusion $\widetilde{\varepsilon}^{-1}(q_f^{-1}\mathscr{L})_{\leqslant\widetilde{f}^{-1}\varphi}\subset(q_{f'}^{-1}\varepsilon^+\mathscr{L})_{\leqslant\widetilde{f}'^{-1}\varphi}$ (as subsheaves of the pull-back of the local system $\mathscr{L}$). We will show that this is an equality by checking this property at each $y' \in \partial\widetilde{X}'$. Let us set $y = \widetilde{\varepsilon}(y')$ and $\theta = \widetilde{f}(y)$. A computation similar to (14.5 $*$) gives

$$\begin{aligned}(q_{f'}^{-1}\varepsilon^+\mathscr{L})_{\leqslant \widetilde{f}'^{-1}\varphi, y'} &= \bigoplus_{\substack{\eta\in\widetilde{\Sigma}_y\\ \varepsilon^*\eta\leqslant_{y'} f'^*\varphi}} \operatorname{gr}_\eta \mathscr{L}_y,\\ \big(\widetilde{\varepsilon}^{-1}(q_f^{-1}\mathscr{L})_{\leqslant\widetilde{f}^{-1}\varphi}\big)_{y'} = (q_f^{-1}\mathscr{L})_{\leqslant\widetilde{f}^{-1}\varphi, y} &= \bigoplus_{\substack{\eta\in\widetilde{\Sigma}_y\\ \eta\leqslant_y f^*\varphi}} \operatorname{gr}_\eta \mathscr{L}_y,\end{aligned}\tag{14.12}$$

so the desired equality is a consequence of Lemma 9.34 applied to $\eta - f^*\varphi$ (due to the assumption of goodness, $\eta - f^*\varphi$ is purely monomial) and the map $\varepsilon$.

We also have an inclusion $\widetilde{\varepsilon}^{-1}(q_f^{-1}\mathscr{L})_{<\widetilde{f}^{-1}\varphi} \subset (q_{f'}^{-1}\varepsilon^+\mathscr{L})_{<\widetilde{f}'^{-1}\varphi}$, by using the same argument as for $\leqslant$, but we do not claim that it is an equality (the argument for $\leqslant$ used the goodness property of $\varphi$ with respect to $\widetilde{\Sigma}$, a property that one cannot use for $\psi <_\theta \varphi$). In any case, we conclude that there is a surjective morphism

$$\lambda : \widetilde{\varepsilon}^{-1}\widetilde{\psi}_f^\varphi(\mathscr{F}, \mathscr{F}_\bullet) \longrightarrow \widetilde{\psi}_{f'}^\varphi \varepsilon^+(\mathscr{F}, \mathscr{F}_\bullet)$$

and we will compute the kernel in the neighbourhood of $\widetilde{\varepsilon}^{-1}(y)$ for each $y \in \partial\widetilde{X}$. According to Lemma 14.6 and the computation (14.12), the kernel at $y' \in \widetilde{\varepsilon}^{-1}(y)$ is identified with the sum of $\operatorname{gr}_\eta \mathscr{L}_y$ for those $\eta \in \widetilde{\Sigma}_y$ such that $\varepsilon^*(\eta - f^*\varphi)$ has poles along all the components of $D'$ going through $x' = \varpi'(y')$, but $\eta - f^*\varphi$ has no poles along some components of $D$ going through $x = \varpi(y)$, whose union is denoted by $D(I)$. Since $\eta - f^*\varphi$ is purely monomial, $\varepsilon^*(\eta - f^*\varphi)$ has no poles exactly along the strict transform $D'(I)$ of $D(I)$. Therefore, the $\eta$-component of the kernel of $\lambda_{|\widetilde{\varepsilon}^{-1}(y)}$ is equal to the pull-back of $\operatorname{gr}_\eta \mathscr{L}_y$ on $\widetilde{\varepsilon}^{-1}(y) \smallsetminus D'(I)$ and is zero on $\widetilde{\varepsilon}^{-1}(y) \cap D'(I)$. This holds in some small neighbourhood of $y$. Applying Lemma 14.10 together with the projection formula for the proper morphism $\widetilde{\varepsilon}$ implies that $\boldsymbol{R}\varepsilon_*$ of this $\eta$-component is zero. Since this holds for any $\eta \in \widetilde{\Sigma}_y$, we conclude that we have an isomorphism in the neighbourhood of $y$:

$$\boldsymbol{R}\varepsilon_*\lambda : \boldsymbol{R}\varepsilon_*\widetilde{\varepsilon}^{-1}\widetilde{\psi}_f^\varphi(\mathscr{F}, \mathscr{F}_\bullet) \longrightarrow \boldsymbol{R}\varepsilon_*\widetilde{\psi}_{f'}^\varphi\varepsilon^+(\mathscr{F}, \mathscr{F}_\bullet),$$

and since $y$ was arbitrary, this is an isomorphism all over $\partial\widetilde{X}$. On the other hand, applying once more the projection formula and (14.1 $**$) we have $\boldsymbol{R}\varepsilon_*\widetilde{\varepsilon}^{-1}\widetilde{\psi}_f^\varphi(\mathscr{F}, \mathscr{F}_\bullet) = \widetilde{\psi}_f^\varphi(\mathscr{F}, \mathscr{F}_\bullet)$. This ends the proof of (14.9 $*$).

By using the commutative diagram

$$\begin{tikzcd} \partial\widetilde{X}' \arrow[r, "\widetilde{\varepsilon}"] \arrow[d, "{(\varpi', \widetilde{f}')}"'] & \partial\widetilde{X} \arrow[d, "{(\varpi, \widetilde{f})}"] \\ D' \times \partial\widetilde{S} \arrow[r, "\varepsilon \times \mathrm{Id}"'] & D \times \partial\widetilde{S} \end{tikzcd}$$

one obtains (14.9 $**$). $\square$

**14.d. Nearby cycles along a function (dimension two).** — We keep the setting of §14.c. Our aim is to define the nearby cycles functors $\widetilde{\psi}_f^\varphi$ and $\psi_f^\varphi$ for *good* Stokes-filtered local systems $(\mathscr{F},\mathscr{F}_\bullet)$ on $(X,D)$, *without assuming that* $\varphi$ *is good* with respect to the stratified $\mathcal{I}$-covering $\widetilde{\Sigma}$ associated to $(\mathscr{F},\mathscr{F}_\bullet)$. One should not use Definition 14.7 in general, since Proposition 14.9 does not hold in general without the goodness property of $\varphi$, and we expect that the property proved in this proposition to be satisfied by nearby cycles. The idea is then to define nearby cycles $\widetilde{\psi}_f^\varphi$ by choosing a modification $\varepsilon:(X',D')\to(X,D)$ so that $\varphi$ becomes good with respect to $\varepsilon^+(\mathscr{F},\mathscr{F}_\bullet)$ and take the formulas of Proposition 14.9 as a definition. This is similar to the notion of *good cell* introduced in [**62**]. The proposition itself is useful to prove that this definition does not depend on the choice of $\varepsilon$, provided that the goodness property is fulfilled. We will now restrict to the case where $\dim X=2$ in order to apply this program.

***Proposition 14.13.*** — *Let us fix* $x\in D$ *and let* $\eta$ *be a local section of* $\mathcal{I}$ *in some neighbourhood of* $y\in\varpi^{-1}(x)$*. Then there exists a finite sequence of point blowing-ups* $\varepsilon:(X',D')\to(X,D)$ *with centers projecting to* $x$ *such that* $\varepsilon^*\eta$ *(see Proposition* 9.30*) is good on* $\varepsilon^{-1}(U)$*, where* $U$ *is some open neighbourhood of* $x$ *in* $X$*.*

*Proof.* — If $\eta$ is not ramified, that is, $\eta\in\mathscr{O}_{X,x}(*D)/\mathscr{O}_{X,x}$, goodness means pure monomiality, and the assertion is that of Lemma 9.11, according to the well-known property that any proper modification of a complex surface is dominated by a sequence of point blowing-ups.

In general, by definition of $\mathcal{I}$, there exists a local ramification $\rho_{\boldsymbol{d}}:U_{\boldsymbol{d}}\to U$ such that $\rho_{\boldsymbol{d}}^*\eta$ is not ramified, i.e., belongs to $\mathscr{O}_{U_{\boldsymbol{d}},0}(*D)/\mathscr{O}_{U_{\boldsymbol{d}},0}$. For any automorphism $\sigma$ of $\rho_{\boldsymbol{d}}$, $\sigma^*\rho_{\boldsymbol{d}}^*\eta$ is also not ramified and has the same polar locus (with the same multiplicities) as $\rho_{\boldsymbol{d}}^*\eta$. The product of all $\sigma^*\rho_{\boldsymbol{d}}^*\eta$ when $\sigma$ varies in the Galois group of $\rho_{\boldsymbol{d}}$ can be written $\rho_{\boldsymbol{d}}^*\xi$, where $\xi$ belongs to $\mathscr{O}_{U,0}(*D)/\mathscr{O}_{U,0}$.

Note now that the product of elements of $\mathscr{O}_{U_{\boldsymbol{d}},0}(*D)/\mathscr{O}_{U_{\boldsymbol{d}},0}$ having the same polar divisor is purely monomial if and only if each term is purely monomial.

We can apply the first part of the proof to $\xi$ and get a sequence $\varepsilon$ of point blowing-ups such that $\varepsilon^*\xi$ is purely monomial. Let $x'\in\varepsilon^{-1}(x)$ and let $D'=\varepsilon^{-1}(D)$. One can choose a local ramification $\rho_{\boldsymbol{d}'}$ in the neighbourhood of $x'$ which dominates $U_{\boldsymbol{d}}$ by a generically finite map $\varepsilon'$. We can apply the first property above to $\rho_{\boldsymbol{d}'}^*\varepsilon^*\xi=\prod_\sigma\varepsilon'^*\sigma^*\rho_{\boldsymbol{d}}^*\eta$ to conclude that $\varepsilon^*\eta$ is purely monomial after the local ramification $\rho_{\boldsymbol{d}'}$. $\square$

***Corollary 14.14.*** — *Let* $\widetilde{\Sigma}\subset\mathcal{I}^{\text{ét}}$ *be a good stratified* $\mathcal{I}$*-covering. Then, for any local section* $\varphi$ *of* $\mathcal{I}_{\partial\widetilde{S}}$*, there exists, over any compact set* $K$ *of* $D$*, a finite sequence of point blowing-ups* $\varepsilon:(X',D')\to(X,D)$ *such that* $\varphi$ *is good with respect to* $\varepsilon^*\widetilde{\Sigma}$ *in some neighbourhood of* $\varepsilon^{-1}(K)$*.*

***Definition 14.15.*** — Let $f:(X,D)\to(S,0)$ be a proper holomorphic map to a disc $S$, where $D=f^{-1}(0)$ is a divisor with normal crossings and smooth components. Let $(\mathscr{F},\mathscr{F}_\bullet)$ be a good Stokes-filtered local system on $(X,D)$ with associated stratified $\mathcal{I}$-covering $\widetilde{\Sigma}$. For any local section $\varphi$ of $\mathcal{I}_{\partial\widetilde{S}}$, we *define*

$$\widetilde{\psi}_f^{\varphi}(\mathscr{F},\mathscr{F}_\bullet):=\boldsymbol{R}\widetilde{\varepsilon}_*\widetilde{\psi}_{f'}^{\varphi}\varepsilon^+(\mathscr{F},\mathscr{F}_\bullet), \tag{14.15 $*$}$$

where $f'=f\circ\varepsilon$ and $\varepsilon:(X',D')\to(X,D)$ is any finite sequence of point blowing-ups such that $\varphi$ is good with respect to $\varepsilon^*\widetilde{\Sigma}$, and we set, as above,

$$\psi_f^{\varphi}(\mathscr{F},\mathscr{F}_\bullet)=\boldsymbol{R}(\varpi,\widetilde{f})_*\widetilde{\psi}_f^{\varphi}(\mathscr{F},\mathscr{F}_\bullet)$$

***Remark 14.16.*** — That the choice of $\varepsilon$ is irrelevant follows from Proposition 14.9. We could also avoid the properness assumption on $f$, by working on an exhaustive sequence of compact subsets of $D$. Lastly, notice the formula

$$\psi_f^{\varphi}(\mathscr{F},\mathscr{F}_\bullet)=\boldsymbol{R}(\varepsilon,\mathrm{Id}_{\partial\widetilde{S}})_*\psi_{f'}^{\varphi}\varepsilon^+(\mathscr{F},\mathscr{F}_\bullet). \tag{14.16 $**$}$$

### 14.e. Moderate nearby cycles of holonomic $\mathscr{D}$-modules

*Restriction and de Rham complex of holonomic $\mathscr{D}$-modules.* — We refer to the book [**34**] and to the introductory article [**57**] for the basic results we need concerning holonomic $\mathscr{D}$-modules. Let $X$ be a complex manifold and let $Z$ be a closed analytic subset. We denote by $i_Z:Z\hookrightarrow X$ the closed inclusion and by $j_Z:X\smallsetminus Z\hookrightarrow X$ the complementary open inclusion. We will denote by $i_{Z,+}i_Z^+$ the functor of local algebraic cohomology, denoted by $\boldsymbol{R}\Gamma_{[Z]}$ in [**34**] and $\boldsymbol{R}\mathrm{alg}\Gamma_Z$ in [**57**], and by $j_{Z,+}j_Z^+$ the localization functor denoted by $\boldsymbol{R}\Gamma_{[X\smallsetminus Z]}$ in [**34**] and $\boldsymbol{R}\bullet(*Z)$ in loc. cit. There is a distinguished triangle in $D^b(\mathscr{D}_X)$ (see [**34**, Th. 3.29(2)]):

$$i_{Z,+}i_Z^+\mathscr{M}\longrightarrow\mathscr{M}\longrightarrow j_{Z,+}j_Z^+\mathscr{M}\xrightarrow{+1}$$

By a theorem of Kashiwara, these functors preserve the bounded derived category $D^b_{\mathrm{hol}}(\mathscr{D}_X)$ of complexes of $\mathscr{D}_X$-modules with holonomic cohomology. We denote by $\boldsymbol{D}$ the duality (contravariant) functor, from $D^b_{\mathrm{coh}}(\mathscr{D}_X)$ to itself. It also preserves $D^b_{\mathrm{hol}}(\mathscr{D}_X)$. In particular, $\boldsymbol{D}\mathscr{M}$ is a holonomic $\mathscr{D}_X$-module if $\mathscr{M}$ is so. We also have a functorial isomorphism $\boldsymbol{D}\circ\boldsymbol{D}\simeq\mathrm{Id}$. We then set

$$i_{Z,+}i_Z^{\dagger}=\boldsymbol{D}(i_{Z,+}i_Z^+)\boldsymbol{D},\qquad j_{Z,\dagger}j_Z^+=\boldsymbol{D}(j_{Z,+}j_Z^+)\boldsymbol{D}.$$

If $\mathscr{O}_{\widehat{Z}}$ denotes the formalization of $\mathscr{O}_X$ along $Z$, we have

***Proposition 14.17* (see [57, Cor. 2.7-2]).** — *There is a canonical functorial isomorphism in $D^b(\mathscr{D}_X)$:*

$$\boldsymbol{R}\mathscr{H}\!om_{\mathscr{D}_X}(i_{Z,+}i_Z^+\bullet,\mathscr{O}_X)\simeq\boldsymbol{R}\mathscr{H}\!om_{\mathscr{D}_X}(\bullet,\mathscr{O}_{\widehat{Z}}).$$

***Corollary 14.18.*** — *There is a canonical functorial isomorphism in $D^b_{\mathrm{hol}}(\mathscr{D}_X)$:*

$$\mathrm{DR}(i_{Z,+}i_Z^{\dagger}\mathscr{M})\simeq\mathrm{DR}(\mathscr{O}_{\widehat{Z}}\otimes_{\mathscr{O}_X}\mathscr{M}).$$

*Proof.* — Proposition 14.17 applied to $\boldsymbol{D}\mathscr{M}$ gives an isomorphism

$$\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_X}(\boldsymbol{D}(i_{Z,+}i_Z^\dagger\mathscr{M}),\mathscr{O}_X)\simeq\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_X}(\boldsymbol{D}\mathscr{M},\mathscr{O}_{\widehat{Z}}).$$

On the one hand, since $i_{Z,+}i_Z^\dagger\mathscr{M}$ has coherent cohomology (being holonomic), we have a canonical isomorphism of the left-hand term with $\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_X}(\mathscr{O}_X,i_{Z,+}i_Z^\dagger\mathscr{M})$, according to [**34**, (3.14)] and the canonical isomorphism $\boldsymbol{D}\mathscr{O}_X\simeq\mathscr{O}_X$.

On the other hand, if we set $\mathscr{D}_{\widehat{Z}}=\mathscr{O}_{\widehat{Z}}\otimes_{\mathscr{O}_X}\mathscr{D}_X$ and $\mathscr{M}_{\widehat{Z}}=\mathscr{O}_{\widehat{Z}}\otimes_{\mathscr{O}_X}\mathscr{M}$, we have, since $\mathscr{O}_{\widehat{Z}}$ is $\mathscr{O}_X$-flat,

$$\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_X}(\boldsymbol{D}\mathscr{M},\mathscr{O}_{\widehat{Z}})\simeq\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_{\widehat{Z}}}(\boldsymbol{D}\mathscr{M}_{\widehat{Z}},\mathscr{O}_{\widehat{Z}})$$

and applying the previous argument in $D^b_{\mathrm{coh}}(\mathscr{D}_{\widehat{Z}})$, the latter term is isomorphic to $\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_{\widehat{Z}}}(\mathscr{O}_{\widehat{Z}},\mathscr{M}_{\widehat{Z}})$. We conclude by using that $\boldsymbol{R}\mathscr{H}om_{\mathscr{D}_X}(\mathscr{O}_X,\bullet)\simeq\mathrm{DR}(\bullet)[\dim X]$, and a similar property for $\mathscr{D}_{\widehat{Z}}$-modules. ☐

*Moderate nearby cycles for holonomic $\mathscr{D}$-modules.* — Let $f:X\to S=\mathbb{C}$ be a holomorphic function and let $\mathscr{M}$ be a holonomic $\mathscr{D}_X$-module. We will recall (see e.g. [**59, 58**]) the construction of the $\mathscr{D}_X$-module $\psi_f^{\mathrm{mod}}\mathscr{M}$ supported on $D:=f^{-1}(0)$.

Let $t$ be the coordinate on $S=\mathbb{C}$. For any complex number $\alpha$ and any integer $k\geqslant 0$, denote by $\mathscr{N}_{\alpha,k}$ the rank $k+1$ free $\mathscr{O}_S(*0)$-module $\bigoplus_{j=0}^k\mathscr{O}_S(*0)e_{\alpha,j}$, equipped with the $\mathscr{D}_S$-action defined by $t\partial_t(e_{\alpha,j})=(\alpha+1)e_{\alpha,j}+e_{\alpha,j-1}$, with the convention that $e_{\alpha,j}=0$ for $j<0$. Then $\mathscr{N}_{\alpha,k}$ is equipped with a $\mathscr{D}_S$-linear nilpotent endomorphism $\mathrm{N}$, defined by $\mathrm{N}e_{\alpha,j}=e_{\alpha,j-1}$, and a natural $\mathscr{D}_S$-linear inclusion $\mathscr{N}_{\alpha,k}\hookrightarrow\mathscr{N}_{\alpha,k+1}$. There is a canonical isomorphism $\mathscr{N}_{\alpha+1,k}\simeq\mathscr{N}_{\alpha,k}$ by sending $e_{\alpha+1,j}$ to $te_{\alpha,j}$. We will set $\lambda=\exp(2\pi i\alpha)$.

We will denote by $f^+\mathscr{N}_{\alpha,k}$ the pull-back meromorphic bundle with connection on $X$. Given a holonomic $\mathscr{D}_X$-module $\mathscr{M}$, the $\mathscr{D}_X$-module $\mathscr{M}_{\alpha,k}:=\mathscr{M}\otimes_{\mathscr{O}_X}f^+\mathscr{N}_{\alpha,k}$ remains holonomic, and is equipped with a nilpotent endomorphism $\mathrm{N}$. The $\mathscr{D}_X$-modules $\mathscr{H}^k(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})$ are also holonomic and supported on $D$. Moreover, the inductive system $\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})$ is locally stationary. We denote its limit by $\psi_{f,\lambda}^{\mathrm{mod}}\mathscr{M}$ (since we have a canonical isomorphism between the object defined with $\alpha+1$ and $\alpha$). It is equipped with a nilpotent endomorphism induced by $\mathrm{N}$. Lastly, the inductive system $\mathscr{H}^1(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})$ has limit zero (and the other $\mathscr{H}^k$ are zero since $D$ is a divisor).

Locally on $X$, $\psi_{f,\lambda}^{\mathrm{mod}}\mathscr{M}$ is zero except for a finite number of $\lambda\in\mathbb{C}^*$. We will set $\psi_f^{\mathrm{mod}}\mathscr{M}=\bigoplus_{\lambda\in\mathbb{C}^*}\psi_{f,\lambda}^{\mathrm{mod}}\mathscr{M}$. It is equipped with a semi-simple endomorphism induced by the multiplication by $\lambda$ on $\psi_{f,\lambda}^{\mathrm{mod}}\mathscr{M}$ and a unipotent one, induced by $\exp(2\pi i\mathrm{N})$.

**Corollary 14.19.** — *With the previous notation, we have in restriction to each compact set $K$ of $D$, an isomorphism (a priori depending on $K$)*

$$\mathrm{DR}\,\psi_{f,\lambda}^{\mathrm{mod}}\mathscr{M}_{|K}\simeq\varinjlim_k\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M}_{\alpha,k})_{|K}.$$

*Proof.* — The inductive limit above is meaningful, as it is defined in the category of complexes, but we will not try to give a meaning to the inductive limit of $\mathrm{DR}(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})$. On the other hand, the inductive limits of $\mathrm{DR}\big(\mathscr{H}^k(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})\big)$ are well defined and commute with taking cohomology of these complexes.

The distinguished triangle

$$\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})\longrightarrow i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k}\longrightarrow\mathscr{H}^1(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})[-1]\xrightarrow{+1}$$

induces a distinguished triangle in $D^b(\mathbb{C}_X)$:

$$\mathrm{DR}\,\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})\longrightarrow\mathrm{DR}\,i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k}\longrightarrow\mathrm{DR}\,\mathscr{H}^1(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})[-1]\xrightarrow{+1}$$

and, together with the isomorphism $\mathrm{DR}\,i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k}\simeq\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M}_{\alpha,k})$, we get a morphism

$$\mathrm{DR}\,\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})\longrightarrow\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M}_{\alpha,k}),$$

and thus a morphism

$$\mathrm{DR}\,\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})\longrightarrow\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M}_{\alpha,\infty})=\varinjlim_k\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M}_{\alpha,k}).$$

When restricted to the compact set $K\subset D$, this morphism is an isomorphism for $k$ big enough: indeed, because the inductive limit of $\mathscr{H}^1(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})$ is zero, each cohomology sheaf of $\varinjlim_k\mathrm{DR}\,\mathscr{H}^1(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})$ is zero; one uses then that $\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k})\to\mathscr{H}^0(i_{D,+}i_D^\dagger\mathscr{M}_{\alpha,k+1})$ is an isomorphism when restricted to $K$, when $p$ is big enough. ☐

**14.f. Comparison.** — We now go back to the setting of §14.c. Let $\mathscr{M}$ be a *good* meromorphic bundle with flat connection on $X$ with poles along a divisor with normal crossings $D$, with associated stratified $\mathfrak{I}$-covering denoted by $\widetilde{\Sigma}$. Let $(\mathscr{F},\mathscr{F}_\bullet)$ be the associated Stokes-filtered local system on $\widetilde{X}(D_{j\in J})$, and let $(\mathscr{L},\mathscr{L}_\bullet)$ denote its restriction to $\partial\widetilde{X}(D_{j\in J})$. Let us also assume that $\widetilde{\Sigma}\cup\{0\}$ is good (i.e., each local section of $\widetilde{\Sigma}$ is purely monomial). Then the complex $\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$ has cohomology in degree 0 at most, and $\mathscr{L}_{\leqslant0}:=\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$ is a subsheaf of $\mathscr{L}=\mathscr{H}^0\widetilde{\mathrm{DR}\mathscr{M}}$ (see Corollary 11.22).

Let us now consider the rapid decay de Rham complex of $\mathscr{M}$. This is the complex defined similarly to $\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$ (see §8.c) by replacing the coefficient sheaf $\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$ with the sheaf $\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$ as defined in Remark 8.4. This complex has already been considered in dimension one for the full Riemann-Hilbert correspondence (see Theorem 5.3), but not in dimension bigger than one because the grading has not been analyzed. Our purpose (Corollary 14.22 and Remark 14.23(1)) it to compare this grading process to the nearby cycle functor $\widetilde{\psi}_f^0$ applied to the Stokes-filtered local system attached to $\mathscr{M}$, when $f:X\to\mathbb{C}$ has zero set equal to $D$. We will first prove the natural analogue of Theorem 5.3.

***Proposition 14.20.*** — *For any germ $\mathscr{M}$ along $D$ of good meromorphic connection such that $\widetilde{\Sigma}\cup\{0\}$ is also good, the complexes $\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$, $\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}$ and $\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}$ have cohomology in degree $0$ at most. The natural morphisms $\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}\to\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}\to\widetilde{\mathrm{DR}}\mathscr{M}$ induce inclusions $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}\hookrightarrow\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}\hookrightarrow\mathscr{H}^0\widetilde{\mathrm{DR}}\mathscr{M}$, and $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}$ is equal to $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}/\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}$.*

*Proof.* — The question is local. Assume first that $\mathscr{M}$ has a good decomposition. That $\mathscr{H}^k(\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M})=0$ for $k\neq 0$ is Corollary 11.22. For $\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}$, the similar assertion is proved with the same arguments (see [**72**, §7] when $\dim X=2$). The remaining part of the proposition follows easily.

In order to treat the ramified case, one uses the same argument as in the proof of Lemma 10.8. ☐

***Proposition 14.21.*** — *With the same assumptions as in Proposition 14.20, the subsheaf $(q_f^{-1}\mathscr{L}_{\leqslant})_{<0}$ of $\mathscr{L}$ is identified with $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}$, so that there is a natural isomorphism*

$$\mathscr{H}^0\,\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}\xrightarrow{\ \sim\ }\widetilde{\psi}_f^0(\mathscr{F},\mathscr{F}_\bullet).$$

*Proof.* — Since both sheaves $(q_f^{-1}\mathscr{L}_{\leqslant})_{<0}$ and $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}$ are subsheaves of $\mathscr{L}_{\leqslant 0}=\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{M}$, the comparison can be done locally on $\partial\widetilde{X}$. Working with $\mathscr{A}_{\widetilde{X}}\otimes\mathscr{M}$ and using the Hukuhara-Turrittin-Majima theorem 11.20, we then reduce to proving the assertion for $\mathscr{M}=\mathscr{E}^\eta$, where $\eta$ is a purely monomial local section of $\mathcal{I}$. It is not difficult to show that, in such a case,

$$\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{E}^\eta=\begin{cases}0 & \text{if $\eta$ does not have a pole}\\ & \text{along each components of $D$,}\\ \mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}\mathscr{E}^\eta & \text{otherwise.}\end{cases}$$

According to Lemma 14.6, $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{E}^\eta$ coincides with the corresponding $(q_f^{-1}\mathscr{L}_{\leqslant})_{<0}$. ☐

***Corollary 14.22.*** — *With the same assumptions as in Proposition 14.20, we have for each $\lambda\in\mathbb{C}^*$ and on each compact set $K\subset D$ a functorial isomorphism of objects of $D^b_c(D)$ equipped with an automorphism*

$$(\psi^0_{f,\lambda}(\mathscr{F},\mathscr{F}_\bullet),T)_{|K}\xrightarrow{\ \sim\ }(\mathrm{DR}\,\psi^{\mathrm{mod}}_{f,\lambda}\mathscr{M},T)_{|K}.$$

*Proof.* — Since $\mathscr{M}$ is $\mathscr{O}_X(*D)$-locally free, we can apply [**73**, Cor. II.1.1.19, p. 45] to compute $\boldsymbol{R}\varpi_*\,\mathrm{DR}^{\mathrm{rd}\,D}\mathscr{M}$, and conclude that $\boldsymbol{R}\varpi_*\,\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}$ is isomorphic to $\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M})$. Let us now replace $\mathscr{M}$ with $\mathscr{M}_{\alpha,k}$.

We denote by $L_{\lambda^{-1},k}$ the local system

$$\mathrm{Ker}\left[\partial_t:\mathscr{A}^{\mathrm{mod}\,0}_{\partial\widetilde{S}}\otimes\mathscr{N}_{\alpha,k}\longrightarrow\mathscr{A}^{\mathrm{mod}\,0}_{\partial\widetilde{S}}\otimes\mathscr{N}_{\alpha,k}\right]$$

on $\partial\widetilde{S}$. This is a rank $k+1$ local system with monodromy $\lambda^{-1}\,\mathrm{Id}+\mathrm{N}_{k+1}$, where $\mathrm{N}_{k+1}$ is a Jordan block of size $k+1$.

Similarly, we have

$$\widetilde{f}^{-1}L_{\lambda^{-1},k}=\mathscr{H}^0\,\mathrm{DR}^{\mathrm{mod}\,D}(f^+\mathscr{N}_{\alpha,k}),$$

and the $\mathscr{H}^j$ vanish for $j>0$, according to Proposition 14.20. Moreover, it is immediate to check that $\mathscr{H}^0\,\mathrm{DR}^{\mathrm{rd}\,D}(f^+\mathscr{N}_{\alpha,k})=0$, and Proposition 14.20 also implies

$$\widetilde{f}^{-1}L_{\lambda^{-1},k}=\mathscr{H}^0\,\mathrm{DR}^{\mathrm{gr}\,D}(f^+\mathscr{N}_{\alpha,k}).$$

We conclude that there is a natural morphism of complexes

$$\widetilde{f}^{-1}L_{\lambda^{-1},k}\otimes_{\mathbb{C}}\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}\longrightarrow\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}_{\alpha,k},$$

and one checks by a local computation on $\partial\widetilde{X}$, by using the Hukuhara-Turrittin-Majima theorem 11.20, that this morphism is a quasi-isomorphism. Taking its inductive limit we get a quasi-isomorphism

$$\widetilde{f}^{-1}L_{\lambda^{-1},\infty}\otimes_{\mathbb{C}}\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}\longrightarrow\varinjlim_k\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}_{\alpha,k},$$

We therefore get an isomorphism

$$\boldsymbol{R}\varpi_*\big(\widetilde{f}^{-1}L_{\lambda^{-1},\infty}\otimes_{\mathbb{C}}\mathrm{DR}^{\mathrm{gr}\,D}\mathscr{M}\big)\longrightarrow\varinjlim_k\mathrm{DR}(\mathscr{O}_{\widehat{D}}\otimes\mathscr{M}_{\alpha,k}).$$

According to the projection formula, the left-hand term is $\psi^0_{f,\lambda}(\mathscr{F},\mathscr{F}_\bullet)$ and the right-hand term, restricted to the compact set $K$, is identified with $\mathrm{DR}\,\psi^{\mathrm{mod}}_{f,\lambda}\mathscr{M}$. The comparison of monodromies is straightforward. ☐

***Remarks 14.23***

(1) If $\dim X=2$, we can define $(\psi^0_f(\mathscr{F},\mathscr{F}_\bullet),T)$ with the only assumption that $(\mathscr{F},\mathscr{F}_\bullet)$ is good by the procedure of §14.d. On the other hand, $\psi^{\mathrm{mod}}_f$ commutes with direct images of $\mathscr{D}$-modules (see e.g. [**59**]), and DR commutes with direct images by $\varepsilon$ (since $X'$ and $X$ have the same dimension, the shifts in the de Rham complexes cancel). Therefore, in such a case, we get a comparison isomorphism as in Corollary 14.22.

(2) One can extend in a straightforward way the comparison of Corollary 14.22 to the various $\psi^{\varphi}_f(\mathscr{F},\mathscr{F}_\bullet)$ provided $\varphi$ is good with respect to $\widetilde{\Sigma}$. If $\varphi$ is not ramified, the left-hand side is replaced with $(\mathrm{DR}\,\psi^{\mathrm{mod}}_f(\mathscr{M}\otimes\mathscr{E}^{-f^*\varphi}),T)$. Similarly, in dimension two, one can relax the goodness assumption on $\varphi$ with respect to $\widetilde{\Sigma}$.

# BIBLIOGRAPHY

[1] Y. André – Structure des connexions méromorphes formelles de plusieurs variables et semi-continuité de l'irrégularité, *Invent. Math.* **170** (2007), p. 147–198.

[2] D.G. Babbitt & V.S. Varadarajan – *Local moduli for meromorphic differential equations*, Astérisque, vol. 169-170, Société Mathématique de France, Paris, 1989.

[3] W. Balser, B.L.J. Braaksma, J.-P. Ramis & Y. Sibuya – Multisummability of formal power series solutions of linear ordinary differential equations, *Asymptotic Anal.* **5** (1991), p. 27–45.

[4] W. Balser, W.B. Jurkat & D.A. Lutz – Birkhoff invariants and Stokes' multipliers for meromorphic linear differential equations, *J. Math. Anal. Appl.* **71** (1979), no. 1, p. 48–94.

[5] D. Barlet – Fonctions de type trace, *Ann. Inst. Fourier (Grenoble)* **33** (1983), no. 2, p. 43–76.

[6] A.A. Beilinson, J.N. Bernstein & P. Deligne – Faisceaux pervers, in *Analyse et topologie sur les espaces singuliers*, Astérisque, vol. 100, Société Mathématique de France, 1982, p. 7–171.

[7] A.A. Beilinson, S. Bloch, P. Deligne & H. Esnault – Periods for irregular connections on curves, manuscript, 2005.

[8] J.-E. Björk – *Analytic $\mathscr{D}$-modules and applications*, Kluwer Academic Publisher, Dordrecht, 1993.

[9] S. Bloch & H. Esnault – Homology for irregular connections, *J. Théor. Nombres Bordeaux* **16** (2004), p. 357–371.

[10] A. Borel (ed.) – *Algebraic $\mathscr{D}$-modules*, Perspectives in Math., vol. 2, Boston, Academic Press, 1987.

[11] ———, Chap. VI-IX, in *Algebraic $\mathscr{D}$-modules*, Perspectives in Math., vol. 2, Academic Press, Boston, 1987, p. 207–352.

[12] L. Boutet de Monvel, A. Douady & J.-L. Verdier (eds.) – *Séminaire E.N.S. Mathématique et Physique*, Progress in Math., vol. 37, Birkhäuser, Basel, Boston, 1983.

[13] J.-L. Brylinski – Transformations canoniques, dualité projective, théorie de Lefschetz, transformation de Fourier et sommes trigonométriques, in *Géométrie et analyse microlocales*, Astérisque, vol. 140-141, Société Mathématique de France, 1986, p. 3–134.

[14] V.I. Danilov – The geometry of toric varieties, *Russian Math. Surveys* **33** (1978), p. 97–154.

[15] P. Deligne – Lettre à B. Malgrange du 19/4/1978, in *Singularités irrégulières, Correspondance et documents* [**19**], p. 25–26.

[16] ———, Lettre à B. Malgrange du 8/05/1984, in *Singularités irrégulières, Correspondance et documents* [**19**], p. 43–46.

[17] ———, Lettre à V.S. Varadarajan du 4/01/1986, in *Singularités irrégulières, Correspondance et documents* [**19**], p. 47–51.

[18] ———, Théorie de Hodge irrégulière (mars 1984 & août 2006), in *Singularités irrégulières, Correspondance et documents* [**19**], p. 109–114 & 115–128.

[19] P. Deligne, B. Malgrange & J.-P. Ramis – *Singularités irrégulières, Correspondance et documents*, Documents mathématiques, vol. 5, Société Mathématique de France, Paris, 2007.

[20] A. Dimca – *Sheaves in topology*, Universitext, Springer-Verlag, Berlin, New York, 2004.

[21] W. Fulton – *Introduction to toric varieties*, Ann. of Math. Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993.

[22] R. Gérard & Y. Sibuya – Étude de certains systèmes de Pfaff avec singularités, in *Équations différentielles dans le champ complexe*, Lect. Notes in Math., vol. 712, Springer-Verlag, 1979, p. 131–288.

[23] R. Godement – *Topologie algébrique et théorie des faisceaux*, Hermann, Paris, 1964.

[24] C. Hertling & C. Sabbah – Examples of non-commutative Hodge structures, *Journal de l'Institut mathématique de Jussieu* **10** (2011), no. 3, p. 635–674.

[25] M. Hien – Periods for irregular singular connections on surfaces, *Math. Ann.* **337** (2007), p. 631–669.

[26] ———, Periods for flat algebraic connections, *Invent. Math.* **178** (2009), no. 1, p. 1–22.

[27] M. Hien & C. Roucairol – Integral representations for solutions of exponential Gauss-Manin systems, *Bull. Soc. math. France* **136** (2008), no. 4, p. 505–532, arXiv: `0704.1739v1`.

[28] H. Hironaka – Subanalytic sets, in *Number theory, algebraic geometry and commutative algebra, in honor of Yasuo Akizuki*, Kinokuniya, Tokyo, 1973, p. 453–493.

[29] W.B. Jurkat – *Meromorphe Differentialgleichungen*, Lect. Notes in Math., vol. 637, Springer, Berlin, 1978.

[30] N. Kachi, K. Matsumoto & M. Mihara – The perfectness of the intersection pairings for twisted cohomology and homology groups with respect to rational 1-forms, *Kyushu J. Math.* **53** (1999), no. 1, p. 163–188.

[31] M. Kashiwara – B-functions and holonomic systems, *Invent. Math.* **38** (1976), p. 33–53.

[32] ———, On the holonomic systems of differential equations II, *Invent. Math.* **49** (1978), p. 121–135.

[33] ———, Regular holonomic $\mathscr{D}$-modules and distributions on complex manifolds, in *Complex analytic singularities*, Adv. Stud. Pure Math., vol. 8, North-Holland, Amsterdam, 1987, p. 199–206.

[34] ———, *D-modules and microlocal calculus*, Translations of Mathematical Monographs, vol. 217, American Mathematical Society, Providence, RI, 2003.

[35] M. Kashiwara & P. Schapira – *Sheaves on Manifolds*, Grundlehren der mathematischen Wissenschaften, vol. 292, Springer-Verlag, 1990.

[36] L. Katzarkov, M. Kontsevich & T. Pantev – Hodge theoretic aspects of mirror symmetry, in *From Hodge theory to integrability and TQFT: tt*-geometry* (R. Donagi & K. Wendland, eds.), Proc. Symposia in Pure Math., vol. 78, American Mathematical Society, Providence, RI, 2008, arXiv: `0806.0107`, p. 87–174.

[37] K. Kedlaya – Good formal structures for flat meromorphic connections, I: surfaces, *Duke Math. J.* **154** (2010), no. 2, p. 343–418, arXiv: `0811.0190`.

[38] ———, Good formal structures for flat meromorphic connections, II: excellent schemes, *J. Amer. Math. Soc.* **24** (2011), no. 1, p. 183–229, arXiv: `1001.0544`.

[39] M. Loday-Richaud & G. Pourcin – Théorèmes d'indice pour les opérateurs différentiels linéaires ordinaires, *Ann. Inst. Fourier (Grenoble)* **47** (1997), p. 1379–1424.

[40] Ph. Maisonobe & L. Narváez Macarro (eds.) – *Éléments de la théorie des systèmes différentiels géométriques, Cours du C.I.M.P.A., École d'été de Séville (1996)*, Séminaires & Congrès, vol. 8, Paris, Société Mathématique de France, 2004.

[41] Ph. Maisonobe & C. Sabbah – Aspects of the theory of $\mathscr{D}$-modules, Lecture Notes (Kaiserslautern 2002), version de juillet 2011, disponible à `http://math.polytechnique.fr/cmat/sabbah/livres.html`, 2011.

[42] H. Majima – *Asymptotic analysis for integrable connections with irregular singular points*, Lect. Notes in Math., vol. 1075, Springer-Verlag, 1984.

[43] B. Malgrange – *Lectures on the theory of functions of several complex variables*, T.I.F.R. Lectures on mathematics and physics, T.I.F.R., Bombay, 1958.

[44] ———, *Ideals of differentiable functions*, Oxford University Press, 1966.

[45] ———, Sur les points singuliers des équations différentielles, *Enseign. Math.* **20** (1974), p. 147–176.

[46] ———, La classification des connexions irrégulières à une variable, in *Séminaire E.N.S. Mathématique et Physique* [**12**], p. 381–399.

[47] ———, Sur les déformations isomonodromiques, I, II, in *Séminaire E.N.S. Mathématique et Physique* [**12**], p. 401–438.

[48] ———, Sur les images directes de $\mathscr{D}$-modules, *Manuscripta Math.* **50** (1985), p. 49–71.

[49] ———, *Équations différentielles à coefficients polynomiaux*, Progress in Math., vol. 96, Birkhäuser, Basel, Boston, 1991.

[50] ———, De Rham Complex and Direct Images of $\mathscr{D}$-Modules, in *Éléments de la théorie des systèmes différentiels (Images directes et constructibilité)*, Les cours du CIMPA, Travaux en cours, vol. 46, Hermann, Paris, 1993, p. 1–13.

[51] ———, Connexions méromorphes, in *Congrès Singularités (Lille, 1991)*, Cambridge University Press, 1994, p. 251–261.

[52] ———, Connexions méromorphes, II: le réseau canonique, *Invent. Math.* **124** (1996), p. 367–387.

[53] ———, On the extension of holonomic systems, in *Structure of solutions of differential equations (Katata-Kyoto, 1995)* (M. Morimoto & T. Kawai, eds.), World Sci. Publishing, 1996, p. 279–285.

[54] ———, On irregular holonomic $\mathscr{D}$-modules, in *Éléments de la théorie des systèmes différentiels géométriques*, Séminaires & Congrès, vol. 8, Société Mathématique de France, Paris, 2004, p. 391–410.

[55] B. Malgrange & J.-P. Ramis – Fonctions multisommables, *Ann. Inst. Fourier (Grenoble)* **42** (1992), p. 353–368.

[56] Z. Mebkhout – *Le formalisme des six opérations de Grothendieck pour les $\mathscr{D}$-modules cohérents*, Travaux en cours, vol. 35, Hermann, Paris, 1989.

[57] ______, Le théorème de positivité, le théorème de comparaison et le théorème d'existence de Riemann, in *Éléments de la théorie des systèmes différentiels géométriques* [**40**], p. 165–310.

[58] Z. Mebkhout & Ph. Maisonobe – Le théorème de comparaison pour les cycles évanescents, in *Éléments de la théorie des systèmes différentiels géométriques* [**40**], p. 311–389.

[59] Z. Mebkhout & C. Sabbah – § III.4 $\mathscr{D}$-modules et cycles évanescents, in *Le formalisme des six opérations de Grothendieck pour les $\mathscr{D}$-modules cohérents*, Travaux en cours, vol. 35, Hermann, Paris, 1989, p. 201–239.

[60] T. Mochizuki – Good formal structure for meromorphic flat connections on smooth projective surfaces, in *Algebraic Analysis and Around*, Advanced Studies in Pure Math., vol. 54, Math. Soc. Japan, Tokyo, 2009, p. 223–253, arXiv: `0803.1346`.

[61] ______, On Deligne-Malgrange lattices, resolution of turning points and harmonic bundles, *Ann. Inst. Fourier (Grenoble)* **59** (2009), no. 7, p. 2819–2837.

[62] ______, Holonomic $\mathscr{D}$-modules with Betti structure, arXiv: `1001.2336`, 2010.

[63] ______, Asymptotic behaviour of variation of pure polarized TERP structures, *Publ. RIMS, Kyoto Univ.* **47** (2011), no. 2, p. 419–534, arXiv: `0811.1384`.

[64] ______, Stokes structure of good meromorphic flat bundle, *Journal de l'Institut mathématique de Jussieu* **10** (2011), no. 3, p. 675–712.

[65] ______, *Wild harmonic bundles and wild pure twistor D-modules*, Astérisque, Société Mathématique de France, Paris, 2011, to appear; arXiv: `0803.1344`.

[66] L. Narváez Macarro – The local duality theorem in $\mathscr{D}$-module theory, in *Éléments de la théorie des systèmes différentiels géométriques*, Séminaires & Congrès, vol. 8, Société Mathématique de France, Paris, 2004, p. 59–87.

[67] T. Oda – *Convex bodies and algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete Folge 3 Band 15, Springer-Verlag, 1988.

[68] J.-P. Ramis – Dévissage Gevrey, in *Journées Singulières de Dijon (Dijon, 1978)*, Astérisque, vol. 59, Société Mathématique de France, Paris, 1978, p. 173–204.

[69] C. Roucairol – Irregularity of an analogue of the Gauss-Manin systems, *Bull. Soc. math. France* **134** (2006), no. 2, p. 269–286.

[70] ______, The irregularity of the direct image of some $\mathscr{D}$-modules, *Publ. RIMS, Kyoto Univ.* **42** (2006), no. 4, p. 923–932.

[71] ______, Formal structure of direct image of holonomic $\mathscr{D}$-modules of exponential type, *Manuscripta Math.* **124** (2007), no. 3, p. 299–318.

[72] C. Sabbah – Équations différentielles à points singuliers irréguliers en dimension 2, *Ann. Inst. Fourier (Grenoble)* **43** (1993), p. 1619–1688.

[73] ______, *Équations différentielles à points singuliers irréguliers et phénomène de Stokes en dimension* 2, Astérisque, vol. 263, Société Mathématique de France, Paris, 2000.

[74] ______, *Déformations isomonodromiques et variétés de Frobenius*, Savoirs Actuels, CNRS Éditions & EDP Sciences, Paris, 2002, English Transl.: Universitext, Springer & EDP Sciences, 2007.

[75] ______, Développement asymptotique de distributions holonomes d'une variable complexe, arXiv: `math.CA/0611474`, 10 pages, 2006.

[76] ______, *Isomonodromic deformations and Frobenius manifolds*, Universitext, Springer & EDP Sciences, 2007, (in French: 2002).

[77] ______, An explicit stationary phase formula for the local formal Fourier-Laplace transform, in *Singularities, vol. 1*, Contemp. Math., American Mathematical Society, Providence, RI, 2008, p. 300–330, arXiv: `0706.3570`.

[78] ______, Fourier-Laplace transform of a variation of polarized complex Hodge structure, *J. reine angew. Math.* **621** (2008), p. 123–158, arXiv: `math.AG/0508551`.

[79] M. Saito – Induced $\mathscr{D}$-modules and differential complexes, *Bull. Soc. math. France* **117** (1989), p. 361–387.

[80] Y. Sibuya – Asymptotic solutions of a system of linear ordinary differential equations containing a parameter, *Funkcial. Ekvac.* **4** (1962), p. 83–113.

[81] ______, *Uniform simplification in a full neighborhood of a transition point*, Mem. Amer. Math. Soc., vol. 149, American Mathematical Society, Providence, RI, 1974.

[82] S.L. Svensson – Singular differential operators and distributions, *Israel J. Math.* **38** (1981), no. 1-2, p. 131–153.

[83] W. Wasow – *Asymptotic expansions for ordinary differential equations*, Interscience, New York, 1965.

# INDEX OF NOTATION

$\mathscr{A}_{\widetilde{X}}$, 63
$\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$, 63, 110
$\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$, 110
$\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$, 64
$\mathscr{A}_{\mathcal{I}\text{ét}}^{\mathrm{mod}\,D}$, $\mathscr{A}_{\mathcal{I}\text{ét}}^{\mathrm{mod}\,D}$, $\mathscr{A}_{\mathcal{I}\text{ét}}^{\mathrm{gr}\,D}$, 64
$\beta_{\leqslant}$, 10
$\beta_{\psi\leqslant\varphi}$, $\beta_{\psi<\varphi}$, 28
$C_X\mathscr{M}$ (Hermitian dual), 77
$\mathfrak{Db}_X$, 76
$\mathfrak{Db}_X^{\mathrm{mod}\,D}$, 77
$\mathrm{DR}\,\mathscr{M}$, 65
$\mathrm{DR}^{\mathrm{mod}\,0}\,\mathscr{M}$, $\mathrm{DR}^{\mathrm{mod}\,D}\,\mathscr{M}$, 66, 113
$\mathrm{DR}^{\mathrm{rd}\,0}\,\mathscr{M}$, $\mathrm{DR}^{\mathrm{rd}\,D}\,\mathscr{M}$, 66
$\widetilde{\mathrm{DR}}\mathscr{M}$, 65
$\mathrm{DR}_{\leqslant\varphi}$, $\mathrm{DR}_{<\varphi}$, $\mathrm{DR}_{\mathrm{gr}\,\varphi}$, 67
$\boldsymbol{D}$, $\boldsymbol{D}'$, 32
$\mathscr{E}^{\varphi}$, 66
$\Phi$ (exponential factors), 28
$\psi_{f,\lambda}$, 182
$\psi_f^{\varphi}(\mathscr{F},\mathscr{F}_{\bullet})$, 181
$\widetilde{\psi}_f^{\varphi}(\mathscr{F},\mathscr{F}_{\bullet})$, 181
$\psi_{f,\lambda}^{\mathrm{mod}}\,\mathscr{M}$, 188
$f_+$ (push-forward), 14
$f^+$ (pull-back), 17
$\mathcal{I}$, $\mathcal{I}^{\text{ét}}$, 7, 34, 122
$\mathscr{L}_{\bullet}$ (Stokes filtration), 28
$\leqslant_U$, $<_U$, $\leqslant_y$, $<_y$, 8
$\leqslant_\theta$, $<_\theta$, 10, 27, 123
$\mathcal{P}$, 9
$\mathscr{P}$, 61
$S^1_{\psi\leqslant\varphi}$, $S^1_{\psi<\varphi}$, 27
$\mathrm{St}(\psi,\varphi)$, 9, 123
$\Sigma$, $\Sigma_\alpha$, 22
$\widetilde{\Sigma}$, 20, 35, 126
$\mathrm{Tr}_{\leqslant}$, 15
$\mathrm{Tr}_{\leqslant q}$, 17
$\mathrm{Tr}_q$, 15
$\widetilde{X}(D)$, 9, 51
$\partial\widetilde{X}(D)$, 36, 51
$\widetilde{X}(D_{j\in J})$, 108

# INDEX

adjunction, 178

co-ℐ-filtration, 19
co-St-ℂ-constructible sheaf, 53

de Rham complex
  moderate –, 71, 113
  rapid-decay –, 71
distribution, 76
  with moderate growth, 77
duality, 39, 55, 57, 58, 73
  and Laplace, 97
  and topological Laplace, 99
  Poincaré, 29
  Poincaré-Verdier, 32

étale space, 7
exhaustive
  pre-ℐ-filtration, 16
exhaustivity property, 9

flatness, 110

good formal decomposition, 149, 150
good formal structure, 137
good lattice, 152
good meromorphic connection, 151
good set of exponential factors, 125
good stratified ℐ-covering, 127
graded-ℂ-constructible sheaf, 54

Hausdorff, 7
  stratified –, 21
Hermitian dual, 77
holomorphic functions
  with moderate growth ($\mathscr{A}_{\widetilde{X}}^{\mathrm{mod}\,D}$), 63, 110
  with rapid decay ($\mathscr{A}_{\widetilde{X}}^{\mathrm{rd}\,D}$), 64, 110
Hukuhara-Turrittin type, 76
Hukuhara-Turrittin-Majima theorem, 156
Hukuhara-Turrittin-Sibuya theorem, 139

ℐ-covering, 20
  – stratified, 21
ℐ-filtered local system, 20
ℐ-filtration, 18
  graded –, 18
irregularity number, 48

Laplace transform, 85
  topological –, 91
level structure, 42, 128
local system
  ℐ-filtered, 20
  Stokes-filtered, 27, 35, 38, 127
  Stokes-graded, 28

Malgrange-Sibuya theorem, 69, 144, 170
Mellin transformation, 79
moderate growth, 9, 42

Poincaré-Verdier duality, 32
pre-ℐ-filtration, 10
  exhaustive –, 16
  graded –, 17
  of a sheaf, 16
pull-back, 33
punctual good formal decomposition, 150
purely monomial, 123
push-forward, 14, 34

real blow-up
  along a divisor, 51, 107
  along a family of divisors, 108
Riemann-Hilbert (RH), 68, 71, 139, 163, 169

saturation with respect to an order, 15

Stokes data, 38
Stokes directions, 10
Stokes-filtered local system, 35, 127, 178
  good, 127
  non-ramified, 28
Stokes filtration, 35, 127
  graded, 28
  non-ramified, 28
  of level $\geqslant \ell$, 42
  of level $\geqslant \boldsymbol{\ell}$, 128
Stokes-perverse sheaf
  on $\underline{\widetilde{X}}$, 60
  on $\widetilde{X}$, 54
St-$\mathbb{C}$-constructible sheaf, 52
stratified $\mathcal{I}$-covering, 21, 152
strict morphism, 41
subanalytic, 123

tensor product, 11, 13
trace map, 15
twist, 11, 31